

- 

检索试验
- 

按国家、省（市）统计
- 

按疾病代码统计
- 

按试验实施单位统计
- 

按试验主办单位统计
- 

按经费或物资来源统计
- 

按招募研究对象情况统计
- 

按注册状态统计
- 

按干预措施统计
- 

按伦理委员会统计
- 

按研究类型统计

卒中后失眠认知行为治疗（CBTI-S)的疗效研究

下载XML文档

注册号: Registration number:	ChiCTR2400086678
最近更新日期: Date of Last Refreshed on:	2024-07-08
注册时间: Date of Registration:	2024-07-08
注册号状态:	补注册
Registration Status:	Retrospective registration
注册题目:	卒中后失眠认知行为治疗（CBTI-S)的疗效研究
Public title:	Study on the efficacy of cognitive behavioral therapy for insomnia after stroke (CBTI-S)
注册题目简写:	
English Acronym:	
研究课题的正式科学名称:	卒中后失眠认知行为治疗（CBTI-S)的疗效研究
Scientific title:	Study on the efficacy of cognitive behavioral therapy for insomnia after stroke (CBTI-S)
研究课题代号(代码): Study subject ID:	
在二级注册机构或其它机构的注册号: The registration number of the Partner Registry or other registrars:	
申请注册联系人: 李婷婷	研究负责人: 常翼
Applicant: Ting-ting LI	Study leader: Yichang
申请注册联系人电话: Applicant telephone:	研究负责人电话: Study leader's telephone: +86 411 83635963
申请注册联系人传真: Applicant Fax:	研究负责人传真: Study leader's fax:
申请注册联系人电子邮件: Applicant E-mail:	研究负责人电子邮件: Study leader's E-mail: changee99@gmail.com
申请单位网址(自愿提供): Applicant website(voluntary supply):	研究负责人网址(自愿提供): Study leader's website(voluntary supply):
申请注册联系人通讯地址: 辽宁省大连市西岗区滨海西路63号	研究负责人通讯地址: 大连市西岗区中山路222号
Applicant address: 63 Binhai West Road, Xigang District, Dalian City, Liaoning Province, China	Study leader's address: Zhongshan Road 222, Dalian
申请注册联系人邮政编码: Applicant postcode:	研究负责人邮政编码: Study leader's postcode:
申请人所在单位: 大连医科大学附属第一医院	
Applicant's institution: The First Affiliated Hospital of Dalian Medical University	
研究负责人所在单位: 大连医科大学附属第一医院	
Affiliation of the Leader: The First Affiliated Hospital of Dalian Medical University	
是否获伦理委员会批准: 是	
Approved by ethic committee: Yes	

伦理委员会批件文号：		伦理委员会批件附件：			
Approved No. of ethic committee: PJ-KS-KY-2024-283		Approved file of Ethical Committee: 查看附件View			
e:		ee:			
批准本研究的伦理委员会名称：		大连医科大学附属第一医院伦理委员会			
Name of the ethic committee:		Ethics Committee of The First Affiliated Hospital of Dalian Medical University			
伦理委员会批准日期：					
Date of approved by ethic committee: 2024-04-23					
伦理委员会联系人：		徐蕾			
Contact Name of the ethic committee: Xu Lei					
伦理委员会联系地址：		大连市西岗区中山路222号			
Contact Address of the ethic committee: Zhongshan Road 222, Dalian					
伦理委员会联系人电话：		伦理委员会联系人邮箱：			
Contact phone of the ethic committee: +86 411 83010706		Contact email of the ethic committee: dyyyirb@163.com			
研究实施负责（组长）单位：		大连医科大学附属第一医院			
Primary sponsor: The First Affiliated Hospital of Dalian Medical University					
研究实施负责（组长）单位地址：		大连市西岗区中山路222号			
Primary sponsor's address: Zhongshan Road 222, Dalian					
试验主办单位(项目批准或申办者)： Secondary sponsor:	国家：	中国	省(直辖市)：	辽宁	市(区县)：
	Country:	China	Province:	Liaoning	City:
	单位(医院)：	大连医科大学附属第一医院	具体地址：	大连市西岗区中山路222号	
	Institution of hospital:	The First Affiliated Hospital of Dalian Medical University	Address:	Zhongshan Road 222, Dalian	
.....					
经费或物资来源：		自选课题（自筹）			
Source(s) of funding: self-fund					
研究疾病：		卒中后病情稳定情况下仍存在失眠症状，且存在行为因素、条件性觉醒及睡眠卫生差的依据，每周至少有3天出现失眠症状，上述症状持续慢性（3个月以上）/短期（1个月以上）。 1）患者主诉对睡眠质量或睡眠长度不满意，症状包括：无法入睡、早醒、难以维持稳定的睡眠（夜间频繁觉醒、醒后再入睡困难）； 2）失眠障碍导致了明显的临床症状或损害了社会、职业、教育、行为等功能； 3）上述症状发生频率大于3次/周；			
Target disease:		insomnia symptoms at least 3 days per week, and the above symptoms are chronic (more than 3 months)/short-term (more than 1 month). 1) Patients complained of dissatisfaction with sleep quality or sleep length, with symptoms including: inability to fall asleep, early waking, difficulty in maintaining stable sleep (frequent waking at night, difficulty in falling asleep after waking up); 2) insomnia			
研究疾病代码：					
Target disease code:					
研究类型：		干预性研究			
Study type: Interventional study					
研究所处阶段：		其它			
Study phase: N/A					
研究设计：		随机平行对照			
Study design: Parallel					
研究目的：		主要目的：CBTI-S对卒中后失眠改善的疗效； 次要目的：CBTI-S对卒中后患者的焦虑抑郁情绪、生活自理能力、认知、乏力症状及卒中结果的影响； 探索性目的：失眠的改善可能对卒中的康复结果产生有利影响，观察其治疗效果的可持续性，为此类患者治疗提供依据。			
Objectives of Study:		Objective: To evaluate the effect of CBTI-S on insomnia after stroke. Secondary objectives: The effects of CBTI-S on anxiety and depression, self-care ability, cognition, fatigue symptoms and stroke outcomes in patients after stroke; Exploratory objective: The improvement of insomnia may have a favorable impact on the rehabilitation outcome of stroke, and to observe the sustainability of its therapeutic effect, so as to provide evidence for the treatment of such patients.			
药物成份或治疗方案详述：					
Description for medicine or protocol of treatment in detail:					
纳入标准：		1.年龄≥18岁； 2.卒中患者，发病≥1个月，且病情稳定无进展； 3.无意识障碍及严重认知障碍，无严重理解障碍； 4.口服镇静药物后失眠无改善且用药稳定； 5.符合美国《精神疾病诊断与统计手册》第V版（DSM-V）中关于失眠的诊断标准： 1）患者主诉对睡眠质量或睡眠长度不满意，症状包括：无法入睡、早醒、难以维持稳定的睡眠（夜间频繁觉醒、醒后再入睡困难）； 2）失眠障碍导致了明显的临床症状或损害了社会、职业、教育、行为等功能； 3）上述症状发生频率大于3次/周； 4）有充足的睡眠机会和良好的睡眠环境； 5）抱怨至少有一种白天的负面影响(如疲劳、功能受损或情绪紊乱)，可归因于睡眠不良； 6）上述症状不能被其他睡眠觉醒障碍所解释； 7）上述症状不是由药物或物质引起的生理效应（如药物滥用）； 8）失眠症状并非继发于躯体或精神疾病。卒中后病情稳定情况下仍存在失眠症状，且存在行为因素、条件性觉醒及睡眠卫生差的依据，每周至少有3天出现失眠症状，上述症状持续慢性（3个月以上）/短期（1个月以上）；			

Inclusion criteria	1. Age ≥18 years old; 2. Stroke patients, the onset ≥1 month, and the disease is stable without progression; 3. Unconscious disorder and severe cognitive disorder, no serious comprehension disorder; 4. Insomnia did not improve after oral sedation and medication was stable; 1.(5) Meet the diagnostic criteria for insomnia in the Diagnostic and Statistical Manual of Mental Disorders V (DSM-V) : 1) The patient's main complaint is not satisfied with sleep quality or sleep length, including: inability to fall asleep, early wake up, difficulty in maintaining stable sleep (frequent awakening at night, difficulty in falling asleep after waking up); 2) insomnia disorder leads to significant clinical symptoms or impairs social, occupational, educational, behavioral and other functions; 3) The occurrence frequency of the above symptoms is greater than 3 times/week; 4) Have sufficient sleep opportunities and a good sleep environment; 5) Complaining of at least one negative daytime effect (such as fatigue, impaired functioning, or mood disturbances) that can be attributed to poor sleep; 6) The above symptoms cannot be explained by other sleep-wake disorders; 7) The above symptoms are not physiological effects caused by drugs or substances (such as drug abuse); 8) Insomnia symptoms are not secondary to physical or mental illness. Symptoms of insomnia still exist in the stable condition after stroke, and there are behavioral factors, conditioned awakening and poor sleep hygiene evidence, insomnia symptoms at least 3 days per week, and the above symptoms are chronic (more than 3 months)/short-term (more than 1 month);				
	排除标准： 1.严重认知障碍（MMSE<10分）； 2.病情不稳定，卒中进展期，或无法控制的卒中外疾病，近期有手术无法完成治疗； 3.怀孕期及哺乳期妇女； 4.有意识障碍及未控制的精神障碍，既往存在精神疾病病史者； 5.其他睡眠障碍：异态睡眠、睡眠呼吸暂停综合征及不安退综合征；无法控制的疼痛影响睡眠的，既往存在躁狂、双向情感障碍、焦虑障碍、抑郁障碍、精神分裂症、创伤后应激障碍等影响睡眠的精神障碍障碍； 6.长期卧床患者； 7.目前从事跨时区、夜班或倒班工作。				
	Exclusion criteria: 1. Severe cognitive impairment (MMSE<10); 2. The condition is unstable, the stroke is in the advanced stage, or the disease outside stroke is uncontrollable, and the surgery cannot be completed recently; 3. Pregnant and lactating women; 4. Conscious disorder, uncontrolled mental disorder, and previous history of mental illness; 5. Other sleep disorders: parasomnia, sleep apnea syndrome and restless regression syndrome; 1.If uncontrollable pain affects sleep, there are previous mental disorders affecting sleep such as mania, bipolar disorder, anxiety disorder, depression disorder, schizophrenia, post-traumatic stress disorder; 6. Patients in bed for a long time; 7. Currently engaged in cross-time zone, night or shift work.				
	研究实施时间： Study execute time:		从 From 2024-06-01至 To 2025-12-01 招募观察对象时间： Recruiting time: 从 From 2024-06-01 至 To 2025-12-01		
	干预措施： Interventions:	组别：	研究组	样本量：	22
		Group：	research group	Sample size：	
		干预措施：	卒中后失眠认知行为治疗(CBTI-S)	干预措施代码：	
		Intervention：	Cognitive Behavioral Therapy for post-Stroke Insomnia (CBTI-S)	Intervention code：	
组别：		对照组	样本量：	22	
Group：		control group	Sample size：		
干预措施：		睡眠卫生教育（SHE）	干预措施代码：		
Intervention：		Sleep Hygiene Education (SHE)	Intervention code：		
研究实施地点： Countries of recruitment and research settings:	国家：	中国	省(直辖市)：	辽宁省(区县)：	
	Country：	China	Province：	Liaoning City：	
	单位(医院)：	大连医科大学附属第一医院	单位级别：	三级甲等	
	Institution hospital：	The First Affiliated Hospital of Dalian Medical University	Level of the institution：	Tertiary A	
测量指标： Outcomes:	指标中文名：	夜间睡眠情况	指标类型：	次要指标	
	Outcome：	Nocturnal sleep condition	Type：	Secondary indicator	
	测量时间点：	基线、干预结束及干预结束后3个月随访	测量方法：	针对夜间睡眠情况的测量，使用睡眠日记进行测量。	
	Measure time point of outcome：	At baseline, at the end of intervention, and at 3 months after intervention	Measure method：	For the measurement of sleep at night, a sleep diary was used.	
	指标中文名：	嗜睡程度	指标类型：	次要指标	
	Outcome：	Degree of sleepiness	Type：	Secondary indicator	
	测量时间点：	基线、干预结束及干预结束后3个月随访	测量方法：	针对嗜睡程度的测量，使用Epworth嗜睡量表进行测量	
	Measure time point of outcome：	At baseline, at the end of intervention, and at 3 months after intervention	Measure method：	The Epworth sleepiness Scale was used to measure the degree of sleepiness	
	指标中文名：	疲劳	指标类型：	次要指标	

	Outcome:	fatigue	Type:	Secondary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	针对疲劳程度的测量，使用疲劳严重程度量表（FSS)进行测量。
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	The fatigue Severity Scale (FSS) was used to measure the fatigue degree.
	指标中文名:	抑郁	指标类型:	次要指标
	Outcome:	depression	Type:	Secondary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	针对抑郁程度的测量，使用抑郁症筛查量表（PHQ-9）进行测量。
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	To measure the degree of depression, the Depression Screening Scale (PHQ-9) was used.
	指标中文名:	焦虑	指标类型:	次要指标
	Outcome:	anxiety	Type:	Secondary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	针对焦虑程度的测量，使用焦虑症筛查量表（GAD-7）进行测量。
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	To measure the degree of anxiety, the Anxiety Screening Scale (GAD-7) was used.
	指标中文名:	智能状况	指标类型:	次要指标
	Outcome:	Intelligence condition	Type:	Secondary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	通过简易智能精神状态检查量表（MMSE）进行测量。
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	It was measured by the Simple Intelligent Mental State Examination Scale (MMSE).
	指标中文名:	生活自理能力	指标类型:	次要指标
	Outcome:	self-care ability of daily living	Type:	Secondary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	通过日常生活能力量表（MBI）测量
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	It was measured by the Daily Living Ability Scale (MBI)
	指标中文名:	卒中严重程度	指标类型:	次要指标
	Outcome:	Stroke severity	Type:	Secondary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	通过美国国立卫生研究院卒中量表（NIHSS评分）测评。
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	Measured by the National Institutes of Health Stroke Scale (NIHSS score).
	指标中文名:	失眠严重程度	指标类型:	主要指标
	Outcome:	Severity of insomnia	Type:	Primary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	通过失眠严重程度指数量表(ISI)进行测量。
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	The insomnia Severity Scale (ISI) was used.
	指标中文名:	睡眠质量	指标类型:	主要指标
	Outcome:	sleep quality	Type:	Primary indicator
	测量时间点:	基线、干预结束及干预结束后3个月随访	测量方法:	针对睡眠质量的测量，使用匹茨堡睡眠质量指数量表进行测量。
	Measure time point of outcome:	At baseline, at the end of intervention, and at 3 months after intervention	Measure method:	For the measurement of sleep quality, the Pittsburgh Sleep Quality Index was used.
采集人体标本： Collecting sample(s) from participants:	标本中文名:	无	组织:	
	Sample Name:	NA	Tissue:	
	人体标本去向	其它	说明	
	Fate of sample:	Others	Note:	
征募研究对象情况： Recruiting status:		正在进行 Recruiting	年龄范围： Participant age:	最小 Min age 18 岁 years

性别：	男女均可	Gender：	Both
随机方法（请说明由何人用什么方法产生随机序列）：	电脑随机生成的随机数字表		
Randomization Procedure (please state who generates the random number sequence and by what method)：	A table of random numbers randomly generated by a computer		
是否公开试验完成后的统计结果：Calculated Results after the Study Completed public access:	不公开/Private		
盲法：	单盲		
Blinding：	Single blind study		
是否共享原始数据：IPD sharing	否No		
共享原始数据的方式（说明：请填写公开原始数据日期和方式，如采用网络平台，需填该网络平台名称和网址）：	临床试验公共管理平台 http://www.medresman.org.cn/login.aspx ，在试验结束6个月内上传试验数据。		
The way of sharing IPD“(include metadata and protocol, If use web-based public database, please provide the url)：	http://www.medresman.org.cn/login.aspx Upload the test data within 6 months after the end of the experiment.		
数据采集和管理（说明：数据采集和管理由两部分组成，一为病例记录表（Case Record Form, CRF），二为电子采集和管理系统(Electronic Data Capture, EDC)，如ResMan即为一种基于互联网的EDC：	1.研究数据使用专门的病例报告表（CRF），所有数据均遵循数据管理指南。 2.研究者有责任保证病例报告表上数据记录及时、准确、完整。病例报告表修改必须注明修改原因、修改时间并签名。 3.负责数据监查工作的成员对每份原始病历和 CRF/eCRF进行审核。 4.所有病例数据完成录入、负责数据监查工作的成员完成原始数据核查、数据管理人员完成核查、数据完成清理、主要研究者审核批准并完成签名后，由数据管理员进行数据锁定。数据锁定后，所有人员将不可对研究数据进行任何修改。		
Data collection and Management (A standard data collection and management system include a CRF and an electronic data capture)：	1. Special case report Form (CRF) was used for study data, and all data were in accordance with data management guidelines. 2. It is the responsibility of the researcher to ensure that the data recorded on the case report form is timely, accurate and complete. Case report form modification must indicate the reason for modification, modification time and signature. 3. Each original medical record and CRF/eCRF are reviewed by the member responsible for data monitoring. 4. After the data of all cases are entered, the original data is checked by the member responsible for data monitoring, the data is checked by the data manager, the data is cleared, and the main investigator is reviewed and approved and signed, the data manager will lock the data. After the data is locked, all personnel will not be able to make any changes to the research data.		
数据与安全监察委员会：Data and Safety Monitoring Committee：	无/No		

返回列表



网站首页 | ChiCTR简介 | 检索入口 | 重要文件 | 注册指南 | 常见问题

中国临床试验注册中心 - 世界卫生组织国际临床试验注册平台一级注册机构
Copyright(c) (2005 - 2023) Chictr.org.cn.All rights reserved. 中国临床试验注册中心 版权所有
[蜀ICP备16010396号-9](#)

提示：推荐使用IE8.0以上版本 宽屏显示分辨率下使用系统