

A Development in the Dynamical Approach to General Relativity and Emergent Spacetime Physics: Background-Free Quantum Field Theory

Zachary Hall

December 5, 2025

“In my opinion, it would be unsatisfactory if there existed a world without matter. Rather, the $g_{\mu\nu}$ field should *be fully determined by matter and not able to exist without the matter.*”

Albert Einstein,
letter to de Sitter, 24 March 1917 ^a

^a Einstein (1998, 309)

“There is no such thing as an empty space, i.e. a space without field. Space-time does not claim existence on its own, but only as a structural quality of the field.”

Albert Einstein,
Relativity and the Problem of Space^a

^a Einstein (1992, 155)

“The right side [of $G_{\mu\nu} = \kappa T_{\mu\nu}$] is a formal condensation of all things whose comprehension in the sense of a field theory is still problematic. Not for a moment, of course, did I doubt that this formulation was merely a makeshift in order to give the general principle of relativity a preliminary closed form expression. For it was essentially *no more* than a theory of the gravitational field, which was isolated somewhat artificially from a total field of as yet unknown structure.”

Albert Einstein,
Autobiographical Notes^a

^a Einstein (1979, 71)

Introduction

In this paper, I offer a development friendly to Harvey Brown’s “dynamical approach” to understanding General Relativity (GR) and its background independence by attention to the two-point correlator of quantum field theory (QFT). In doing so, I will offer the beginnings of a *background-free* version of QFT that also proves supportive of the “emergent spacetime” project of Cao, Carroll, and Michalakis (Cao et al., 2017), (Carroll, 2016). In particular, I show that if we identify the metric and affine structure of spacetime with the singular structure of the two-point correlator and do not fix that structure *ab initio*, that structure is, in the case of a free Klein-Gordon field, automatically that of the EFE sourced by inhomogenieties in the smooth contributions to the two-point correlator at coincidence. This result is thus also a background-free recasting of the free Klein-Gordon field of QFT.

In Section 1, I exposit GR’s background independence, the dynamical approach to understanding it before registering limitations of some existing resources for developing the dynamical approach (semiclassical gravity and the spin-2 approach). In Section 3, I motivate the two-point correlator as an alternative object of study for Brown’s explanatory project. I then offer the result in Section 4. To conclude, I indicate directions for developing the result further and highlight its significance for the emergentist project.

1 GR's Background Independence

1. GR, through the Einstein Field Equations (EFE)

$$G_{\alpha\beta} + \Lambda g_{\alpha\beta} = \kappa T_{\alpha\beta} \quad (1)$$

with $G_{\alpha\beta} := R_{\alpha\beta} - \frac{1}{2}Rg_{\alpha\beta}$ the Einstein tensor, is background independent: some of the structure of spacetime is itself a *solution* to certain field equations, evolving together with the matter within spacetime.

“instead of thinking of space and time as a stage on which the drama of matter unfolds, we have to imagine some ultramodern theater in which the stage itself becomes one of the actors, changing as the drama unfolds” (Stachel, 1979, 32).




Figure 1: The spacetime structure of GR is dynamically determined by the matter-energy content of the spacetime. Image source: Gribben (2021)

2. Specifically, the *metric* g (“ $g_{\mu\nu}$ ”, measuring lengths of curve segments and angles between vectors) and the (curvature of the) *connection* ∇ (encoding “sameness of direction” along curves)—the tensor $R^\nabla_{\text{Riemann}}$ —are so determined (Misner et al., 1973, 244-277, 404-408).

- In GR, the the connection is required to be “compatible with” the metric:
 $\nabla_{\mathbf{u}}g = 0 \quad \forall \mathbf{u}$, so that lengths and angles are uniformly defined along any curve.

2 The Dynamical Explanation of GR's Background Independence

1. One way of explaining the dynamical dependence of the connection and metric on matter is the constructive or dynamical approach championed by Harvey Brown and Oliver Pooley:¹ gravity is a universal attractive force, thus the natural [read: "default"] ways things move can be represented by a connection with curvature; and the behaviors of real rods and clocks (temporal and spatial measures represented by the metric) is systematically deformed because they themselves are gravitationally influenced.
2. Einstein helps drive the point about the metric home:

...strictly speaking, measuring rods and clocks should be represented as solutions of the fundamental equations (objects consisting of moving atomic configurations), not, as it were, as theoretically self-sufficient entities (Einstein, 1979, 54-7).²

 - Real temporal and spatial measures are read off from real physical systems.
 - Thus successful cases of chronogeometric representation in GR must be explained by the behaviors of those systems.
 - Thus, fundamentally, the metric's dependence on matter and energy must be explained by the dependence of how real rods and clocks operate on their relations to matter and energy.
3. This explanation is *programmatic* and only completed insofar as one can manifestly recover the empirical successes of GR through theory that interfaces (i) quantum-theoretic description of real systems used as rods and clocks with (ii) gravity as a universally attractive force.
4. Efforts in this direction need not involve "solving quantum gravity."

¹ Brown (2005, Introduction, Ch. 3, Ch. 8-9), Brown and Pooley (2001), Brown and Pooley (2006).

² Maßstäbe und Uhren müssten eigentlich als Lösungen der Grundgleichungen (Gegenstände aus bewegten atomistischen Gebilden) dargestellt werden, nicht als gewissermaßen theoretisch selbständige Wesen.

5. Existing such efforts have defects by the lights of Brown's explanatory program.

(a) Semiclassical gravity feeds the expectation value of the stress energy tensor $\langle T_{\mu\nu} \rangle$ into Einstein's Field Equations (EFE) to determine a metric and connection that is then used as a curved background for the equations of standard QFT generalized to curved spaces, which then determines $\langle T_{\mu\nu} \rangle$ more precisely in that region, and so on—it is a mixed, iterative application of GR and QFT (Birrell and Davies, 1982), (Wald, 1994), (Kuo and Ford, 1993).

- This *works* well in making sense of systems where both quantum theoretic effects and gravitational effects are relevant. But it is not an *explanation* from gravity as a universal attractive force at the quantum level to the empirical successes of GR. It takes the EFE as a *given*, then finds some mathematically tractable way of iteratively relating the structures of QFT and the EFE, rather than recovering the EFE as an effective description of gravitation-involving quantum-theoretic systems.

(b) The spin-2 approach theorizes a universal attractive coupling in the framework of Quantum Field Theory (QFT) via a [self-interacting] “massless spin-2 field” h mediating the gravitational force by coupling to the stress energy tensor T , and gives the EFE under certain natural assumptions. Schematically, one starts with a flat-space QFT (metric = η) with Lagrangian $\mathcal{L}_{\text{total}}$, then adds a term $\mathcal{L}_{\text{total}} \rightarrow \mathcal{L}_{\text{total}} + h_{\mu\nu}T^{\mu\nu}$, and consistency arguments show h must self-interact so that $S_{\text{total}} \rightarrow S_{\text{total}} + S_{\text{EH}}$, from which the EFE follow by the variational principle, $G_{\mu\nu} = \kappa T^{\mu\nu}$ with $G_{\mu\nu}$ computed from g in the standard way (the connection is the Levi-Civita connection) (Gupta, 1954), (Deser, 1970).

- This plays well with QFT formally, but crucially *does not show that the microphysical dynamics of quantum fields reflects the curved metric given by h* . More specifically, none of the dynamics for the matter fields ϕ_i are required by the self-consistency argument to use a connection compatible with the variable metric, $\nabla^{R(g)}$.³ The dynamical approach requires that the microphysical dynamics of rods and clocks *give rise to* the metric g of the EFE, but that those microphysical dynamics even track g at all can only be stipulated by fiat after spin-2 delivers its result, thus the spin-2 approach does not deliver for the dynamical explanatory strategy, *pace* Salimkhani (2020).

6. Thus, neither semiclassical gravity nor spin-2 has the resources to execute the dynamical explanatory strategy. My aim is to see if we can get some traction here by attention to some formally geometric structure in the microphysical details of QFT (the two-point correlator and its singular structure).

³ The matter Lagrangian \mathcal{L} begins formulated with the flat connection ∇^η , and the self-consistency argument simply gives a non-matter Lagrangian from which $G_{\mu\nu} = \kappa T_{\mu\nu}$ is derived—the consistency condition is that the total stress energy $T_{\mu\nu}^{\text{total}} = T_{\mu\nu}^{\text{matter}} + T_{\mu\nu}^{\text{gravitational}}(h)$ is conserved *with respect to the flat connection*, $\nabla_\mu^\eta T_{\text{total}}^{\mu\nu} = 0$. This leaves the matter Lagrangian completely untouched, and thus all microphysical quantum dynamics untouched. One could motivate changing the connection in $\mathcal{L}_{\text{matter}}$ to $\nabla^{R(g)}$ because otherwise we do not have $\nabla_\mu^{R(g)} T_{\text{matter}}^{\mu\nu} = 0$, which is implied by the EFE which the consistency argument has forced. However, $\nabla_\mu^{R(g)} T^{\mu\nu} = 0$ is forced by the EFE *only by utilizing a geometric identity involving $G_{\mu\nu}$* , namely $\nabla_\mu^{R(g)} T^{\mu\nu} = 0$, but the whole point of the spin-2 approach is not to treat h/g as a metric in the first place, but to show that matter only “sees” an effective metric g from the universal dynamics of the self-interacting h . Nor does $\nabla_\mu^\eta (T_{\mu\nu}^{\text{matter}} + T_{\mu\nu}^{\text{gravitational}}(h)) = 0$ analytically imply $\nabla_\mu^{R(g)} T_{\text{matter}}^{\mu\nu} = 0$ on its own, as the only way to get an expression for $T_{\mu\nu}^{\text{gravitational}}(h)$ in the first place is, apparently, by *assuming* $\nabla_\mu^{R(g)} T_{\text{matter}}^{\mu\nu} = 0$ (Padmanabhan, 2008), (Linnemann et al., 2026) (forthcoming). Therefore, the only way to motivate the transition from $\nabla_\mu^\eta T_{\text{total}}^{\mu\nu} = 0$, which the self-consistency argument implies, to $\nabla_\mu^{R(g)} T_{\text{matter}}^{\mu\nu} = 0$, which it does not, is to illicitly treat g as the geometry of the spacetime substantively, such that fluxes of stress-energy are divergence-free with respect to its geometry; but this is precisely not to give a microphysical, quantum field theoretic explanation of the dynamics of GR, but to postulate that those dynamics are specified by the geometry of GR.

3 The Two-Point Correlator, Hadamard States

1. A natural object of study for universal, geometrically representable microphysical dynamics is the (time-symmetric) two-point correlator $W(x, y) = \langle S | \phi(x) \phi(y) | S \rangle$, where $\phi(x)$ is a quantum field of certain kind and S is either the vacuum ($S = 0$) or an excitation thereof for that field-kind. $W(x, x')$ represents how correlated a field is with itself between two points x and x' in the vacuum state or in an excited state (Wald, 1994, 43-45, 74-75).
2. One motivation is that it is the simplest object that represents how quantum fields exist over arbitrarily small spacetime regions, as they are ill-defined pointwise (Wald, 1994, 35, 45), (Haag, 1996, 45). (They are operator-valued distributions rather than classical fields.)
3. Another is that the mathematical structure of $W(x, y)$ for all physically admissible states of all fields (i) is universal and (ii) reflects metric and causal relations (Wald, 1977, 14-16), (Fulling et al., 1981, 243-244), (Birrell and Davies, 1982, 23), (Wald, 1994, 91-93); (Wald, 1994, 91), (Fulling, 1989, 176).
4. Roughly, $W(x, y)$ is the following form:

$$W(x, y) \simeq \frac{U(x, y)}{\sigma(x, y)} + S(x, y),$$

where $\frac{U(x, y)}{\sigma(x, y)}$ is the universal, geometric-dependent part, with Synge's world function $\sigma(x, y)$ equal to half the geodesic distance squared, and $S(x, y)$ is the state-dependent part that encodes difference from the geometric baseline structure. $\frac{U(x, y)}{\sigma(x, y)}$ is *singular*—it diverges as $y \rightarrow x$. $S(x, y)$ is *smooth*: it never diverges, and is distributionally differentiable at all orders in both arguments, $S(x, y) \in C^\infty(M \times M)$ (Friedlander and Joshi, 1998, Ch. 1-2). The universal part is the generalization of the singular structure of the quantum vacuum of flat-space QFT to QFT in curved spacetime (QTFCS) (Wald, 1994, 94). Thus:

$$W(x, y) \simeq \overbrace{\frac{U(x, y)}{\sigma(x, y)}}^{\text{singular, reflects short distance and causal structure (universal)}} + \overbrace{S(x, y)}^{\text{smooth (state dependent)}}$$

- The singular part reflects short-distance structure in that as geodesic distance on timelike or spacelike curves approaches 0, the field becomes infinitely self-correlated. Intuitively: this reflects infinitely energetic vacuum fluctuations at short distance (Wald, 1994, 94). Mathematically, this reflects the distributional character of quantum fields (Wald, 1994, 35, 45), (Haag, 1996, 45).

- The singular part reflects causal structure because along null geodesics (along which causal signals propagate), $\sigma(x, y) = 0$. This reflects that the singularity of $W(x, y) = \langle S | \phi(x)\phi(y) | S \rangle$ at coincidence ($x = y$) must propagate along the bicharacteristic flow of the kinetic differential operator \mathcal{P} when that field propagates in accordance with $\mathcal{P}\phi = 0$, i.e., precisely along null cones. This was proved by [Duistermaat and Hörmander \(1972, 194-197\)](#) as the “propagation of singularities theorem.” The key conceptual upshot for our purposes: this result allows both the time-like and spacelike distance structure and the causal-kinetic structure of fields to be represented in the universal part of the two-point function $W(x, y)$.
- In summary: that $W(x, y)$ of any physically well-behaved quantum field ϕ is singular at coincidence ($x = y$) mathematically encodes the short-distance structure of such fields; that it is singular away from coincidence ($x \neq y$) precisely along null cones (geodesic distance equals 0) reflects those fields’ being internally causally connected by null cones.

5. More precisely, for all physically admissible states,

$$W(x, y) = \frac{U(x, y)}{\sigma(x, y)} + V(x, y) \ln \sigma + S(x, y),$$

where the $V(x, y) \ln \sigma$ term is included to cancel unwanted terms resulting from applying the kinetic differential operator \mathcal{P} to $\frac{U(x, y)}{\sigma(x, y)}$ when applying it to $W(x, y)$ ([Zelditch, 2017, 125](#))—this ensures that $\mathcal{P}_x W(x, y) = \mathcal{P}_y W(x, y) = 0$ as it must when $\mathcal{P}\phi = 0$.⁴ $U(x, y)$ and $V(x, y)$ are coefficients that obey transport equations such that those unwanted terms cancel at arbitrary $x \neq y$ ([Poisson et al., 2011, 71](#)).

6. It is standard to collect the U and V terms into one geometry-representing, singular bidistribution, $H(x, y) := \frac{U(x, y)}{\sigma(x, y)} + V(x, y) \ln \sigma$ ([Wald, 1994](#)). Thus we have, for all physically admissible states,

$$W(x, y) = \overbrace{H(x, y)}^{\text{singular, reflects short distance and causal structure (universal)}} + \overbrace{S(x, y)}^{\text{smooth (state dependent)}}.$$

Such a state is called a *Hadamard state*.

7. Only Hadamard states are thought to be physically admissible because only they allow identification of finite, local, state-dependent quantities ([Wald, 1977](#)), ([Kay et al., 1997](#)), ([Fewster and Verch, 2002](#)), ([Fewster and Verch, 2013](#)).

⁴ If $\mathcal{P}\phi = 0$, then $\mathcal{P}_x W(x, y) = \langle S | \mathcal{P}_x \phi(x) \phi(y) | S \rangle = \langle S | 0 \phi(y) | S \rangle = 0$ and $\mathcal{P}_y W(x, y) = \langle S | \phi(x) \mathcal{P}_y \phi(y) | S \rangle = \langle S | \phi(x) 0 | S \rangle = 0$.

4 The Free Bisolution of the Two-Point Correlator as a Fixed Point Law

1. In QFTCS, the metric g is taken as given, and then properties of e.g. the two-point correlator $W(x, y)$ is studied, with $\sigma(x, y)$ computed with g . See e.g. [Wald \(1977\)](#).
2. This allows representation of the microphysical dynamics of quantum fields as obeying the geometry specified by the EFE, but leaving it as it stands undermines the dynamical explanatory strategy: there is no explanation of why microphysical dynamics give rise to the macroscopic effective spacetime geometry successfully represented by the EFE, but a stipulation that they do.
3. Is there a way to affect this strategy by attention to the geometric structure of the two-point correlator $W(x, y)$ *itself*, without supplying the metric from elsewhere?
4. Consider, the universal, singular part of $W(x, y)$, i.e. $H(x, y)$, already encodes geometric information and, by the propagation of singularities theorem, causal information. For this reason, given that such an $H(x, y)$ exists with a suitable real function $\sigma(x, y)$, the metric can be defined from $\sigma(x, y)$ itself: $g_{\alpha\beta} = \lim_{y \rightarrow x} \nabla_\alpha \nabla_\beta \sigma(x, y)$ with ∇ acting on the x -slot ([Poisson et al., 2011](#), 37). Perhaps we can think of it *as* the spacetime geometry, rather than merely reflecting it. This suggestion is consonant with Carroll et. al.'s program that seeks to define spacetime structure from entanglement entropy ([Cao et al., 2017](#)): perhaps, just as [Carroll \(2016\)](#) suggests, we can think of *spacetime geometry as in the quantum state*—specifically in how it is self-correlated—rather than the quantum state as in spacetime. Carroll's result showing EFE-like dependence on the curvature defined from the entanglement entropy under perturbations of the system further encourages looking in this direction, given the close connection of the two-point correlator to entanglement entropy ([Lin et al., 2024](#)).
5. Now we must ask: are there any principles holding of $W(x, y)$ itself that put constraints on the geometry encoded by $\sigma(x, y)$? If not, it is difficult to see how the dynamical explanatory strategy can succeed.
6. Let us first consider the structure of the two-point correlator for a free field, i.e. $\mathcal{P}\phi = 0$. In this case, beyond the Hadmard-form of $W(x, y)$, we have that $W(x, y)$ is a bisolution of \mathcal{P} : $\mathcal{P}_x W(x, y) = \mathcal{P}_y W(x, y) = 0$.⁵ I will henceforth abbreviate both conditions by $\mathcal{P}_g W(x, y) = 0$.
 - The subscript g marks that the kinetic operator \mathcal{P}_g is defined with the metric g : it always includes a (Levi-Cevita) connection ∇ .
 - [Bär and Ginoux \(2012, Ch. 1.5\)](#) gives the general form of all such operators. In the simplest case—the massless scalar field— $\mathcal{P}_g = \square_g = g^{\alpha\beta} \nabla_\alpha \nabla_\beta$ ([Wald, 1994](#), 54). This is a hyperbolic operator with wavelike solutions to $\mathcal{P}\phi = 0$ travelling at the speed of light c —in flat space, $\mathcal{P} = -c^2 \partial_t^2 + \sum_i \partial_i^2$.

⁵ See ([Birrell and Davies, 1982](#), 10, 20-21), ([Wald, 1994](#), 91), and footnote 4.

7. [Bär et al. \(2007, 34-36\)](#) prove that the Levi-Civita connection—the unique torsion-free metric compatible connection—is the unique connection for all normally hyperbolic operators; together with [Duistermaat and Hörmander \(1972, 194-197\)](#)’s Propagation of Singularities theorem, this implies that the Levi-Civita connection is required for the definition of Hadamard states—otherwise, the singularities do not propagate along the null cones, and $H(x, y)$ singular away from coincidence ($x \neq y$) cannot encode the causal structure of the field ϕ . I will thus henceforth require the connection ∇ to be Levi-Civita.

8. When $\mathcal{P}_g\phi = 0$,

$$\mathcal{P}_g W(x, y) = \mathcal{P}_g (H(x, y) + S(x, y)) = 0.$$

$H(x, y)$ encodes the spacetime geometry/metric (both distance structure and causal relatedness), and $S(x, y)$ encodes the excitation content of the quantum field. Thus we have

$$\overbrace{\mathcal{P}_g H(x, y)}^{\text{geometry}} + \overbrace{\mathcal{P}_g S(x, y)}^{\text{matter}} = 0$$

when $\mathcal{P}_g\phi = 0$.

9. The equation

$$\overbrace{\mathcal{P}_g H(x, y)}^{\text{geometry}} + \overbrace{\mathcal{P}_g S(x, y)}^{\text{matter}} = 0 \tag{2}$$

is valid in QFTCS. Again, QFTCS places no *a priori* restrictions on g — σ —whatsoever; the geometry is given from without. Thus QFTCS treats g as input, and Eq. (2) as a linear differential equation governing the propagation of the smooth part S given that geometry. As far as QFTCS is concerned, g is specified freely, and Eq. (2) specifies the admissible S given g .

10. One might think the free field law $\mathcal{P}_g\phi = 0$ is thus insufficient for the dynamical explanatory strategy: we will have to include interactions ($\mathcal{P}_g\phi \neq 0$), ultimately gravitational interactions, to constrain the universal geometry encoded by σ . But consider: if we treat g not as externally specifiable, but as a *variable to be determined* by Eq. (2)—signify g with this standing by $g[W]$ —then Eq. (2) takes on a different mathematical character: it is a non-linear differential equation of second order in g —the propagation operator in general (for arbitrary fields) contains $\nabla_\alpha \nabla_\beta$ to leading order—and ∇ and g are nonlinearly codetermining), restricting the admissible *pairs* (W, g) .

11. Thus the equation

$$\mathcal{P}_{g[W]} W = 0 \tag{3}$$

is equivalent to

$$\overbrace{\mathcal{P}_{g[W]} H(x, y)}^{\text{geometry}} + \overbrace{\mathcal{P}_{g[W]} S(x, y)}^{\text{matter}} = 0$$

and is therefore a nonlinear, second-order coupling of g to the excitation content S that is just as background-free as GR. It is a *fixed point law*: its solutions, thus all

possible ways excitation data and geometric data can interrelate, are consistent pairs $(W, g[W])$ where $\mathcal{P}_{g[W]}$ propagates the excitation data of W consistently with its total structure and W Hadamard form.

4.1 Local Geometric Consequences of $\mathcal{P}_{g[W]}W = 0$

III.2.1.i - The Balance Law

1. For simplicity in the following, I will restrict my attention to Klein-Gordon fields, where $\mathcal{P} = g^{\alpha\beta}\nabla_{\alpha\beta} - m^2$.
2. Some local geometric consequences of the coupling $\mathcal{P}_{g[W]}W = 0$ can be unpacked by defining a divergence-free correlator current from W . But

$$J_W^\alpha(x, y) := i(W^*(x, y)\nabla_x^\alpha W(x, y) - W(x, y)\nabla_x^\alpha W^*(x, y)).$$

is divergence-free *only away from coincidence* because of the singularity in $W(x, x)$. Because we want local geometric consequences, we need a current that is divergence-free and pointwise defined. Now, mathematically, for a given g , there are an infinite number of Hadamard bisolutions of \mathcal{P}_g , $\{B_c\}_g$, each differing from any other by a smooth bisolution (Lewandowski, 2022, §1.B). As a renormalization trick, we can use some reference $W_0 \neq W$, $W_0 \in \{B_c\}_{g[W]}$ to subtract from W to obtain a smooth bisolution on the geometry $g[W]$: $S_B(x, y) := W(x, y) - W_0(x, y)$. Now we have $\mathcal{P}S_B(x, y) = 0$, and we can define a correlation current

$$J_{S_B}^\alpha(x, y) := i(S_B^*(x, y)\nabla_x^\alpha S_B(x, y) - S_B(x, y)\nabla_x^\alpha S_B^*(x, y)).$$

3. $J_{S_B}^\alpha(x, y)$ is smooth everywhere, including at coincidence, and divergence free everywhere: it is mathematically the same as the divergence-free Klein-Gordon current with an extra y index floating around, $S(x, y)$ is a bisolution of the same \mathcal{P} that ϕ solves, and $\mathcal{P}\phi$ is the reason the Klein-Gordon current is divergence free.⁶
4. Therefore,

$$\nabla_\alpha J_S^\alpha(x) = 0 \tag{4}$$

with $J_{S_B}^\alpha(x) := \lim_{x \rightarrow y} J_{S_B}^\alpha(x, y)$. (As per Bär and Ginoux (2012, Ch. 1), when a bidistribution $A(x, y)$ is smooth, covariant differentiation and the limit-taking $x \rightarrow y$ commute—the only thing that would spoil this is singular behavior at the diagonal.)

5. Thus, by Gauss' law,

$$\int_V \nabla_\alpha J_{S_B}^\alpha = \oint_{\partial V} J_{S_B}^\alpha n_\alpha d\Sigma = 0.$$

6. Now for point p , pick a null direction k^α (let k^α be dimensionless) and arbitrarily small space-like 2-surface S_0 transverse to k^α , and consider the family of null geodesics emanating from S_0 . We may choose S_0 to be a small disc of radius ϵ . This defines a small “null pencil” with each null geodesic parametrizable by some λ , generating a small cross-sectional area $A_\epsilon(\lambda)$ defined by the metric g . Write $J_{S_B}^\alpha = \rho V^\alpha$ with V^α unit magnitude and dimensionless, and define $\rho_{||} := J_{S_B}^\alpha k_\alpha$ with k_α null and dimensionless. Thus ρ is the magnitude of $J_{S_B}^\alpha$ and $\rho_{||}$ is its magnitude projected along the null vector chosen.

⁶ See the proof by Gingrich (2004) and simply substitute $\partial_\mu \rightarrow \nabla_\mu$.

7. Now consider the pencil segment generated by λ ranging from λ_1 to λ_2 . We may make a closed four-volume by moving off the 3-volume by small magnitude Δz . By Gauss' law, the surface integral over the surface enclosing this 4-volume vanishes. This integral has four parts: the two caps of the pencil—call these pieces $-\Theta(\lambda_1)$ and $\Theta(\lambda_2)$ —, the null-oriented 2-surface boundary of the pencil segment, and the additional boundary generated by Δz . Thus we have

$$\oint_{\partial V} J_{S_B}^\alpha n_\alpha d\Sigma = \Theta(\lambda_2) - \Theta(\lambda_1) + \Theta_{\text{null boundary}} + \Theta_{\Delta z} = 0.$$

In the $\Delta z \rightarrow 0$ limit,

$$\lim_{\Delta z \rightarrow 0} \oint_{\partial V} J_{S_B}^\alpha n_\alpha d\Sigma = \Theta(\lambda_2) - \Theta(\lambda_1) + \Theta_{\text{null boundary}} = 0,$$

as the surface contributed by Δz becomes vanishingly small. But the null-oriented boundary has zero surface area, because each of its infinitesimal areas is spanned by a null vector k^α which has zero length according to the metric, $g_{\alpha\beta} k^\alpha k^\beta = 0$, and therefore there is zero flux of $J_{S_B}^\alpha$ through it. Therefore

$$\lim_{\Delta z \rightarrow 0} \oint_{\text{pencil segment}} J_{S_B}^\alpha n_\alpha d\Sigma = \Theta(\lambda_2) - \Theta(\lambda_1) = 0.$$

8. Now we may write $\Theta_\lambda = \bar{\rho}_{||}(\lambda) A_\epsilon(\lambda)$, and the above shows us that $\bar{\rho}_{||} A_\epsilon = \text{const.}$ along any null pencil. Therefore

$$\frac{1}{\bar{\rho}_{||} A_\epsilon} \frac{d}{d\lambda} (\bar{\rho}_{||} A_\epsilon) = \frac{1}{A_\epsilon} \frac{dA_\epsilon}{d\lambda} + \frac{1}{\bar{\rho}_{||}} \frac{d\bar{\rho}_{||}}{d\lambda} = 0.$$

9. $\frac{1}{A_\epsilon} \frac{dA_\epsilon}{d\lambda}$ has a standard name—the *expansion scalar* θ (more precisely θ_ϵ), so called because it encodes how cross-sectional areas of geodesic congruences expand (or contract) given the curvature of g (Wald, 1984, 216-222), (Poisson, 2004, 34). Thus we have

$$\theta_\epsilon + \frac{1}{\bar{\rho}_{||}} \frac{d\bar{\rho}_{||}}{d\lambda} = 0,$$

therefore

$$\frac{d\theta_\epsilon}{d\lambda} + \frac{d}{d\lambda} \left(\frac{1}{\bar{\rho}_{||}} \frac{d\bar{\rho}_{||}}{d\lambda} \right) = 0. \quad (5)$$

10. Eq. (5) is a balance law: it says that changes in how the cross-sectional area of a null congruence change along the parameter must be balanced by changes in normalized changes of the correlation density $\bar{\rho}_{||}$ along that parameter.
11. It is well known that, for a null pencil chosen with zero initial expansion or shear at point p —such a pencil always exists (Jacobson, 1995, 1261)—, in the limit $\epsilon \rightarrow 0$, $\frac{d\theta_\epsilon}{d\lambda} = -R_{\alpha\beta} k^\alpha k^\beta$, where $R_{\alpha\beta}$ is the Ricci curvature tensor for metric g (Wald, 1984, 216-222), (Poisson, 2004, 34, 40-51)—intuitively, $R_{\alpha\beta}$ controls focusing or defocusing

of geodesics, so it controls $\frac{d\theta_\epsilon}{d\lambda}$. In the $\epsilon \rightarrow 0$ limit, $\bar{\rho}_{||} \rightarrow \rho_{||}$. Thus the balance law (5) becomes

$$-R_{\alpha\beta}k^\alpha k^\beta + \frac{d}{d\lambda} \left(\frac{1}{\rho_{||}} \frac{d\rho_{||}}{d\lambda} \right) = 0. \quad (6)$$

12. Now, $\frac{d\rho_{||}}{d\lambda}$ is the same as the covariant derivative of ρ in direction k^α , as $k^\alpha = \frac{dx^\alpha}{d\lambda}$ for coordinates $x^\alpha(\lambda)$ of any null geodesic in the congruence. (Note that there is uniquely one such $k^\alpha(\lambda)$ in the $\epsilon \rightarrow 0$ limit.) Thus

$$-R_{\alpha\beta}k^\alpha k^\beta + \nabla_k \left(\frac{\nabla_k \rho}{\rho} \right) = 0.$$

Some algebra—noting that $k^\alpha \nabla_\alpha k^\beta$ vanishes because k^α picks out a null geodesic congruence—leads to

$$-R_{\alpha\beta}k^\alpha k^\beta + Q_{\alpha\beta}k^\alpha k^\beta = 0$$

where

$$Q_{\alpha\beta} := \frac{1}{\rho} \nabla_\alpha \nabla_\beta \rho - \frac{1}{\rho^2} \nabla_\alpha \rho \nabla_\beta \rho = 0. \quad (7)$$

13. Therefore the balance law Eq. (6) is

$$R_{\alpha\beta}k^\alpha k^\beta = Q_{\alpha\beta}k^\alpha k^\beta. \quad (8)$$

14. Eq. (8) expresses a local, null-contracted tensorial relation between geometric data of W and some Q_{ab} constructed from $W(x, y)$ at coincidence.
15. This law was derived from the smooth data of the bisolution W along null directions. Thus we can take it to encode all geometry-excitation data only if we endorse

Structural Axiom 1: all coupling encoded by $\mathcal{P}_{g[W]}W = 0$ involves geometric and excitation data visible on null directions.

4.1.1 Comment on the Balance Law

1. $Q_{\alpha\beta}k^\alpha k^\beta > 0$ just in case $\nabla_k \left(\frac{\nabla_k \rho}{\rho} \right) > 0$. I.e., $Q_{ab}k^\alpha k^\beta$ is a measure of the change in the normalized change in correlational density ρ along null direction k^α . When it is positive, this normalized second derivative is positive.

4.1.2 Uncontracting the Balance Law - The EFE as the Fixed Point Law for the Free KG Field

1. We may re-write Eq. (8) as

$$(R_{\alpha\beta} - Q_{\alpha\beta})k^\alpha k^\beta = 0.$$

Now, eq. (8) is valid at *all* points in the manifold for *all* vectors k^α that are null with respect to g —its derivation was with respect to any g -null k^α WLOG. Thus we have

$$(R_{\alpha\beta} - Q_{\alpha\beta})k^\alpha k^\beta = 0 \quad \forall k^\alpha \text{ null w.r.t. } g.$$

Thus $R_{\alpha\beta} - Q_{\alpha\beta}$ and $g_{\alpha\beta}$ have precisely the same null cones: $(R - Q)(k, k) = 0$ just in case $g(k, k) = 0$. [Beem et al. \(1996, §2.3\)](#) prove that in such a case, $(R - Q)$ must be conformally related to g , i.e.

$$R_{\alpha\beta} - Q_{\alpha\beta} = \Phi_n(x)g_{\alpha\beta}$$

or

$$R_{\alpha\beta} = Q_{\alpha\beta} + \Phi_n(x)g_{\alpha\beta}. \quad (9)$$

2. Now consider the Einstein tensor $G_{\alpha\beta} := R_{\alpha\beta} - \frac{1}{2}Rg_{\alpha\beta}$. Substituting Eq. (9), it quickly follows that

$$G_{\alpha\beta} + \Phi(x)g_{\alpha\beta} = Q_{\alpha\beta} \quad (10)$$

with $\Phi(x) := \frac{1}{2}R - \Phi_n(x)$.

3. Suppose we postulate

Structural Axiom 2: Any geometric term that appears in any of local-geometric consequences of $P_{g[W]}W = 0$ must be locally constructable from the metric and all of its derivatives.

Then note that [Hollands and Wald \(2001, §5\)](#) show that the only such term including just the metric is a global constant times the metric. Therefore, it must be that $\Phi(x) = \Lambda = \text{const.}$ and, as a consequence $\nabla^\alpha Q_{\alpha\beta} = 0$. That is,

$$\boxed{G_{\alpha\beta} + \Lambda g_{\alpha\beta} = Q_{\alpha\beta}} \quad (11)$$

follows for $P_{g[W]}W = 0$ for the Klein-Gordon field together with the above two structural axioms. This is just the EFE with the matter source in units of curvature—we could define $T_{\alpha\beta} := \frac{Q_{\alpha\beta}}{\kappa}$ to bring Eq. (11) into identical form as Eq. (1).

- Note that this implies that the $W_0 \neq W$ that is a bisolution on $g[W]$ used in our renormalization trick is in fact not a solution of the fixed-point law $P_{g[W]}W = 0$. But this is benign, it was just used as an auxilliary mathematical object to unpack the local geometric consequences of that law.
- Changing the choice of W_0 to another $W'_0 \neq W$ that is a bisolution of $\mathcal{P}_{g[W]}$ changes Q in a way that can be absorbed into the cosmological constant Λ : if $W_0 \rightarrow W'_0$, $J_{S_B}(x) \rightarrow J_{S'_B}(x)$, thus $\rho \rightarrow \rho'$, therefore Eq. (8) changes to $R_{\alpha\beta}k^\alpha k^\beta = Q'_{\alpha\beta}k^\alpha k^\beta$, so we have $(Q_{\alpha\beta} - Q'_{\alpha\beta})k^\alpha k^\beta = 0$; combined with $\nabla^\alpha Q_{\alpha\beta} = \nabla^\alpha Q'_{\alpha\beta} = 0$, this shows that $Q_{\alpha\beta} - Q'_{\alpha\beta} = c_1\Lambda$. Thus $W_0 \rightarrow W'_0$ corresponds to $\Lambda \rightarrow (1 - c_1)\Lambda$ for some c_1 .
- The various $W_0 \neq W$, $W_0 \in \{B_c\}_{g[W]}$ therefore belong together with corresponding Λ to a gauge orbit $\{(W_\Lambda, \Lambda)\} | W_\Lambda \neq W, W_\Lambda \in \{B_c\}_{g[W]}$; it is plausible that each member of such an orbit represents the same solutions of Eq. (11).

Conclusion

I have shown that the EFE sourced by $Q_{\alpha\beta}$ defined from the density of the smooth correlation current $J_{S_B}^\alpha$ result from the fixed-point law $\mathcal{P}_{g[W]}W = 0$ for the free Klein-Gordon field's two-point correlator if we enforce two structural axioms concerning that fixed-point law: first, all of the excitation data and geometric data it couples is visible in some null direction; (ii) all of the geometric data are local functionals of the metric and its derivatives.

This result naturally sits alongside results in “emergent gravity” style research programs, e.g. those of (Cao et al., 2017) and (Jacobson, 1995), which suggest in their own ways that GR is a structural consequence of quantum-theoretic dynamics, rather than a force of its own standing in need of quantization. Whereas (Jacobson, 1995) derives GR as quantum-thermodynamic equation of state, and (Cao et al., 2017) find a GR-like relation concerning spatial curvature as defined from entanglement entropy in a Hilbert space, this result finds the geometry of GR to be that of the two-point correlator of Hadamard states when we identify that geometry with the singular structure ($g_{\alpha\beta} = \lim_{x \rightarrow y} \nabla_{\alpha,x} \nabla_{\beta,x} \sigma(x, y)$), at least for the free Klein Gordon field. Imagining for a moment that the world were only populated by free Klein-Gordon fields, the above result tells us the following: the spacetime internal to those fields—that of its two-point correlator—is automatically governed by Einstein’s Field Equations sourced by local inhomogenities in the smooth part of that two-point correlator. This result thus supports Carroll’s speculative suggestion— independently supported by his co-authored results in Cao et al. (2017)—that “gravity isn’t hard to obtain” in quantum theory, but rather “automatic” (Carroll, 2016). In other words, this result is a step towards finding gravity *in* the quantum world (namely in the $W(x, y)$ defined on $M \times M$), rather than quantizing gravity.

To be clear, a lot of work remains before it can be seen whether this result is more than a structural curiosity, but genuinely relevant for gravitational research. First, the two structural axioms should be argued for. Second, energy conditions on $Q_{\alpha\beta}$ need to be investigated: if $Q_{\alpha\beta}$ is to be identified as real stress-energy, it needs to be positive definite in the field configurations in which we know gravitation occurs. Third, investigation of whether the result generalizes is needed, as the actual world is populated by interacting non-Klein-Gordon fields. One strategy to generalize for other free fields would be to define conserved correlation currents for those fields and proceed in parallel fashion as above. Another would be to investigate whether an algebraic argument from $\mathcal{P}_{g[W]}W = 0$ together with structural axioms 1 + 2 can derive the EFE with a more abstract parsing of the geometric and excitation data it encodes, for example, with a bidifferential operator (as is done in the point-splitting definition of stress-energy in QFTCS, see e.g. Wald (1994, Ch. 4)). This is already desirable, as it should be possible to analyze the local geometric consequences of $\mathcal{P}_{g[W]}W = 0$ without use of an auxiliary bisolution W_0 . This investigation, if yielding positive results, could also be undertaken for interacting fields after suitably generalizing the equation $\mathcal{P}_{g[W]}W = 0$. This latter investigation would also certainly benefit from engagement with the results of Cao et al. (2017) showing the dependence of spatial curvature on the entanglement entropy of real excitations, given the close relation of entanglement entropy and the two-point correlator (Lin et al., 2024).

Pending this further investigation, however, the result is promising for delivering

on Brown's dynamical approach: if the result generalizes with reasonable energy conditions, the empirical support of GR can plausibly be understood as the coarse-grained consequences of microphysical propagation laws of QFT.

References

- Bär, C. and N. Ginoux (2012). Green-hyperbolic operators on globally hyperbolic space-times. *Communications in Mathematical Physics* 313, 485–502.
- Bär, C., N. Ginoux, and F. Pfäffle (2007). *Wave Equations on Lorentzian Manifolds and Quantization*. ESI Lectures in Mathematics and Physics. European Mathematical Society.
- Beem, J. K., P. E. Ehrlich, and K. L. Easley (1996). *Global Lorentzian Geometry* (2 ed.), Volume 202 of *Pure and Applied Mathematics*. New York: Marcel Dekker.
- Birrell, N. D. and P. C. W. Davies (1982). *Quantum Fields in Curved Space*. Cambridge: Cambridge University Press.
- Brown, H. R. (2005). *Physical Relativity: Space-time Structure from a Dynamical Perspective*. Oxford: Oxford University Press.
- Brown, H. R. and O. Pooley (2001). *The origin of the spacetime metric: Bell's 'Lorentzian pedagogy' and its significance in general relativity*, pp. 256–272. Cambridge University Press.
- Brown, H. R. and O. Pooley (2006). Minkowski space-time: A glorious non-entity. In D. G. B. J. Dieks (Ed.), *Ontology of Spacetime*, pp. 67–89. Elsevier.
- Cao, C., S. M. Carroll, and S. Michalakis (2017, Jan). Space from hilbert space: Recovering geometry from bulk entanglement. *Phys. Rev. D* 95, 024031.
- Carroll, S. M. (2016). Space emerging from quantum mechanics. Blog post on Preposterous Universe.
- Deser, S. (1970). Self-interaction and gauge invariance. *General Relativity and Gravitation* 1(1), 9–18.
- Duistermaat, J. J. and L. Hörmander (1972). Fourier integral operators. II. *Acta Mathematica* 128(1), 183–269.
- Einstein, A. (1979). *Autobiographical Notes*. La Salle, Illinois: Open Court Publishing Company. Translated and edited by Paul Arthur Schilpp; originally published in *Albert Einstein: Philosopher–Scientist*.
- Einstein, A. (1992). *Relativity: The Special and General Theory*. London: Routledge. Translated by Robert W. Lawson; first published 1916; Routledge edition reprinted 1993, 1994, 1996.
- Einstein, A. (1998). *The Collected Papers of Albert Einstein, Volume 8: The Berlin Years: Correspondence, 1914–1918*. Princeton, NJ: Princeton University Press. Translation consultant: Klaus Hentschel.
- Fewster, C. J. and R. Verch (2002, February). A quantum weak energy inequality¶for dirac fields in curved spacetime. *Communications in Mathematical Physics* 225(2), 331–359.

- Fewster, C. J. and R. Verch (2013, November). The necessity of the hadamard condition. *Classical and Quantum Gravity* 30(23), 235027.
- Friedlander, F. G. and M. Joshi (1998). *Introduction to the Theory of Distributions* (2 ed.), Volume 14 of *Cambridge Studies in Advanced Mathematics*. Cambridge: Cambridge University Press.
- Fulling, S., F. Narcowich, and R. M. Wald (1981). Singularity structure of the two-point function in quantum field theory in curved spacetime, ii. *Annals of Physics* 136(2), 243–272.
- Fulling, S. A. (1989). *Aspects of Quantum Field Theory in Curved Spacetime*. Cambridge University Press.
- Gingrich, D. M. (2004, March). Conserved current and charge. <https://sites.ualberta.ca/~gingrich/courses/phys512/node26.html>. Accessed: 2025-11-29.
- Gribben, J. (2021). Curved geometries, from john gribben’s “einstein’s masterwork: 1915 and the general theory of relativity.”.
- Gupta, S. N. (1954, Dec). Gravitation and electromagnetism. *Phys. Rev.* 96, 1683–1685.
- Haag, R. (1996). *Local Quantum Physics: Fields, Particles, Algebras* (2 ed.). Texts and Monographs in Physics. Berlin: Springer.
- Hollands, S. and R. M. Wald (2001). Local wick polynomials and time-ordered products of quantum fields in curved spacetime. *Communications in Mathematical Physics* 223, 289–326.
- Jacobson, T. (1995, Aug). Thermodynamics of spacetime: The einstein equation of state. *Phys. Rev. Lett.* 75, 1260–1263.
- Kay, B. S., M. J. Radzikowski, and R. M. Wald (1997, February). Quantum field theory on spacetimes with a compactly generated cauchy horizon. *Communications in Mathematical Physics* 183(3), 533–556.
- Kuo, C.-I. and L. H. Ford (1993, May). Semiclassical gravity theory and quantum fluctuations. *Phys. Rev. D* 47, 4510–4519.
- Lewandowski, M. (2022, 01). Hadamard states for bosonic quantum field theory on globally hyperbolic spacetimes. *Journal of Mathematical Physics* 63(1), 013501.
- Lin, Z. K. et al. (2024). Measuring entanglement entropy and its topological signature. *Nature Communications* 15(1).
- Linnemann, N., C. Smeenk, and M. R. Baker (2026). Gr as a classical spin-2 theory? *Philosophy of Science*.

- Misner, C. W., K. S. Thorne, and J. A. Wheeler (1973). *Gravitation*. San Francisco, CA: W.H. Freeman and Company.
- Padmanabhan, T. (2008, March). From gravitons to gravity: Myths and reality. *International Journal of Modern Physics D* 17(03n04), 367–398.
- Poisson, E. (2004). *A Relativist's Toolkit: The Mathematics of Black-Hole Mechanics*. Cambridge University Press.
- Poisson, E., A. Pound, and I. Vega (2011). The motion of point particles in curved spacetime. *Living Reviews in Relativity* 14(7), 1–187.
- Salimkhani, K. (2020). The dynamical approach to spin-2 gravity. *Studies in History and Philosophy of Science Part B: Studies in History and Philosophy of Modern Physics* 72, 101–112.
- Stachel, J. (1979). Einstein's odyssey: His journey from special to general relativity. *The Sciences* 19(3), 14–34.
- Wald, R. M. (1977). The back reaction effect in particle creation in curved spacetime. *Communications in Mathematical Physics* 54(1), 1–19.
- Wald, R. M. (1984). *General Relativity*. University of Chicago Press.
- Wald, R. M. (1994). *Quantum Field Theory in Curved Spacetime and Black Hole Thermodynamics*. University of Chicago Press.
- Zelditch, S. (2017). Eigenfunctions of the laplacian on riemannian manifolds (cbms lecture notes). Technical report, Northwestern University, Department of Mathematics. Updated: August 15, 2017. Available at <https://sites.math.northwestern.edu/~zelditch/Eigenfunction.pdf>.