

Algebraic letters and NMHV last entry conditions from \bar{Q} -equation

Based on oncoming and recent works with Song He and Zhenjie Li

Chi Zhang

September 12, 2020

Institute of Theoretical Physics, CAS

Last entry conditions

N²MHV Yangian invariants

The N²MHV Yangian invariants have already been classified.

[Arkani-Hamed, Bourjaily, Cachazo, Goncharov, Postnikov, Trnka]

They are

$$Y_1^{(2)} = [1, 2, (23) \cap (456), (234) \cap (56), 6][2, 3, 4, 5, 6]$$

$$Y_2^{(2)} = [1, 2, (34) \cap (567), (345) \cap (67), 7][3, 4, 5, 6, 7]$$

$$Y_3^{(2)} = [1, 2, 3, (345) \cap (67), 7][3, 4, 5, 6, 7]$$

$$Y_4^{(2)} = [1, 2, 3, (456) \cap (78), 8][4, 5, 6, 7, 8]$$

$$Y_5^{(2)} = [1, 2, 3, 4, 8][4, 5, 6, 7, 8]$$

$$Y_6^{(2)} = [1, 2, 3, (45) \cap (678), 8][4, 5, 6, 7, 8]$$

$$Y_7^{(2)} = [1, 2, 3, (45) \cap (678), (456) \cap (78)][4, 5, 6, 7, 8]$$

$$Y_8^{(2)} = [1, 2, 3, 4, (456) \cap (78)][4, 5, 6, 7, 8]$$

$$Y_9^{(2)} = [1, 2, 3, 4, 9][5, 6, 7, 8, 9]$$

$$Y_{10}^{(2)} = [1, 2, 3, 4, (567) \cap (89)][5, 6, 7, 8, 9]$$

$$Y_{11}^{(2)} = [1, 2, 3, 4, (56) \cap (789)][5, 6, 7, 8, 9]$$

$$Y_{12}^{(2)} = \varphi[1, 2, 3, (45) \cap (789), (46) \cap (789)][(45) \cap (123), (46) \cap (123), 7, 8, 9]$$

$$Y_{13}^{(2)} = [1, 2, 3, 4, 5][6, 7, 8, 9, 10]$$

$$Y_{14}^{(2)} = \psi[A, 1, 2, 3, 4][B, 5, 6, 7, 8]$$

where

$$(ij) \cap (klm) = \mathcal{Z}_i \langle j k l m \rangle - \mathcal{Z}_j \langle i k l m \rangle$$

NMHV last entry conditions

For N^2 MHV yangian invariants, this operation gives three kinds of last entries

1.

$$[i\ j\ k\ l\ m] \bar{Q} \log \frac{\langle \bar{n}a \rangle}{\langle \bar{n}b \rangle}$$

2.

$$[1\ i_1\ i_2\ i_3\ i_4] \bar{Q} \log \frac{\langle 1(n-1\ n)(i_1\ i_2)(i_3\ i_4) \rangle}{\langle \bar{n}i_1 \rangle \langle 1i_2i_3i_4 \rangle}, \quad [i_1\ i_2\ i_3\ i_4\ n-1] \bar{Q} \log \frac{\langle n-1(n\ 1)(i_1\ i_2)(i_3\ i_4) \rangle}{\langle \bar{n}i_1 \rangle \langle n-1\ i_2i_3i_4 \rangle}$$

$$[i_1\ i_2\ i_3\ i_4\ n] \bar{Q} \log \frac{\langle n(1\ n-1)(i_1\ i_2)(i_3\ i_4) \rangle}{\langle \bar{n}i_1 \rangle \langle n\ i_2i_3i_4 \rangle} \quad \text{with } 1 < i_1 < i_2 < i_3 < i_4 < n-1$$

where $\langle a(bc)(de)(fg) \rangle := \langle abde \rangle \langle acfg \rangle - \langle acde \rangle \langle abfg \rangle$

3.

$$[i_1\ i_2\ i_3\ i_4\ i_5] \bar{Q} \log \frac{\langle \bar{n}(i_1i_2) \cap (i_3i_4i_5) \rangle}{\langle \bar{n}i_1 \rangle \langle i_2i_3i_4i_5 \rangle}, \quad [i_1\ i_2\ i_3\ i_4\ i_5] \bar{Q} \log \frac{\langle \bar{n}(i_1i_2i_3) \cap (i_4i_5) \rangle}{\langle \bar{n}i_1 \rangle \langle i_2i_3i_4i_5 \rangle},$$

with $1 < i_1 < i_2 < i_3 < i_4 < i_5 < n$

Algebraic letters and words in two-loop NMHV amplitudes

To compute the 2-loop NMHV n point BDS-normalized amplitude, we need the input of the one-loop N^2 MHV BDS-normalized amplitude $R_{n+1,2}^{(1)}$, which can be obtained from the chiral/scalar box expansion [Bourjaily, Caron-Huot, Trnka]:


$$R_{n+1,2}^{(1)} = \sum_{a < b < c < d} (f_{a,b,c,d} - R_{n+1,2}^{\text{tree}} f_{a,b,c,d}^{\text{MHV}}) \mathcal{I}_{a,b,c,d}^{\text{fin}}$$

where

- $f_{a,b,c,d}$ are linear combinations of N^2 MHV yangian invariants
- $f_{a,b,c,d}^{\text{MHV}}$ are either 1 or 0
- $\mathcal{I}_{a,b,c,d}^{\text{fin}}$ denote the finite part of DCI-regulated box integrals

Four-mass box

The most generic term in box expansion:



$$\left\{ \begin{array}{l} u = \frac{x_{ad}^2 x_{bc}^2}{x_{ac}^2 x_{bd}^2}, \quad v = \frac{x_{ab}^2 x_{cd}^2}{x_{ac}^2 x_{bd}^2}, \quad \Delta_{abcd} = \sqrt{(1-u-v)^2 - 4uv} \\ z_{a,b,c,d} = \frac{1}{2}(1+u-v+\Delta), \quad \bar{z}_{a,b,c,d} = \frac{1}{2}(1+u-v-\Delta), \end{array} \right.$$

For such a box,

$$f_{a,b,c,d} = \sum_{\pm} \frac{1-u-v \pm \Delta}{2\Delta} [\alpha_{\pm}, b-1, b, c-1, c] [\delta_{\pm}, d-1, d, a-1, a]$$


$$\mathcal{I}_{a,b,c,d}^{\text{fin}} = \text{Li}_2(z) - \text{Li}_2(\bar{z}) + \frac{1}{2} \log(z\bar{z}) \log \frac{1-z}{1-\bar{z}}$$

where α_{\pm} and δ_{\pm} are two solutions of Schubert problem

$$\alpha = (a-1 a) \cap (d d-1 \gamma), \quad \gamma = (c-1 c) \cap (b b-1 \alpha)$$

Four-mass box

The most generic term in box expansion:



$$\left\{ \begin{aligned} u &= \frac{x_{ad}^2 x_{bc}^2}{x_{ac}^2 x_{bd}^2}, & v &= \frac{x_{ab}^2 x_{cd}^2}{x_{ac}^2 x_{bd}^2}, & \Delta_{abcd} &= \sqrt{(1-u-v)^2 - 4uv} \\ z_{a,b,c,d} &= \frac{1}{2}(1+u-v+\Delta), & \bar{z}_{a,b,c,d} &= \frac{1}{2}(1+u-v-\Delta), \end{aligned} \right.$$

For such a box,

$$f_{a,b,c,d} = \sum_{\pm} \frac{1-u-v \pm \Delta}{2\Delta} [\alpha_{\pm}, b-1, b, c-1, c] [\delta_{\pm}, d-1, d, a-1, a]$$

$$\mathcal{I}_{a,b,c,d}^{\text{fin}} = \text{Li}_2(z) - \text{Li}_2(\bar{z}) + \frac{1}{2} \log(z\bar{z}) \log \frac{1-z}{1-\bar{z}}$$

where α_{\pm} and δ_{\pm} are two solutions of Schubert problem

$$\alpha = (a-1 \ a) \cap (d \ d-1 \ \gamma), \quad \gamma = (c-1 \ c) \cap (b \ b-1 \ \alpha)$$

The square root will disappear when one mass corner become massless,
e.g. $b = a+1$

Rationalize the square root Δ

Only $f_{a,b,c,n+1}$ and $f_{1,b,c,n}$ survive under the $d^{2|3}Z_{n+1}$ integration, since

$$\Delta_{1,b,c,n} \xrightarrow{\mathcal{Z}_{n+1}||\mathcal{Z}_n} 1 - \frac{x_{1b}^2 x_{cn}^2}{x_{1c}^2 x_{bn}^2}$$

Algebraic letters of two-loop NMHV amplitudes

1-D τ -integrals for these four masses introduce new algebraic letters

$$\mathcal{X}_{a,b,c,d}^* := \frac{(x_{a,b,c,d}^* + 1)^{-1} - \bar{z}_{d,a,b,c}}{(x_{a,b,c,d}^* + 1)^{-1} - z_{d,a,b,c}}, \quad \tilde{\mathcal{X}}_{a,b,c,d}^* := \frac{(x_{a,b,c,d-1}^* + 1)^{-1} - z_{d,a,b,c}}{(x_{a,b,c,d-1}^* + 1)^{-1} - \bar{z}_{d,a,b,c}}$$

with 6 choices $a-1, a, b-1, b, c-1, c$ of the superscript, where

$$\begin{aligned} x_{a,b,c,d}^a &= \frac{\langle \bar{d}(c-1c) \cap (ab-1b) \rangle}{\langle \bar{d}a \rangle \langle b-1bc-1c \rangle}, & x_{a,b,c,d}^{a-1} &= x_{a,b,c,d}^a |_{a \leftrightarrow a-1} \\ x_{a,b,c,d}^b &= \frac{\langle \bar{d}(c-1c) \cap (a-1ab) \rangle}{\langle \bar{d}(a-1a) \cap (bc-1c) \rangle}, & x_{a,b,c,d}^{b-1} &= x_{a,b,c,d}^b |_{b \leftrightarrow b-1} \\ x_{a,b,c,d}^c &= \frac{\langle \bar{d}c \rangle \langle a-1ab-1b \rangle}{\langle \bar{d}(a-1a) \cap (b-1bc) \rangle}, & x_{a,b,c,d}^{c-1} &= x_{a,b,c,d}^c |_{c \leftrightarrow c-1} \end{aligned}$$

Note that $\mathcal{X}_{a,b,c,d}^*$, $\mathcal{X}_{b,c,d,a}^*$, $\mathcal{X}_{c,d,a,b}^*$ and $\mathcal{X}_{d,a,b,c}^*$ involve the same square root $\Delta_{a,b,c,d}$

Naïvely, there are would be $12 \times 4 + 2 = 50$ letters associated with the same $\Delta_{a,b,c,d}$.

However, some degeneracy happens when some mass corners only involve 2 particles, for example

$$\mathcal{X}_{d+2,b,c,d}^{d+1} = \frac{\bar{z}_{d,d+2,b,c}}{z_{d,d+2,b,c}}, \quad \tilde{\mathcal{X}}_{a,b,d-2,c}^{d-2} = \frac{1 - z_{d,a,b,d-2}}{1 - \bar{z}_{d,a,b,d-2}}.$$

This leave us $50 - 2m$ algebraic letters associated with the same Δ where

$m =$ the number of corners that contain only two particles

These \mathcal{X} 's and $\tilde{\mathcal{X}}$'s, together with z/\bar{z} and $(1 - z)/(1 - \bar{z})$ give a cyclic and reflection invariant set of algebraic letters

Multiplicative relations among algebraic letters

33 multiplicative relations:

$$\begin{aligned} \frac{\chi_{a,b,c,d}^{a-1}}{\chi_{a,b,c,d}^a} &= \frac{\chi_{d,a,b,c}^a}{\chi_{d,a,b,c}^{a-1}}, & \frac{\chi_{d,a,b,c}^{a-1}}{\chi_{d,a,b,c}^a} &= \frac{\chi_{c,d,a,b}^a}{\chi_{c,d,a,b}^{a-1}}, \\ \frac{\tilde{\chi}_{a,b,c,d}^{a-1}}{\tilde{\chi}_{a,b,c,d}^a} &= \frac{\tilde{\chi}_{d,a,b,c}^a}{\tilde{\chi}_{d,a,b,c}^{a-1}}, & \frac{\tilde{\chi}_{d,a,b,c}^{a-1}}{\tilde{\chi}_{d,a,b,c}^a} &= \frac{\tilde{\chi}_{c,d,a,b}^a}{\tilde{\chi}_{c,d,a,b}^{a-1}}, \\ \frac{\chi_{a,b,c,d}^{a-1}}{\chi_{a,b,c,d}^a} &= \frac{\tilde{\chi}_{c,d,a,b}^a}{\tilde{\chi}_{c,d,a,b}^{a-1}}, & \frac{\chi_{a,b,c,d}^a}{\chi_{a,b,c,d}^b} &= \frac{\tilde{\chi}_{a,b,c,d}^b}{\tilde{\chi}_{a,b,c,d}^a}, & \frac{\chi_{a,b,c,d}^b}{\chi_{a,b,c,d}^c} &= \frac{\tilde{\chi}_{a,b,c,d}^c}{\tilde{\chi}_{a,b,c,d}^b} \end{aligned}$$

and 21 images under the rotations of $a \rightarrow b \rightarrow c \rightarrow d \rightarrow a$,

$$\begin{aligned} \frac{\chi_{a,b,c,d}^a \chi_{b,c,d,a}^d \chi_{c,d,a,b}^c \chi_{d,a,b,c}^a}{\chi_{a,b,c,d}^b \chi_{b,c,d,a}^c \chi_{c,d,a,b}^c \chi_{d,a,b,c}^b} &= 1, \\ \frac{\chi_{a,b,c,d}^a \chi_{b,c,d,a}^d \chi_{d,a,b,c}^a}{\chi_{a,b,c,d}^c \chi_{b,c,d,a}^c \chi_{d,a,b,c}^d} &= 1, & \frac{\chi_{b,c,d,a}^b \chi_{c,d,a,b}^a \chi_{d,a,b,c}^a}{\chi_{b,c,d,a}^c \chi_{c,d,a,b}^c \chi_{d,a,b,c}^b} &= 1, \\ \frac{\chi_{c,d,a,b}^a \chi_{d,a,b,c}^a}{\chi_{c,d,a,b}^d \chi_{d,a,b,c}^d} &= \frac{z_{a,b,c,d}}{\bar{z}_{a,b,c,d}}, & \frac{\chi_{b,c,d,a}^c \chi_{c,d,a,b}^c}{\chi_{b,c,d,a}^d \chi_{c,d,a,b}^d} &= \frac{1 - z_{a,b,c,d}}{1 - \bar{z}_{a,b,c,d}}. \end{aligned}$$

Taking these relations into account:

number of multiplicatively independent algebraic letters = $17 - 2m$

Two kinds of cuts

The algebraic letters can be rewritten as $(a \pm \sqrt{a^2 - 4b})$.

(a, b) are polynomials of Plücker coordinates.

Such letters indicate two kinds of cuts:

- One arise from the discriminant $a^2 - 4b$.
- The other arise from $b \rightarrow 0$ which is the same as the cut of $\log b$.

That is, b must belong to the alphabet of **rational letters**:

For example

$$|(x_{a,b,c,d}^c + 1)^{-1} - \bar{z}_{d,a,b,c}|^2 \propto \langle c(A)(B)(D) \rangle \langle (A) \cap (\bar{d})B(C) \cap (\bar{d}) \rangle \langle AB \rangle ,$$

$$|(x_{a,b,c,d}^b + 1)^{-1} - \bar{z}_{d,a,b,c}|^2 \propto \langle b(A)(C)(D) \rangle \langle (A) \cap (\bar{d})B(C) \cap (\bar{d}) \rangle ,$$

$$|(x_{a,b,c,d}^a + 1)^{-1} - \bar{z}_{d,a,b,c}|^2 \propto \langle a(B)(C)(D) \rangle \langle (A) \cap (\bar{d})B(C) \cap (\bar{d}) \rangle \langle BC \rangle ,$$

here $A = (a-1 a)$, $B = (b-1 b)$, $C = (c-1 c)$, $D = (d-1 d)$

For non-degenerate $\mathcal{X}_{a,b,c,d}^*$'s, then the algebraic words reads

$$\begin{aligned}
 & \mathcal{S}(I_{a,b,c,d}) \otimes \mathcal{X}_{a,b,c,d}^{c-1} \otimes x_{a,b,c,d}^{c-1} [a-1 \ a \ b-1 \ b \ c-1] \\
 & - \mathcal{S}(I_{a,b,c,d}) \otimes \mathcal{X}_{a,b,c,d}^c \otimes x_{a,b,c,d}^c [a-1 \ a \ b-1 \ b \ c] \\
 & + \mathcal{S}(I_{a,b,c,d}) \otimes \mathcal{X}_{a,b,c,d}^{b-1} \otimes x_{a,b,c,d}^{b-1} [a-1 \ a \ b-1 \ c-1 \ c] \\
 & - \mathcal{S}(I_{a,b,c,d}) \otimes \mathcal{X}_{a,b,c,d}^b \otimes x_{a,b,c,d}^c [a-1 \ a \ b \ c-1 \ c] \\
 & + \mathcal{S}(I_{a,b,c,d}) \otimes \mathcal{X}_{a,b,c,d}^{a-1} \otimes x_{a,b,c,d}^{a-1} [a-1 \ b-1 \ b \ c-1 \ c] \\
 & - \mathcal{S}(I_{a,b,c,d}) \otimes \mathcal{X}_{a,b,c,d}^a \otimes x_{a,b,c,d}^a [a \ b-1 \ b \ c-1 \ c] ,
 \end{aligned}$$


likewise for $\tilde{\mathcal{X}}$. Recall that

$$\mathcal{X}^* := \frac{(x^* + 1)^{-1} - \bar{z}}{(x^* + 1)^{-1} - z} ,$$

A class of special components

The $\chi_i \chi_j \chi_k \chi_l$ component with i, j, k, l nonadjacent

- first show up at the two-loop order,
- **completely free** of algebraic letters,
- correspond to the difference of double-pentagon integrals
[Arkani-Hamed, Bourjaily, Cachazo, Trnka]



each of which depend on **many** algebraic roots.[Bourjaily, McLeod, Vergu, Volk, von Hippel, Wilhelm]

- Octagons and algebraic letters at three-loop MHV and two-loop N^2 MHV.
- The connection to cluster algebra, tropical Grassmannian
- \bar{Q} equations for individual integral and other theories.

Thank You

The kernel of \bar{Q}

When $k=1$,

$$\begin{aligned}\bar{Q}\left([1, 2, 3, 4, 5] \log \frac{\langle 1234 \rangle}{\langle 2345 \rangle}\right) &= [1, 2, 3, 4, 5] \bar{Q} \log \frac{\langle 1234 \rangle}{\langle 2345 \rangle} \\ &= (\bar{3})_a [1, 2, 3, 4, 5] \frac{\langle 1234 \rangle \chi_5^A + \text{cyclic}}{\langle 2345 \rangle \langle 2341 \rangle}\end{aligned}$$

When $k=2$, it's easy to show

$$Y_1^{(2)} = \frac{\delta^{0|4}(\langle 1234 \rangle \chi_5 \chi_6 + \text{cyclic})}{\langle 1234 \rangle \cdots \langle 6123 \rangle} \propto \bar{Q} \log u \bar{Q} \log v \bar{Q} \log w$$

then

$$\bar{Q}(Y_1^{(2)} F(u, v, w)) = 0$$

Outline of derivation of \bar{Q} -equation

By using chiral Lagrangian insertion [Caron-Huot], one can show

$$\bar{Q}_{\dot{\alpha}}^A \langle W_{n,k} \rangle \propto \oint dx_{\dot{\alpha}\alpha} \langle (\psi^A + F\theta^A + \dots)^\alpha W_{n,k} \rangle$$

To obtain the \bar{Q} -equation, there are two powerful facts:

- The fermion insertion is the unique twist-one excitation with the quantum numbers of \bar{Q} .
- Its expectation value can be extracted from any object having a nonzero overlap with it in the OPE limit. [Alday, Gaiotto, Maldacena, Sever, Vieira]

$\langle W_{n+1,k+1} \rangle$ has a nonzero overlap with $\bar{Q}_{\dot{\alpha}}^A \langle W_{n,k} \rangle$ under the collinear limit, while $\int d^2|3 \mathcal{Z}_{n+1}$ has the same quantum number as \bar{Q} ,

$$\bar{Q}_{\dot{\alpha}}^A \langle W_{n,k} \rangle \propto \int d^2|3 \mathcal{Z}_{n+1} \langle W_{n+1,k+1} \rangle$$