

Cross-organ (Tabula Sapiens)
ρ = 0.227, P = 9.5×10−14; permutation P = 1×10−4
a
Median kf
0.106
0.127
0.209
0.244
0.246
0.271
n=69
n=32
n=84
n=179
n=267
n=486
0
1-2
3-4
5-6
7-8
9+
CL lineage distance bin
Cross-region (Siletti brain)
ρ = 0.181, P = 9.9×10−13; permutation P = 1×10−4
b
Median kf (fixed 5,000-gene panel)
0.138
0.136
0.136
0.141
0.126
n=173
n=164
n=594
n=516
n=90
1-2
3-4
5-6
7-8
9+
CL lineage distance bin
Oligodendrocyte-lineage chain
52/52 regions monotone; Wilcoxon P = 1.8×10−10
c
Per-region kf
0.063
0.075
0.132
OPC<->COP (dist 2)
COP<->Oligo (dist 5)
OPC<->Oligo (dist 5)
kf replicates; kn reverses across designs
d
Spearman ρ with CL lineage distance
Cross-organ (Tabula Sapiens)
Cross-region (Siletti brain)
***
NS
kf
kn
ω
Germ-layer prediction
Mann–Whitney P = 4.5×10−146
e
Median kf to other classes
0.163
0.123
Microglia (mesoderm) n = 420
Macroglia (ectoderm) n = 652
Distance-0 anchors sit far below any cross-class pair
f
closest cross-class pair
OPC<->COP (0.063)
COP
Choroid plexus
Astrocyte
Vascular
Ependymal
Fibroblast
Bergmann glia
Microglia
Oligodendrocyte
OPC
Median distance-0 kf
(same class, cross-region; 31,764 pairs)