

(a)
Strong (<0.3)
Moderate (<0.5)
Weak (<0.75)
Frequency
Multiplicative residual
(observed / expected ω)
(b)
Top 10 Strong candidates
OL-lineage (n=12 of 39 Strong)
non-OL (n=27)
glia: A14 -- Pul
glia: A14 -- LG
glia: Cla -- LG
ocyte: LP -- TF
yte: STH -- TF
cyte: Pul -- VA
yte: A13 -- MG
glia: DTg -- TF
r: MD-Re -- Pul
te: A13 -- STH
0.204 (ω=7.2)
0.209 (ω=5.6)
0.209 (ω=11.7)
0.213 (ω=11.6)
0.214 (ω=5.4)
0.215 (ω=6.5)
0.224 (ω=9.7)
0.228 (ω=3.4)
0.228 (ω=13.0)
Multiplicative residual
(c)
Microglia
Oligodendrocyte
Fibroblast
Astrocyte
Ependymal
Oligodendrocyte precursor
Committed oligodendrocyte precursor
Vascular
Choroid plexus
Bergmann glia
Strong 16 (0.96%)
Strong 10 (0.95%)
Strong 6 (0.62%)
Strong 3 (0.47%)
Strong 2 (1.25%)
Strong 1 (0.10%)
Strong 1 (0.26%)
0.00%
0.00%
0.00%
Confidence tier (Strong/Moderate/Weak)
Strong
Moderate
Weak
Number of cross-region pairs
(d)
All pairs (n = 31,764)
Block-shuffle null (B = 1,000)
Observed
expected_ω = μ_ct · μ_pair / μ_grand
μ_grand = 38.55
Density
Multiplicative residual (ω / expected_ω)
Lower tail: Strong-tier candidates
Block-shuffle null (B = 1,000)
Observed Strong candidates (n = 39)
Strong cutoff
residual = 0.30
Observed Strong
39 pairs
Null mean
149 ± 37
Density
Multiplicative residual (ω / expected_ω)