

A
17 cross-organ cell-type ranks (n=59 pairs)
Well-sampled (n ≥ 5)
Sparse (n < 5)
CD8+ T
Plasma
Neutrophil
Macrophage
NK
Hepatocyte
B cell
HSC
Smooth muscle
Monocyte
CD4+ T
Classical Mono.
Naive B
Inter. Mono.
Memory B
Endothelial
Erythrocyte
0
5
10
15
20
25
30
Mean CKI ω
B
ω distribution across cross-organ pairs
Conserved (<15), n=29
Variable (≥15), n=30
Number of pairs
0
2
4
6
8
10
12
0
20
40
60
80
CKI ω
C
Cross-organ ω gradient
Mean CKI ω ± SD
0
10
20
30
40
Bone_Marrow-Heart
Heart-Liver
Heart-Lung
Heart-Spleen
Heart-Kidney
Kidney-Lung
Bone_Marrow-Kidney
Liver-Lung
Bone_Marrow-Liver
Bone_Marrow-Spleen
Bone_Marrow-Lung
Kidney-Liver
Lung-Spleen
Kidney-Spleen
Liver-Spleen
D
Top 5 conserved cross-organ pairs
Macrophage
Heart-Bone_Marrow
4.5
Macrophage
Liver-Heart
5.7
Macrophage
Heart-Lung
5.7
Neutrophil
Liver-Bone_Marrow
6.1
CD8+ T
Bone_Marrow-Spleen
6.2
0
1
2
3
4
5
6
CKI ω