

A MEROMORPHIC FUNCTION OMITTING THREE VALUES IN A HALF-PLANE BUT NOT OF BOUNDED TYPE

YIXIN HE AND TENG ZHANG

ABSTRACT. We construct a nonconstant meromorphic function F on the complex plane for which

$$F^{-1}\{0, 1, \infty\} \subset \mathbb{R} \quad \text{and} \quad F|_{\mathbb{H}} \notin N(\mathbb{H}).$$

Thus, omitting three distinct spherical values in a half-plane does not force a meromorphic function on $\mathbb{C}$ to be of bounded type there. The construction combines the modular lambda function with a locally finite Jordan graph of Farey edges. Finite Stern–Brocot refinements, chosen by a Green-kernel diagonal argument, force infinite Green-weighted spherical area while retaining meromorphic Schwarz reflection across the graph.

1. INTRODUCTION

Let $\mathbb{H} = \{z \in \mathbb{C} : \operatorname{Im} z > 0\}$. The Nevanlinna class $N(\mathbb{H})$ consists of the meromorphic functions in $\mathbb{H}$ representable as quotients of two bounded analytic functions. Equivalently, for an analytic function g , membership in $N(\mathbb{H})$ is characterized by the existence of a harmonic majorant of $\log^+ |g|$; see [Gar07, Chapter II, Section 5, pp. 66–71]. Eremenko, Kulikov, and Sodin formulated the following question, tracing it to R. Nevanlinna [EKS26, Question 2, p. 2].

Question 1.1. *Suppose that a meromorphic function on $\mathbb{C}$ omits three distinct values of $\widehat{\mathbb{C}}$ in a half-plane. Must its restriction to that half-plane belong to the Nevanlinna class?*

Their theorem settles related problems for functions meromorphic in variable enlargements of a half-plane, with a sharp logarithmic-integral dichotomy [EKS26, Theorem 1, p. 2]. It does not settle Question 1.1 for a function meromorphic on the whole plane. The purpose of this paper is to give a negative answer.

Theorem 1.2. *There exists a nonconstant meromorphic function $F: \mathbb{C} \rightarrow \widehat{\mathbb{C}}$ such that*

$$F(z) \notin \{0, 1, \infty\}, \quad z \in \mathbb{H},$$

but $F|_{\mathbb{H}} \notin N(\mathbb{H})$. In fact,

$$F^{-1}\{0, 1, \infty\} \subset \mathbb{R}.$$

The construction begins with the modular lambda function

$$\lambda: \mathbb{H} \longrightarrow \widehat{\mathbb{C}} \setminus \{0, 1, \infty\}.$$

Although λ is not of bounded type in $\mathbb{H}$, it cannot itself serve as the required example because it has no meromorphic continuation through the whole real axis. We instead construct a simply connected domain $\Omega \subset \mathbb{H}$ whose boundary is a locally finite chain of Farey edges. If $W: \mathbb{H} \rightarrow \Omega$ is a suitably normalized Riemann map, then $\lambda \circ W$ has finite real boundary values away from a locally finite set. At the exceptional points, finite-width modular cusps become

Date: August 24, 2026.

2020 Mathematics Subject Classification. Primary 30D30; Secondary 30D35, 30C35, 30F20.

Key words and phrases. Nevanlinna class; bounded type; modular lambda function; Farey tessellation; spherical derivative; Green function.

zeros, one-points, or poles after reflection. Consequently, $\lambda \circ W$ extends meromorphically to $\mathbb{C}$.

The domain Ω is selected by a diagonal procedure. In pairwise disjoint bays we make finite Stern–Brocot refinements of increasing, individually chosen depths. The limiting infinite refinement of one bay reaches a real interval. Near that interval its Green function is bounded below by a positive multiple of the height, whereas the height-weighted spherical area of λ is infinite. Monotone convergence therefore permits a finite refinement whose local Green energy exceeds one. The final domain contains each selected one-bay domain, so these disjoint contributions add to infinity.

Throughout the paper, the Green function is normalized by

$$G_D(p, z) > 0, \quad -\Delta_z G_D(p, z) = 2\pi\delta_p,$$

and the spherical derivative of a meromorphic function g is

$$g^\#(z) = \frac{|g'(z)|}{1 + |g(z)|^2}.$$

AI tool disclosure. This counterexample was obtained through AI-assisted exploration under the authors' mathematical supervision and guidance. The authors take responsibility for the statements, citations, and proofs in the final manuscript.

2. GREEN ENERGY AND THE NEVANLINNA CLASS

We first isolate the one direction of the Ahlfors–Shimizu criterion needed later. We use the Riesz decomposition theorem in the normalization of [Ran95, Theorem 3.7.9, p. 76].

Proposition 2.1. *Let D be a Greenian plane domain, let $p \in D$, and let g be holomorphic in D . If $\log^+ |g|$ has a harmonic majorant in D , then*

$$\int_D G_D(p, z) (g^\#(z))^2 dA(z) < \infty.$$

Consequently, if the integral is infinite for $D = \mathbb{H}$, then $g \notin N(\mathbb{H})$.

Proof. Set

$$u(z) = \frac{1}{2} \log(1 + |g(z)|^2).$$

Then

$$0 \leq u \leq \log^+ |g| + \frac{1}{2} \log 2, \quad \Delta u = 2(g^\#)^2.$$

Thus u has a finite harmonic majorant H . Let $D_j \Subset D$ be a smooth exhaustion with $p \in D_1$. The nonnegative function $H - u$ is superharmonic, and the Riesz decomposition theorem on D_j gives

$$\frac{1}{2\pi} \int_{D_j} G_{D_j}(p, z) \Delta u(z) dA(z) \leq H(p) - u(p).$$

As $j \rightarrow \infty$, the Green kernels increase to G_D ; hence monotone convergence yields

$$\frac{1}{\pi} \int_D G_D(p, z) (g^\#(z))^2 dA(z) \leq H(p) - u(p) < \infty.$$

For the last assertion, suppose that $g = h_1/h_2$ with bounded analytic h_1, h_2 in $\mathbb{H}$. Because g is holomorphic, the zeros of h_2 occur among those of h_1 , with at least the same multiplicities. Dividing both functions by the corresponding Blaschke product leaves a bounded zero-free denominator. After a harmless common scaling, minus the logarithm of its modulus is a positive harmonic majorant of $\log^+ |g|$. This is also an immediate consequence of the canonical

factorization in [Gar07, Theorem 5.5, p. 71]. The first part of the proposition now proves the contrapositive. $\square$

We shall repeatedly use domain monotonicity and exhaustion convergence. If $D_m \uparrow D$, if D is Greenian, and if $p \in D_1$, then

$$G_{D_m}(p, z) \uparrow G_D(p, z), \quad z \in D. \quad (2.1)$$

Here and below a Green function is extended by zero outside its domain when integrated over a fixed ambient set. The required monotonicity and convergence are [Ran95, Corollary 4.4.5 and Theorem 4.4.6, p. 108].

3. THE MODULAR LAMBDA FUNCTION

Put $q = e^{\pi i \tau}$ and define

$$\vartheta_2(\tau) = \sum_{n \in \mathbb{Z}} e^{\pi i (n+1/2)^2 \tau}, \quad \vartheta_3(\tau) = \sum_{n \in \mathbb{Z}} e^{\pi i n^2 \tau}, \quad \lambda(\tau) = \frac{\vartheta_2(\tau)^4}{\vartheta_3(\tau)^4}.$$

This fixes our normalization. The following standard facts are recorded in [Ahl79, Chapter 7, Sections 3.4–3.5, pp. 278–282] and, in the theta-function normalization used above, in [NIS26, Equations 23.15.6, 23.17.4, 23.18.1, and 23.18.3]:

- (i) λ is holomorphic in $\mathbb{H}$, is 2-periodic, and omits 0, 1, and ∞ there;
- (ii) λ is invariant under the principal congruence group $\Gamma(2)$;
- (iii) for every $\gamma \in \text{PSL}_2(\mathbb{Z})$ there is an anharmonic transformation R_γ satisfying $\lambda \circ \gamma = R_\gamma \circ \lambda$, where

$$R_\gamma \in \mathcal{A} := \left\{ t \mapsto t, t \mapsto \frac{1}{t}, t \mapsto 1-t, t \mapsto \frac{1}{1-t}, t \mapsto \frac{t}{t-1}, t \mapsto \frac{t-1}{t} \right\}; \quad (3.1)$$

- (iv) uniformly modulo the period 2,

$$\lambda(\tau) = 16q + O(q^2), \quad \text{Im } \tau \rightarrow +\infty; \quad (3.2)$$

- (v) $\lambda(iy)$ is real for $y > 0$.

The quantitative input for the construction is a lower bound for the logarithmic mean of λ . We include the orbit-counting argument, following the idea of [EKS26, Lemma 6, pp. 10–12], in order to make every normalization and multiplicity explicit.

Lemma 3.1. *There exist constants $c > 0$ and $y_0 > 0$ such that*

$$\int_0^2 \log^+ |\lambda(x + iy)| dx \geq c \log \frac{1}{y}, \quad 0 < y < y_0.$$

Proof. Periodicity and (3.2) show that

$$\Lambda(q) := \lambda(\tau), \quad q = e^{\pi i \tau},$$

is well defined and holomorphic in the unit disk, with $\Lambda(0) = 0$. Let $r = e^{-\pi y}$. For $a \in \mathbb{C} \setminus \{0\}$, the elementary inequality

$$\log^+ |u - a| \leq \log^+ |u| + C_a$$

and Jensen's formula applied to $\Lambda - a$ give

$$\int_{-1}^1 \log^+ |\lambda(x + iy)| dx \geq 2 \sum_{|q_k| \leq r} \log \frac{r}{|q_k|} - C'_a,$$

where the q_k are the zeros of $\Lambda - a$, counted with multiplicity. If $|q_k| = r$ for some k , perturb r and pass to the limit; such a boundary zero makes zero contribution. These zeros correspond

to the a -points τ_k of λ in the period strip $-1 \leq \operatorname{Re} \tau < 1$, and $|q_k| = e^{-\pi \operatorname{Im} \tau_k}$. If $\operatorname{Im} \tau_k \geq 2y$, then

$$2 \log \frac{r}{|q_k|} = 2\pi(\operatorname{Im} \tau_k - y) \geq \pi \operatorname{Im} \tau_k.$$

Consequently,

$$\int_{-1}^1 \log^+ |\lambda(x + iy)| dx \geq \pi \sum_{\substack{-1 \leq \operatorname{Re} \tau_k < 1 \\ \operatorname{Im} \tau_k \geq 2y}} \operatorname{Im} \tau_k - C'_a. \quad (3.3)$$

We next supply sufficiently many a -points. For every primitive pair $(c, d) \in \mathbb{Z}^2$, choose $a_0, b_0 \in \mathbb{Z}$ with $a_0 d - b_0 c = 1$. Replacing (a_0, b_0) by $(a_0 + nc, b_0 + nd)$ for a suitable $n \in \mathbb{Z}$, we obtain

$$M_{c,d} = \begin{pmatrix} a_0 & b_0 \\ c & d \end{pmatrix} \in \operatorname{SL}_2(\mathbb{Z})$$

such that $z_{c,d} = M_{c,d}i$ satisfies

$$-\frac{1}{2} \leq \operatorname{Re} z_{c,d} < \frac{1}{2}, \quad \operatorname{Im} z_{c,d} = \frac{1}{c^2 + d^2}. \quad (3.4)$$

The stabilizer of i in $\operatorname{SL}_2(\mathbb{Z})$ is $\{\pm I, \pm S\}$, where $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Thus matrices sending i to the same point have lower rows among

$$(c, d), \quad (-c, -d), \quad (d, -c), \quad (-d, c),$$

and each orbit point is counted at most four times.

For completeness, let

$$L(R) = \#\{(c, d) \in \mathbb{Z}^2 : 0 < c^2 + d^2 \leq R^2\}.$$

Comparison with unit lattice squares gives $L(R) = \pi R^2 + O(R)$. Möbius inversion therefore yields

$$\begin{aligned} N(R) &:= \#\{(c, d) \in \mathbb{Z}^2 : \gcd(c, d) = 1, c^2 + d^2 \leq R^2\} \\ &= \sum_{k \leq R} \mu(k) L(R/k) = \frac{6}{\pi} R^2 + O(R \log R). \end{aligned}$$

This is the standard primitive lattice-point estimate; compare [HW08, Section 18.3, Theorem 332, p. 354]. Subtracting the estimates at radii $\sqrt{X}$ and $\sqrt{X}/2$ shows that, for all large X , the annulus $X/2 < c^2 + d^2 \leq X$ contains at least $c_0 X$ primitive pairs. Summation over dyadic annuli gives

$$\sum_{\substack{\gcd(c,d)=1 \\ 0 < c^2 + d^2 \leq X}} \frac{1}{c^2 + d^2} \geq c_1 \log X \quad (3.5)$$

for all sufficiently large X .

Take $X = (2y)^{-1}$. Equations (3.4) and (3.5), together with the fourfold multiplicity bound, imply

$$\sum_{\substack{z \in \operatorname{PSL}_2(\mathbb{Z}) \cdot i \\ -1 < \operatorname{Re} z < 1, \operatorname{Im} z \geq 2y}} \operatorname{Im} z \geq c_2 \log \frac{1}{y}.$$

Let

$$\mathcal{S} = \{R(\lambda(i)) : R \in \mathcal{A}\}.$$

This is a fixed finite subset of $\mathbb{C} \setminus \{0, 1\}$. By (3.1), every point of the orbit $\operatorname{PSL}_2(\mathbb{Z}) \cdot i$ is an a -point of λ for some $a \in \mathcal{S}$. Partitioning the last sum according to this value, we may choose $a = a_y \in \mathcal{S}$ whose a -points contribute at least $|\mathcal{S}|^{-1}$ of the total. The constants C'_a in (3.3) are

uniformly bounded because $\mathcal{S}$ is finite. After decreasing y_0 and choosing a smaller constant $c > 0$, (3.3) gives the asserted estimate on $[-1, 1]$, and hence on $[0, 2]$ by periodicity. $\square$

We now turn the logarithmic mean into the weighted spherical-area divergence used in the diagonal construction.

Lemma 3.2. *For every $a \in \mathbb{R}$ and every $\varepsilon > 0$,*

$$\int_0^\varepsilon \int_a^{a+2} y(\lambda^\#(x + iy))^2 dx dy = \infty. \quad (3.6)$$

Proof. It suffices to consider small ε . Put

$$v(\tau) = \frac{1}{2} \log(1 + |\lambda(\tau)|^2), \quad M(y) = \int_a^{a+2} v(x + iy) dx.$$

Since λ is 2-periodic and $\Delta v = 2(\lambda^\#)^2$, integration of the x -derivative over one period gives

$$M''(y) = 2 \int_a^{a+2} (\lambda^\#(x + iy))^2 dx.$$

Lemma 3.1 and periodicity imply

$$M(y) \geq c \log \frac{1}{y}$$

for all sufficiently small $y > 0$. On the other hand, (3.2) gives $M(y) \rightarrow 0$ as $y \rightarrow \infty$. The preceding identity makes M convex, and a convex function with a finite limit at $+\infty$ has $M' \leq 0$. Hence, for $0 < \delta < \varepsilon$,

$$\begin{aligned} 2 \int_\delta^\varepsilon \int_a^{a+2} y(\lambda^\#(x + iy))^2 dx dy &= \int_\delta^\varepsilon y M''(y) dy \\ &= \varepsilon M'(\varepsilon) - \delta M'(\delta) - M(\varepsilon) + M(\delta) \\ &\geq M(\delta) - C_\varepsilon, \end{aligned}$$

where $C_\varepsilon = M(\varepsilon) - \varepsilon M'(\varepsilon)$. The right-hand side tends to $+\infty$ as $\delta \downarrow 0$, proving (3.6). $\square$

4. FAREY REFINEMENTS

Write every rational number in reduced form. Two rationals

$$\alpha = \frac{p}{q} < \beta = \frac{r}{s}, \quad q, s > 0,$$

are Farey neighbors if $rq - ps = 1$. Let $e(\alpha, \beta)$ be the upper semicircle with diameter $[\alpha, \beta]$. Its diameter and maximal height are

$$\beta - \alpha = \frac{1}{qs}, \quad \text{height } e(\alpha, \beta) = \frac{1}{2qs}. \quad (4.1)$$

The mediant

$$\alpha \oplus \beta = \frac{p+r}{q+s}$$

is a Farey neighbor of both endpoints. A single refinement replaces $e(\alpha, \beta)$ by

$$e(\alpha, \alpha \oplus \beta) \cup e(\alpha \oplus \beta, \beta).$$

Both children lie in the closed half-disk bounded by the parent semicircle and its diameter. Indeed, on $[\alpha, \alpha \oplus \beta]$ the squared heights of the left child and the parent are, respectively,

$$(x - \alpha)(\alpha \oplus \beta - x) \quad \text{and} \quad (x - \alpha)(\beta - x),$$

and the former is no larger; the other child is analogous. Refining every current leaf edge m times is called the full depth- m refinement.

Lemma 4.1. *Every leaf edge in the full depth- m refinement of the root edge $e(k, k+1)$ has Euclidean diameter at most $(m+1)^{-1}$ and height at most $(2m+2)^{-1}$. Consequently, the corresponding boundary height functions decrease uniformly to zero on $[k, k+1]$; equivalently, the boundary graphs converge to that interval in Hausdorff distance.*

Proof. At depth zero the endpoint denominators are $(1, 1)$. Passing from a pair (q, s) to a child replaces it by $(q, q+s)$ or $(q+s, s)$. Along each branch, the sum of the denominators therefore increases by at least one at every step. At depth m ,

$$q + s \geq m + 2.$$

Since $(q-1)(s-1) \geq 0$, we have $qs \geq q+s-1 \geq m+1$. The conclusion follows from (4.1). $\square$

We next install these finite refinements in separated bays. The initial proper arc chain is

$$\gamma_0 = \bigcup_{k \in \mathbb{Z}} e(k, k+1).$$

It is the graph of a continuous function $h_0 : \mathbb{R} \rightarrow [0, 1/2]$. Define

$$\Omega_0 = \{x + iy : y > h_0(x)\}.$$

For $n \geq 1$, choose the pairwise disjoint bays and their inner intervals

$$B_n = [6n, 6n+4], \quad J_n = [6n+1, 6n+3] \subseteq \text{int } B_n.$$

For fixed n and $m \geq 0$, refine to depth m each of the four root edges over B_n , leaving every other root edge unchanged. Denote the resulting boundary function by $h_{n,m}$ and put

$$D_{n,m} = \{x + iy : y > h_{n,m}(x)\}.$$

On B_n , Lemma 4.1 gives

$$0 \leq h_{n,m}(x) - h_{n,\infty}(x) \leq \frac{1}{2m+2},$$

while the two functions agree off B_n . The child arcs lie below their parents, so the boundary heights decrease with m . Hence $\bigcup_m D_{n,m} = D_{n,\infty}$, and

$$D_{n,m} \uparrow D_{n,\infty}, \tag{4.2}$$

where

$$h_{n,\infty}(x) = \begin{cases} 0, & x \in B_n, \\ h_0(x), & x \notin B_n, \end{cases} \quad D_{n,\infty} = \{x + iy : y > h_{n,\infty}(x)\}.$$

All these domains contain i and lie in $\mathbb{H}$.

5. THE GREEN-ENERGY DIAGONAL CONSTRUCTION

The next lemma converts the limiting divergence in one bay into a finite refinement. Its proof is the point at which boundary regularity of the Green function enters.

Lemma 5.1. *For every $n \geq 1$ there exist $\varepsilon_n > 0$ and an integer $m_n \geq 0$ such that, with*

$$Q_n = \{x + iy : x \in J_n, 0 < y < \varepsilon_n\},$$

one has

$$\int_{Q_n \cap D_{n,m_n}} G_{D_{n,m_n}}(i, z) (\lambda^\#(z))^2 dA(z) > 1.$$

Proof. The limiting domain $D_{n,\infty}$ agrees locally with $\mathbb{H}$ along the interior of B_n . The function

$$z \mapsto G_{D_{n,\infty}}(i, z)$$

is positive and harmonic near J_n in the upper half-plane and has no pole there. Every point of J_n is a regular boundary point because the domain agrees locally with a half-disk. The Green function therefore extends continuously by zero to J_n . Odd reflection across the real segment makes it harmonic in a full neighborhood of J_n . The Hopf boundary point lemma [GT01, Lemma 3.4, p. 34] gives

$$\partial_y G_{D_{n,\infty}}(i, x) > 0, \quad x \in J_n.$$

The reflected function has a continuous derivative. Compactness of J_n and Taylor's formula therefore provide constants $c_n, \varepsilon_n > 0$ such that

$$G_{D_{n,\infty}}(i, x + iy) \geq c_n y, \quad x \in J_n, \quad 0 < y < \varepsilon_n. \quad (5.1)$$

By (3.6) and (5.1),

$$\int_{Q_n} G_{D_{n,\infty}}(i, z) (\lambda^\#(z))^2 dA(z) = \infty.$$

Every point of Q_n belongs to $D_{n,\infty}$ because the limiting boundary vanishes on B_n . Extend $G_{D_{n,m}}(i, \cdot)$ by zero on $Q_n \setminus D_{n,m}$. The domain exhaustion (4.2), the Green-kernel convergence (2.1), and the uniform collapse of the boundary imply pointwise monotone convergence on all of Q_n to $G_{D_{n,\infty}}(i, \cdot)$. Monotone convergence now shows that the finite-depth integrals tend to infinity. Some finite $m = m_n$ gives the required inequality. $\square$

We choose the integer m_n supplied by Lemma 5.1 in the n th bay and make no other change to γ_0 . More precisely, set

$$h(x) = \begin{cases} h_{n,m_n}(x), & x \in B_n \text{ for some } n, \\ h_0(x), & x \notin \bigcup_{n \geq 1} B_n, \end{cases}$$

and define

$$\gamma = \{x + ih(x) : x \in \mathbb{R}\}, \quad \Omega = \{x + iy : y > h(x)\}.$$

At each endpoint of a bay, both adjacent boundary arcs have height zero; hence h is continuous. Each m_n is finite, and every compact real interval meets only finitely many bays, so γ is a locally finite chain of Farey edges. Moreover, $|x + ih(x)| \geq |x|$ and $0 \leq h \leq 1/2$. Thus γ is a continuous proper graph, $\Omega \subset \mathbb{H}$, and $i \in \Omega$.

Proposition 5.2. *The final domain satisfies*

$$\int_{\Omega} G_{\Omega}(i, z) (\lambda^\#(z))^2 dA(z) = \infty. \quad (5.2)$$

Proof. Fix n . On B_n the final boundary agrees with h_{n,m_n} . Outside B_n , every other bay refinement only lowers h_0 , whereas $h_{n,m_n} = h_0$ there. Thus

$$h \leq h_{n,m_n} \quad \text{on } \mathbb{R}, \quad D_{n,m_n} \subset \Omega.$$

Domain monotonicity gives

$$G_{\Omega}(i, z) \geq G_{D_{n,m_n}}(i, z), \quad z \in D_{n,m_n}.$$

The rectangles Q_n are pairwise disjoint. Hence, for $N \geq 1$, Lemma 5.1 yields

$$\begin{aligned} \int_{\Omega} G_{\Omega}(i, z) (\lambda^{\#}(z))^2 dA(z) &\geq \sum_{n=1}^N \int_{Q_n \cap D_{n, m_n}} G_{\Omega}(i, z) (\lambda^{\#}(z))^2 dA(z) \\ &\geq \sum_{n=1}^N \int_{Q_n \cap D_{n, m_n}} G_{D_{n, m_n}}(i, z) (\lambda^{\#}(z))^2 dA(z) > N. \end{aligned}$$

Letting $N \rightarrow \infty$ proves (5.2). $\square$

6. REFLECTION ACROSS THE FAREY BOUNDARY

It remains to show that uniformization followed by Schwarz reflection produces a function meromorphic at every reflected vertex. We begin with the open edges.

Lemma 6.1. *The function λ takes finite real values on the interior of every Farey edge.*

Proof. The positive imaginary axis is the hyperbolic geodesic joining 0 to ∞ , and $\lambda(iy)$ is real for $y > 0$. Every Farey edge is the image of this geodesic under an element of $\mathrm{PSL}_2(\mathbb{Z})$. Each transformation in (3.1) has real coefficients and preserves $\widehat{\mathbb{R}}$. Thus λ is real on every open Farey edge. It is finite there because it is holomorphic and omits ∞ in $\mathbb{H}$. $\square$

Let V be the vertex set of γ . Every vertex is rational, V is locally finite in $\mathbb{C}$, and its only accumulation point in the Riemann sphere is ∞ . The local geometry at a vertex becomes especially transparent after a modular transformation.

Lemma 6.2. *Let $v \in V$, and let $e(v, u_1)$ and $e(v, u_2)$ be the two boundary edges incident to v . There exist $A \in \mathrm{PSL}_2(\mathbb{Z})$ with $A(v) = \infty$ and two distinct integers k_1, k_2 such that $A(e(v, u_j))$ is the vertical geodesic $\{\mathrm{Re} \zeta = k_j, \mathrm{Im} \zeta > 0\}$. Moreover, above some height, the end of $A(\Omega)$ corresponding to v is exactly the finite-width strip between these two geodesics. In particular,*

$$\mathrm{Im} A(w) \longrightarrow +\infty \quad \text{as } w \rightarrow v \text{ within } \Omega. \quad (6.1)$$

Proof. Write $v = p/q$ in reduced form, with $q > 0$, and choose $a, b \in \mathbb{Z}$ so that $ap + bq = -1$. Then

$$A(w) = \frac{aw + b}{qw - p}$$

belongs to $\mathrm{PSL}_2(\mathbb{Z})$ and sends v to ∞ . The modular group preserves Farey adjacency. A Farey neighbor of $\infty = 1/0$ has denominator one and is therefore an integer. Hence $A(u_1)$ and $A(u_2)$ are distinct integers, and the incident geodesics become the corresponding vertical lines.

We determine the relevant side. Label the adjacent vertices so that $u_1 < v < u_2$. If $u = r/s$ is either neighbor, direct calculation gives

$$A(u) - \frac{a}{q} = -\frac{s}{q(qr - ps)}.$$

The determinant $qr - ps$ equals -1 for the left neighbor and 1 for the right neighbor, so the two integers $A(u_1)$ and $A(u_2)$ lie on opposite sides of a/q . Furthermore,

$$A(v + it) = \frac{a}{q} + \frac{i}{q^2 t}, \quad t > 0,$$

and $v + it \in \Omega$.

Local finiteness and the graph property provide a disk U about v meeting γ only in the two incident edge germs. The component of $\Omega \cap U$ accumulating at v is the region between those germs. Its image under A therefore lies between the two vertical geodesics. If $w \rightarrow v$ in Ω ,

then $|A(w)| \rightarrow \infty$, while the real part remains between the two corresponding integers. This proves (6.1).

Finally, the sets $A^{-1}\{\operatorname{Im} \zeta > Y\}$ are horodisks shrinking to v as $Y \rightarrow \infty$. Choose Y so large that such a horodisk lies in U . Above height Y , the only boundary pieces of $A(\Omega)$ are the two vertical geodesics. Since the middle interval contains $A(v + it)$ for small t , crossing either boundary geodesic exits the Jordan domain. Thus, with $k_- = \min(k_1, k_2)$ and $k_+ = \max(k_1, k_2)$,

$$A(\Omega) \cap \{\operatorname{Im} \zeta > Y\} = \{\zeta : k_- < \operatorname{Re} \zeta < k_+, \operatorname{Im} \zeta > Y\}.$$

□

The cusp description makes every exceptional reflected point an isolated meromorphic singularity.

Proposition 6.3. *There is a conformal map $W: \mathbb{H} \rightarrow \Omega$ satisfying*

$$W(i) = i, \quad W(\infty) = \infty$$

in the sense of prime ends. For such a map, $f = \lambda \circ W$ extends by reflection to a meromorphic function F on $\mathbb{C}$, and

$$F^{-1}\{0, 1, \infty\} \subset \mathbb{R}.$$

Proof. Because γ is a continuous proper graph, its one-point compactification $\gamma \cup \{\infty\}$ is a Jordan curve in the Riemann sphere. Thus Ω is a Jordan domain on the sphere, with ∞ as a boundary point. After applying a Möbius transformation to reduce to bounded Jordan domains, the Riemann mapping theorem and Carathéodory's boundary theorem [Ahl79, Chapter 6, Sections 1.1–1.2, especially p. 232] give a conformal map from $\mathbb{H}$ onto Ω that extends homeomorphically to the compactified boundaries. Precomposition with an automorphism of $\mathbb{H}$ first arranges that ∞ corresponds to ∞ . If $z_0 = x_0 + iy_0$ is then the preimage of i , further precomposition with $z \mapsto x_0 + y_0 z$ fixes ∞ and sends i to z_0 . This gives the stated normalization.

Let

$$E = W^{-1}(V) \subset \mathbb{R},$$

where the boundary extension is understood. Since V has no finite accumulation point and $W(\infty) = \infty$, the set E is locally finite in $\mathbb{R}$. On every component of $\mathbb{R} \setminus E$, the boundary values of W lie in the interior of one Farey edge. Lemma 6.1 therefore gives finite real boundary values for f there. Define

$$F(z) = \begin{cases} f(z), & \operatorname{Im} z > 0, \\ \overline{f(\bar{z})}, & \operatorname{Im} z < 0, \end{cases}$$

and fill in $\mathbb{R} \setminus E$ by continuous extension. Schwarz reflection [Ahl79, Chapter 4, Section 6.5, Theorem 24, pp. 172–173] shows that F is holomorphic on $\mathbb{C} \setminus E$.

Fix $x_v \in E$, with $W(x_v) = v \in V$, and choose a disk about x_v containing no other point of E . Reflection across the two punctured real intervals makes F holomorphic on the full punctured disk. Let A be as in Lemma 6.2. Boundary continuity and (6.1) imply

$$\operatorname{Im} A(W(z)) \longrightarrow +\infty \quad (z \rightarrow x_v, z \in \mathbb{H}).$$

Apply (3.1) to A^{-1} and put $\zeta = A(W(z))$. Then

$$f(z) = \lambda(W(z)) = R_{A^{-1}}(\lambda(A(W(z)))).$$

Equation (3.2) yields, in the spherical metric,

$$f(z) \longrightarrow a_v := R_{A^{-1}}(0) \in \{0, 1, \infty\}.$$

The same spherical limit holds in the reflected lower half-disk. If $a_v = 0$ or 1 , the singularity is removable. If $a_v = \infty$, then $|F(z)| \rightarrow \infty$ as $z \rightarrow x_v$ through the punctured disk. Thus $1/F$ is holomorphic there, tends to zero, and has a removable singularity. Since $1/F$ is not identically zero, its zero has finite order, and F has a pole of finite order at x_v .

Because E is locally finite, filling in all its points produces a meromorphic function on $\mathbb{C}$. It is nonconstant because $\lambda \circ W$ is nonconstant in $\mathbb{H}$. Both open half-planes omit 0 , 1 , and ∞ , so every preimage of these values is real. $\square$

The normalization at the boundary point ∞ is not cosmetic. We record its role explicitly.

Remark 6.4. The condition $W(\infty) = \infty$ ensures that the preimages of vertices escaping to infinity cannot accumulate at a finite point of $\mathbb{R}$. Without it, the reflected function need not be meromorphic at such an accumulation point.

7. PROOF OF THE MAIN THEOREM

We now combine the reflection theorem with the Green-energy obstruction.

Proof of Theorem 1.2. Let Ω be the domain constructed in Sections 4–5, let W be the map in Proposition 6.3, and set

$$f = \lambda \circ W \quad \text{in } \mathbb{H}.$$

Proposition 6.3 supplies a nonconstant meromorphic extension F to $\mathbb{C}$ and shows that f omits 0 , 1 , and ∞ in $\mathbb{H}$.

Green functions and spherical area are conformally invariant. Since $W(i) = i$, the change of variables $w = W(z)$ gives

$$\int_{\mathbb{H}} G_{\mathbb{H}}(i, z) (f^{\#}(z))^2 dA(z) = \int_{\Omega} G_{\Omega}(i, w) (\lambda^{\#}(w))^2 dA(w) = \infty$$

by (5.2). Proposition 2.1 implies $f \notin N(\mathbb{H})$, completing the proof. $\square$

8. CONTEXT AND CONSEQUENCES

We use the standard Nevanlinna notation

$$\rho(F) = \limsup_{r \rightarrow \infty} \frac{\log T(r, F)}{\log r}, \quad \delta(a, F) = 1 - \limsup_{r \rightarrow \infty} \frac{N(r, a; F)}{T(r, F)}.$$

Theorem 1.2 does not conflict with Nevanlinna's classical positive theorem. Nevanlinna proved that a finite-order meromorphic function on $\mathbb{C}$ which omits three values in a half-plane is of bounded type there; the precise location is [Nev25, Satz II and the following Hilfssatz, p. 18]. See also the modern discussion in [EKS26, Section 2.1, pp. 3–4]. Consequently, the function constructed above has infinite order.

Nor does Edrei's theorem on radially distributed values rule out the example. Specialized to derivative order $l = 0$ and to the two rays with arguments 0 and π , [Edr55, Theorem 1, p. 276] says that if all zeros, one-points, and poles of a meromorphic function lie on these rays and

$$\delta(0, F) + \delta(1, F) + \delta(\infty, F) > 0, \tag{8.1}$$

then F has finite order. More precisely, both complementary angular gaps are π , so Edrei's bound is

$$\rho(F) \leq \max \left\{ \frac{\pi}{\pi}, \frac{\pi}{\pi} \right\} = 1.$$

The positive-deficiency hypothesis is part of the original theorem and is reproduced explicitly in [WZ14, Theorem A, p. 409]. The informal summary in [EKS26, Section 2.3, p. 5] does not display this hypothesis and should not be used in place of Edrei's exact statement.

If (8.1) held for our function, Edrei's theorem would make its order finite, and Nevanlinna's theorem would then put $F|_{\mathbb{H}}$ in $N(\mathbb{H})$, a contradiction. Since deficiencies are nonnegative, the construction necessarily satisfies

$$\delta(0, F) = \delta(1, F) = \delta(\infty, F) = 0.$$

Finally, we have neither used nor proved that the complete set of singular values of the reflected function is exactly $\{0, 1, \infty\}$. That stronger assertion would require a separate analysis of possible asymptotic values at infinity.

REFERENCES

- [Ahl79] L. V. Ahlfors, *Complex analysis*, 3rd ed., McGraw-Hill Book Co., New York, 1979.
- [NIS26] NIST Digital Library of Mathematical Functions, F. W. J. Olver et al., eds., Release 1.2.7 of June 15, 2026, Chapter 23, Sections 23.15, 23.17, and 23.18, <https://dlmf.nist.gov/23> (accessed August 24, 2026), doi:[10.18434/T4HW23](https://doi.org/10.18434/T4HW23).
- [Edr55] A. Edrei, *Meromorphic functions with three radially distributed values*, Trans. Amer. Math. Soc. **78** (1955), no. 2, 276–293, doi:[10.1090/S0002-9947-1955-0067982-9](https://doi.org/10.1090/S0002-9947-1955-0067982-9).
- [EKS26] A. Eremenko, A. Kulikov, and M. Sodin, *When a meromorphic function that omits three values has a bounded type*, revised preprint, July 24, 2026, [author's revised PDF](#) (accessed August 24, 2026), arXiv:2604.06136, doi:[10.48550/arXiv.2604.06136](https://doi.org/10.48550/arXiv.2604.06136).
- [Gar07] J. B. Garnett, *Bounded analytic functions*, revised 1st ed., Graduate Texts in Mathematics, vol. 236, Springer, New York, 2007, doi:[10.1007/0-387-49763-3](https://doi.org/10.1007/0-387-49763-3).
- [GT01] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, doi:[10.1007/978-3-642-61798-0](https://doi.org/10.1007/978-3-642-61798-0).
- [HW08] G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, 6th ed., Oxford University Press, Oxford, 2008, doi:[10.1093/oso/9780199219858.001.0001](https://doi.org/10.1093/oso/9780199219858.001.0001).
- [Nev25] R. Nevanlinna, *Über die Eigenschaften meromorpher Funktionen in einem Winkelraum*, Acta Soc. Sci. Fenn. **50** (1925), no. 12, 1–45.
- [Ran95] T. Ransford, *Potential theory in the complex plane*, London Mathematical Society Student Texts, vol. 28, Cambridge University Press, Cambridge, 1995, doi:[10.1017/CBO9780511623776](https://doi.org/10.1017/CBO9780511623776).
- [WZ14] N. Wu and J.-H. Zheng, *On the growth order of an algebroid function with radially distributed values in the unit disk*, Tohoku Math. J. (2) **66** (2014), no. 3, 409–425, doi:[10.2748/tmj/1412783205](https://doi.org/10.2748/tmj/1412783205).

SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, SHANGHAI 200433, P. R. CHINA
 Email address: yixin.he717@gmail.com

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN 710049, P. R. CHINA
 Email address: teng.zhang@stu.xjtu.edu.cn