


195


Preferable examples of the oxime sulfonate-based acid generator include compounds represented by general formula (B-2) or (B-3) shown below.

[Chemical Formula 98]



In the formula (B-2), R³³ represents a cyano group, an alkyl group having no substituent or a halogenated alkyl group; R³⁴ represents an aryl group; and R³⁵ represents an alkyl group having no substituent or a halogenated alkyl group.

[Chemical Formula 99]



In the formula (B-3), R³⁶ represents a cyano group, an alkyl group having no substituent or a halogenated alkyl group; R³⁷ represents a divalent or trivalent aromatic hydrocarbon group; R³⁸ represents an alkyl group having no substituent or a halogenated alkyl group; and p'' represents 2 or 3.

In general formula (B-2), the alkyl group having no substituent or the halogenated alkyl group for R³³ preferably has 1 to 10 carbon atoms, more preferably 1 to 8 carbon atoms, and most preferably 1 to 6 carbon atoms.

As R³³, a halogenated alkyl group is preferable, and a fluorinated alkyl group is more preferable.

The fluorinated alkyl group for R³³ preferably has 50% or more of the hydrogen atoms thereof fluorinated, more preferably 70% or more, and most preferably 90% or more.

Examples of the aryl group for R³⁴ include groups in which one hydrogen atom has been removed from an aromatic hydrocarbon ring, such as a phenyl group, a biphenyl group, a fluorenyl group, a naphthyl group, an anthryl group, and a phenantryl group, and heteroaryl groups in which some of the carbon atoms constituting the ring(s) of these groups are substituted with hetero atoms such as an oxygen atom, a sulfur atom, and a nitrogen atom. Of these, a fluorenyl group is preferable.

The aryl group for R³⁴ may have a substituent such as an alkyl group of 1 to 10 carbon atoms, a halogenated alkyl group, or an alkoxy group. The alkyl group and halogenated alkyl group as the substituent preferably has 1 to 8 carbon atoms, and more preferably 1 to 4 carbon atoms. Further, the halogenated alkyl group is preferably a fluorinated alkyl group.

The alkyl group having no substituent or the halogenated alkyl group for R³⁵ preferably has 1 to 10 carbon atoms, more preferably 1 to 8 carbon atoms, and most preferably 1 to 6 carbon atoms.

As R³⁵, a halogenated alkyl group is preferable, and a fluorinated alkyl group is more preferable.

In terms of enhancing the strength of the acid generated, the fluorinated alkyl group for R³⁵ preferably has 50% or more of the hydrogen atoms fluorinated, more preferably 70% or more, still more preferably 90% or more. A completely fluorinated alkyl group in which 100% of the hydrogen atoms are substituted with fluorine atoms is particularly desirable.

196

In general formula (B-3), as the alkyl group having no substituent and the halogenated alkyl group for R³⁶, the same alkyl group having no substituent and the halogenated alkyl group described above for R³³ can be used.

5 Examples of the divalent or trivalent aromatic hydrocarbon group for R³⁷ include groups in which one or two hydrogen atoms have been removed from the aryl group for R³⁴.

10 As the alkyl group having no substituent or the halogenated alkyl group for R³⁸, the same one as the alkyl group having no substituent or the halogenated alkyl group for R³⁵ can be used.

p'' is preferably 2.

Specific examples of suitable oxime sulfonate-based acid generators include α-(p-toluenesulfonyloxyimino)-benzyl cyanide, α-(p-chlorobenzenesulfonyloxyimino)-benzyl cyanide, α-(4-nitrobenzenesulfonyloxyimino)-benzyl cyanide, α-(4-nitro-2-trifluoromethylbenzenesulfonyloxyimino)-benzyl cyanide, α-(benzenesulfonyloxyimino)-4-chlorobenzyl cyanide, α-(benzenesulfonyloxyimino)-2,4-dichlorobenzyl cyanide, α-(benzenesulfonyloxyimino)-2,6-dichlorobenzyl cyanide, α-(benzenesulfonyloxyimino)-4-methoxybenzyl cyanide, α-(2-chlorobenzenesulfonyloxyimino)-4-methoxybenzyl cyanide, α-(benzenesulfonyloxyimino)-thien-2-yl acetonitrile, α-(4-dodecylbenzenesulfonyloxyimino)benzyl cyanide, α-[(p-toluenesulfonyloxyimino)-4-methoxyphenyl]acetonitrile, α-[(dodecylbenzenesulfonyloxyimino)-4-methoxyphenyl]acetonitrile, α-(tosyloxyimino)-4-thienyl cyanide, α-(methylsulfonyloxyimino)-1-cyclopentenyl acetonitrile, α-(methylsulfonyloxyimino)-1-cyclohexenyl acetonitrile, α-(methylsulfonyloxyimino)-1-cycloheptenyl acetonitrile, α-(methylsulfonyloxyimino)-1-cyclooctenyl acetonitrile, α-(trifluoromethylsulfonyloxyimino)-1-cyclopentenyl acetonitrile, α-(trifluoromethylsulfonyloxyimino)-ethyl cyclohexyl acetonitrile, α-(ethylsulfonyloxyimino)-ethyl acetonitrile, α-(propyl sulfonyloxyimino)-propyl acetonitrile, α-(cyclohexylsulfonyloxyimino)-cyclopentyl acetonitrile, α-(cyclohexylsulfonyloxyimino)-cyclohexyl acetonitrile, α-(cyclohexylsulfonyloxyimino)-1-cyclopentenyl acetonitrile, α-(ethylsulfonyloxyimino)-1-cyclopentenyl acetonitrile, α-(isopropylsulfonyloxyimino)-1-cyclopentenyl acetonitrile, α-(n-butylsulfonyloxyimino)-1-cyclopentenyl acetonitrile, α-(ethylsulfonyloxyimino)-1-cyclohexenyl acetonitrile, α-(isopropylsulfonyloxyimino)-1-cyclohexenyl acetonitrile, α-(n-butylsulfonyloxyimino)-1-cyclohexenyl acetonitrile, α-(methylsulfonyloxyimino)-phenyl acetonitrile, α-(methylsulfonyloxyimino)-p-methoxyphenyl acetonitrile, α-(trifluoromethylsulfonyloxyimino)-phenyl acetonitrile, α-(trifluoromethylsulfonyloxyimino)-p-methoxyphenyl acetonitrile, α-(ethylsulfonyloxyimino)-p-methoxyphenyl acetonitrile, α-(propylsulfonyloxyimino)-p-methylphenyl acetonitrile, and α-(methyl sulfonyloxyimino)-p-bromophenyl acetonitrile.

60 Further, oxime sulfonate-based acid generators disclosed in Japanese Unexamined Patent Application, First Publication No. Hei 9-208554 (Chemical Formulas 18 and 19 shown in paragraphs [0012] to [0014]) and oxime sulfonate-based acid generators disclosed in WO 2004/074242A2 (Examples 1 to 40 described at pages 65 to 86) may be preferably used.

Furthermore, as preferable examples, the following can be used.