Topological Gravity

Xuhui Zhang*1

¹Department of Mathematical Science, Tsinghua University

May 19, 2024

A note of [DW18].

1 Introduction

There are two approaches to study 2D quantum gravity. One is through matrix model, which is developed in 1980's, see [FGZ95] for a comprehensive review. The other one is through topological gravity, whose corelators are given by the intersection numbers on the moduli space of Riemann Surfaces. Witten conjectured that these two approaches are equivalent[Wit90a, Wit90b], and his conjecture is proved by Kontsevich[Kon92]. Mirzakhani also gave a new proof of Witten conjecture by using her recursion relations for Weil-Petersson volume $V_{g,b}$ of moduli space of hyperbolic metrics on Riemann surface of genus g with g boundaries, where boundaries are geodesics of length g be g

2 Matrix Model

References

cicalGravity2018	[DW18]	Robbert Dijkgraaf and Edward Witten. Developments in Topological Gravity. International Journal of Modern Physics A, 33(30):1830029, October 2018.
avityRandom1995	[FGZ95]	P. Di Francesco, P. Ginsparg, and J. Zinn-Justin. 2D gravity and random matrices. <i>Physics Reports</i> , 254(1):1–133, March 1995.
heoryModuli1992	[Kon92]	Maxim Kontsevich. Intersection theory on the moduli space of curves and the matrix airy function. <i>Communications in Mathematical Physics</i> , 147(1):1–23, June 1992.
ogicalPhase1990	[Wit90a]	Edward Witten. On the structure of the topological phase of two-dimensional gravity. Nuclear Physics B, $340(2-3):281-332$, 1990 .
ntersection1990	[Wit90b]	Edward Witten. Two-dimensional gravity and intersection theory on moduli space. Surveys in differential geometry, $1(1):243-310$, 1990 .

^{*}zhangxh.math@gmail.com