Ewald Force

zhaoyihao

zhaoyihao@protonmail.com

2022年2月12日

实空间:

$$U^{R} = \frac{1}{2} \sum_{\mathbf{n}} \sum_{i}^{N} \sum_{j}^{N} q_{i} q_{j} \frac{erfc(\alpha |\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|)}{|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|}$$

所以对于 j 粒子:

$$U_j^R = q_j \sum_{\mathbf{n}} \sum_{i \neq j} q_i \frac{erfc(\alpha |\mathbf{r}_i - \mathbf{r}_j + \mathbf{n}|)}{|\mathbf{r}_i - \mathbf{r}_j + \mathbf{n}|}$$

$$\begin{aligned} \mathbf{F}_{j}^{R} &= -\nabla U_{j}^{R} = -q_{j} \sum_{\mathbf{n}} \sum_{i \neq j} q_{i} \frac{d \frac{erfc(\alpha|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|)}{|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|}}{d\mathbf{r}_{j}} \\ &= -q_{j} \sum_{\mathbf{n}} \sum_{i \neq j} q_{i} \frac{2 \sqrt{\pi} (-\alpha) e^{-\alpha^{2}|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|^{2}} \frac{d|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|}{d\mathbf{r}_{j}} |\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}| - erfc(\alpha|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|) \frac{d|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|}{d\mathbf{r}_{j}}}{|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|^{2}} \\ &= -q_{j} \sum_{\mathbf{n}} \sum_{i \neq j} q_{i} \frac{2\alpha}{\sqrt{\pi}} e^{-\alpha^{2}|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|^{2}} (\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}) + erfc(\alpha|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|) \frac{\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}}{|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|} \\ &= -q_{j} \sum_{\mathbf{n}} \sum_{i \neq j} q_{i} \left[\frac{2\alpha}{\sqrt{\pi}} e^{-\alpha^{2}|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|^{2}} + \frac{1}{|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|} erfc(\alpha|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|) \right] \frac{\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}}{|\mathbf{r}_{i} - \mathbf{r}_{j} + \mathbf{n}|^{2}} \end{aligned} \tag{1}$$

其中:

$$|\mathbf{r}_i - \mathbf{r}_j| = \sqrt{(x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2}$$

$$\frac{d|\mathbf{r}_{i} - \mathbf{r}_{j}|}{d\mathbf{r}_{j}} = \left(\frac{\partial|\mathbf{r}_{i} - \mathbf{r}_{j}|}{\partial x_{j}}, \frac{\partial|\mathbf{r}_{i} - \mathbf{r}_{j}|}{\partial y_{j}}, \frac{\partial|\mathbf{r}_{i} - \mathbf{r}_{j}|}{\partial z_{j}}\right)
= \left(-\frac{x_{i} - x_{j}}{|\mathbf{r}_{i} - \mathbf{r}_{j}|}, -\frac{y_{i} - y_{j}}{|\mathbf{r}_{i} - \mathbf{r}_{j}|}, -\frac{z_{i} - z_{j}}{|\mathbf{r}_{i} - \mathbf{r}_{j}|}\right)
= -\frac{\mathbf{r}_{i} - \mathbf{r}_{j}}{|\mathbf{r}_{i} - \mathbf{r}_{j}|}$$
(2)

k 空间:

$$U^{k} = \frac{2\pi}{V} \sum_{\mathbf{k} \neq \mathbf{0}} \frac{1}{k^{2}} e^{-\frac{k^{2}}{4\alpha^{2}}} \sum_{i}^{N} \sum_{j}^{N} q_{i} q_{j} e^{i\mathbf{k}(\mathbf{r}_{i} - \mathbf{r}_{j})}$$

对于 j 粒子:

$$U_j^k = q_j \frac{2\pi}{V} \sum_{\mathbf{k} \neq \mathbf{0}} \frac{1}{k^2} e^{-\frac{k^2}{4\alpha^2}} \sum_i^N q_i e^{i\mathbf{k}(\mathbf{r}_i - \mathbf{r}_j)}$$

$$\mathbf{F}_{j}^{k} = -\nabla U_{j}^{k} = -q_{j} \frac{2\pi}{V} \sum_{\mathbf{k} \neq 0} \frac{1}{k^{2}} e^{-\frac{k^{2}}{4\alpha^{2}}} \sum_{i}^{N} q_{i} \frac{e^{i\mathbf{k}(\mathbf{r}_{i} - \mathbf{r}_{j})}}{d\mathbf{r}_{j}}$$

$$= -q_{j} \frac{2\pi}{V} \sum_{\mathbf{k} \neq 0} \frac{1}{k^{2}} e^{-\frac{k^{2}}{4\alpha^{2}}} \sum_{i}^{N} q_{i} (-i\mathbf{k}) e^{i\mathbf{k}(\mathbf{r}_{i} - \mathbf{r}_{j})}$$

$$= -q_{j} \frac{2\pi}{V} \sum_{\mathbf{k} \neq 0} \frac{1}{k^{2}} e^{-\frac{k^{2}}{4\alpha^{2}}} \sum_{i}^{N} q_{i} (-i\mathbf{k}) \{\cos[\mathbf{k} \cdot (\mathbf{r}_{i} - \mathbf{r}_{j})] + i\sin[\mathbf{k} \cdot (\mathbf{r}_{i} - \mathbf{r}_{j})]\}$$

$$= -q_{j} \frac{2\pi}{V} \sum_{\mathbf{k} \neq 0} \frac{\mathbf{k}}{k^{2}} e^{-\frac{k^{2}}{4\alpha^{2}}} \sum_{i}^{N} q_{i} \sin[\mathbf{k} \cdot (\mathbf{r}_{i} - \mathbf{r}_{j})]$$

$$+ iq_{j} \frac{2\pi}{V} \sum_{\mathbf{k} \neq 0} \frac{\mathbf{k}}{k^{2}} e^{-\frac{k^{2}}{4\alpha^{2}}} \sum_{i}^{N} q_{i} \cos[\mathbf{k} \cdot (\mathbf{r}_{i} - \mathbf{r}_{j})]$$

$$(3)$$

当 $\mathbf{k} \in [-\mathbf{kmax}, \mathbf{kmax}]$ 时, \cos 是偶函数,此时虚部为 0,则:

$$\mathbf{F}_{j} = -q_{j} \frac{2\pi}{V} \sum_{\mathbf{k} \neq \mathbf{0}} \frac{\mathbf{k}}{k^{2}} e^{-\frac{k^{2}}{4\alpha^{2}}} \sum_{i}^{N} q_{i} \sin[\mathbf{k} \cdot (\mathbf{r}_{i} - \mathbf{r}_{j})]$$