






	1	2	3	4	5
1	5.8	5.6	6.2	6.5	5.8
2	6.5	7.2	8.0	7.2	6.5
3	7.2	8.0	10.0	8.0	7.2
4	8.0	10.0	10.0	10.0	8.0
5	7.2	9.0	10.0	9.0	8.1

(a) Baseline example: $r_{\text{boundary}} = r_{\text{forbidden}} = -1$, $r_{\text{target}} = 1$, $\gamma = 0.9$.




	1	2	3	4	5
1	0.0	0.0	0.0	0.0	0.0
2	0.0	0.0	0.0	0.0	0.1
3	0.0	0.0	2.0	0.1	0.1
4	0.0	2.0	2.0	2.0	0.2
5	0.0	1.0	2.0	1.0	0.5

(b) The discount rate is changed to $\gamma = 0.5$. The other parameters are the same as those in (a).



	1	2	3	4	5
1	0.0	0.0	0.0	0.0	0.0
2	0.0	0.0	0.0	0.0	0.0
3	0.0	0.0	1.0	0.0	0.0
4	0.0	1.0	1.0	1.0	0.0
5	0.0	0.0	1.0	0.0	0.0

(c) The discount rate is changed to $\gamma = 0$. The other parameters are the same as those in (a).



	1	2	3	4	5
1	3.5	3.9	4.3	4.8	5.3
2	3.1	3.5	4.8	5.3	5.9
3	2.8	2.5	10.0	5.9	6.6
4	2.5	10.0	10.0	10.0	7.3
5	2.3	9.0	10.0	9.0	8.1

(d) $r_{\text{forbidden}}$ is changed from -1 to -10 . The other parameters are the same as those in (a).