$D_1(2420)^0$ 


$$I(J^P) = \frac{1}{2}(1^+)$$
I needs confirmation.

## $D_1(2420)^0$ MASS

The fit includes  $D^\pm$ ,  $D^0$ ,  $D_s^\pm$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_s^{*\pm}$ ,  $D_1(2420)^0$ ,  $D_2^*(2460)^0$ , and  $D_{s1}(2536)^\pm$  mass and mass difference measurements.

VALUE (MeV)	EVTS	DOCUMENT ID		TECN	COMMENT
2420.8±0.5 OUR	FIT Error incl	udes scale factor o	f 1.3.		
2420.5±0.6 OUR	<b>AVERAGE</b> Er	ror includes scale f	actor	of 1.3.	See the ideogram below.
$2419.6\!\pm\!0.1\!\pm\!0.7$	210k	AAIJ	<b>13</b> CC	LHCB	$pp \rightarrow D^{*+}\pi^{-}X$
$2423.1\!\pm\!1.5\!+\!0.4\\-1.0$	2.7k	<sup>1</sup> ABRAMOWIC	Z13	ZEUS	$e^{\pm} p \rightarrow D^{(*)+} \pi^- X$
$2420.1\!\pm\!0.1\!\pm\!0.8$	103k	DEL-AMO-SA.			$e^+e^- \rightarrow D^{*+}\pi^- X$
$2426$ $\pm 3$ $\pm 1$	151	ABE	05A	BELL	$B^- \to D^0 \pi^+ \pi^- \pi^-$
$2421.4\!\pm\!1.5\!\pm\!0.9$		<sup>2</sup> ABE	<b>04</b> D	BELL	$B^- \rightarrow D^{*+}\pi^-\pi^-$
$2421 \begin{array}{cc} +1 \\ -2 \end{array} \pm 2$	286	AVERY	<b>94</b> C	CLE2	$e^+e^- \rightarrow D^{*+}\pi^-X$
$2422$ $\pm 2$ $\pm 2$	51	FRABETTI	<b>94</b> B	E687	$\gamma\mathrm{Be}  o D^{*+}\pi^-\mathrm{X}$
$2428$ $\pm 3$ $\pm 2$	279	AVERY	90		$e^+e^- \rightarrow D^{*+}\pi^-X$
$2414 \ \pm 2 \ \pm 5$	171	ALBRECHT	89н	ARG	$e^+e^- \rightarrow D^{*+}\pi^-X$
$2428$ $\pm 8$ $\pm 5$	171	ANJOS	89C	TPS	$\gamma N \rightarrow D^{*+}\pi^{-}X$
$\bullet$ $\bullet$ We do not	use the following	g data for averages	s, fits,	limits, e	etc. • • •
$2420.5\!\pm\!2.1\!\pm\!0.9$	$3110\pm340$	<sup>3</sup> CHEKANOV	09	ZEUS	$e^{\pm} p \rightarrow D^{*+} \pi^{-} X$
$2421.7\!\pm\!0.7\!\pm\!0.6$	7.5k	ABULENCIA	06A	CDF	1900 $p\overline{p} \rightarrow D^{*+}\pi^{-}X$
$2425$ $\pm 3$	235	<sup>4</sup> ABREU	98M	DLPH	$e^+e^-$
WEIGHTE	D AVERAGE				

WEIGHTED AVERAGE 2420.5±0.6 (Error scaled by 1.3)



 $D_1(2420)^0$  mass (MeV)

### $m_{D_1^0} - m_{D^{*+}}$

The fit includes  $D^{\pm}$ ,  $D^{0}$ ,  $D_{s}^{\pm}$ ,  $D^{*\pm}$ ,  $D^{*0}$ ,  $D_{s}^{*\pm}$ ,  $D_{1}(2420)^{0}$ ,  $D_{2}^{*}(2460)^{0}$ , and  $D_{\rm S1}(2536)^{\pm}$  mass and mass difference measurements.

	VALUE	<u>EVTS</u>	DOCUMENT ID		TECN	COMMENT
	410.6±0.5 OUR F	<b>IT</b> Error includ	les scale factor of	1.3.		
411.5±0.8 OUR AVERAGE						
	$410.2\!\pm\!2.1\!\pm\!0.9$	$3110 \pm 340$	CHEKANOV	09	ZEUS	$e^{\pm} p \rightarrow D^{*+} \pi^{-} X$
	$411.7\!\pm\!0.7\!\pm\!0.4$	7.5k	ABULENCIA	06A	CDF	1900 $p\overline{p} \rightarrow D^{*+}\pi^{-}X$

#### $D_1(2420)^0$ WIDTH

VALUE (MeV)	EVTS	DOCUMENT ID	TECN	COMMENT
31.7± 2.5 OUR AVER	AGE Erro	or includes scale fac	tor of 3.5. S	ee the ideogram below.
$35.2 \pm 0.4 \pm 0.9$	210k	AAIJ	13CC LHCB	$pp \rightarrow D^{*+}\pi^{-}X$
$38.8\pm 5.0^{+}_{-} \begin{array}{c} 1.9 \\ 5.4 \end{array}$	2.7k	<sup>1</sup> ABRAMOWICZ	Z13 ZEUS	$e^{\pm} p \rightarrow D^{(*)+} \pi^- X$
$31.4 \pm \ 0.5 \pm \ 1.3$	103k	DEL-AMO-SA	10P BABR	$e^+e^- \rightarrow D^{*+}\pi^- X$
$20.0 \pm \ 1.7 \pm \ 1.3$	7.5k			1900 $p\overline{p} \rightarrow D^{*+}\pi^{-}X$
$24~\pm~7~\pm~8$	151	ABE	05A BELL	$B^- \rightarrow D^0 \pi^+ \pi^- \pi^-$
$23.7 \pm \ 2.7 \pm \ 4.0$		<sup>2</sup> ABE	04D BELL	$B^- \rightarrow D^{*+}\pi^-\pi^-$
$20 \begin{array}{cccc} + & 6 \\ - & 5 \end{array} \pm 3$	286	AVERY	94C CLE2	$e^+e^- \rightarrow D^{*+}\pi^-X$
$15 \pm 8 \pm 4$	51	FRABETTI	94B <b>E687</b>	$\gamma\mathrm{Be}  o D^{*+}\pi^-\mathrm{X}$
$23 \begin{array}{cccccccccccccccccccccccccccccccccccc$	279	AVERY	90 CLEO	$e^+e^- \rightarrow D^{*+}\pi^-X$
$13 \pm 6 \begin{array}{l} +10 \\ -5 \end{array}$	171	ALBRECHT	89н ARG	$e^+e^- \rightarrow D^{*+}\pi^-X$

<sup>• • •</sup> We do not use the following data for averages, fits, limits, etc. • • •

53.2
$$\pm$$
 7.2 $^+$  3.3 3110 $\pm$ 340 CHEKANOV 09 ZEUS  $e^{\pm} p \rightarrow D^{*+} \pi^- X$  58  $\pm$ 14  $\pm$ 10 171 ANJOS 89C TPS  $\gamma N \rightarrow D^{*+} \pi^- X$ 


 $<sup>^1</sup>$  From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions. and  ${\rm A}_{D_2}$  fixed to the theoretical prediction of -1.

<sup>&</sup>lt;sup>2</sup> Fit includes the contribution from  $D_1^*(2430)^0$ .

 $<sup>^3</sup>$  Calculated using the mass difference  $\mathit{m}(\mathit{D}_1^0) - \mathit{m}(\mathit{D}^{*+})_{PDG}$  reported below and  $m(D^{*+})_{PDG}=$  2010.27  $\pm$  0.17 MeV. The 0.17 MeV uncertainty of the PDG mass value should be added to the experimental uncertainty of 0.9 MeV.

<sup>&</sup>lt;sup>4</sup> No systematic error given.

 $<sup>^1</sup>$  From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions. and  ${\rm A}_{D_2}$  fixed to the theoretical prediction of -1. <sup>2</sup> Fit includes the contribution from  $D_1^*(2430)^0$ .



# $D_1(2420)^0$ DECAY MODES

 $\overline{D}_1(2420)^0$  modes are charge conjugates of modes below.

	Mode	Fraction $(\Gamma_i/\Gamma)$
$\overline{\Gamma_1}$	$D^*(2010)^+\pi^-$	seen
$\Gamma_2$	$D^0\pi^+\pi^-$	seen
$\Gamma_3$	$D^0  ho^0$	
$\Gamma_{\Delta}$	$D^0 f_0(500)$	
Γ <sub>5</sub>	$D_0^*(2400)^+\pi^-$	
	$D^{+}\pi^{-} \ D^{*0}\pi^{+}\pi^{-}$	not seen
Γ <sub>7</sub>	$D^{*0} \pi^+ \pi^-$	not seen

# $D_1(2420)^0$ BRANCHING RATIOS

$\Gamma(D^*(2010)^+\pi^-)/\Gamma_{total}$			$\Gamma_1/\Gamma$
VALUE	DOCUMENT ID	TECN	COMMENT
seen		_	$e^+e^- \rightarrow D^{*+}\pi^-X$
seen	AVERY 90	CLEO	$e^+e^-  ightarrow D^{*+}\pi^-X$
seen	ALBRECHT 89	н ARG	$e^+e^-  ightarrow D^*\pi^-X$
seen	ANJOS 89	c TPS	$\gamma N \rightarrow D^{*+} \pi^{-} X$

$\Gamma(D^+\pi^-)/\Gamma(D^*(2010)^+\pi^-)$					$\Gamma_6/\Gamma_1$
VALUE	CL%	DOCUMENT ID		TECN	COMMENT
<0.24	90	AVERY	90	CLEO	$e^+e^- \rightarrow D^+\pi^-X$

### $D_1(2420)^0$ POLARIZATION AMPLITUDE $A_{D_1}$

A polarization amplitude  $A_{D_1}$  is a parameter that depends on the initial polarization of the  $D_1$  and is sensitive to a possible S-wave contribution to its decay. For  $D_1$  decays the helicity angle,  $\theta_h$ , distribution varies like  $1+A_{D_1}\cos^2\theta_h$ , where  $\theta_h$  is the angle in the  $D^*$  rest frame between the two pions emitted by the  $D_1 \to D^*\pi$  and the  $D^* \to D\pi$ .

Unpolarized  $D_1$  decaying purely via D-wave is predicted to give  $A_{D_1}=3$ .

<u>VALUE</u>	<u>EVTS</u>	DOCUMENT ID TECN COMMENT				
5.73±0.25 OUR AVERAGE						
$7.8 \begin{array}{c} +6.7 \\ -2.7 \end{array} \begin{array}{c} +4.6 \\ -1.8 \end{array}$	2.7k	<sup>1</sup> ABRAMOWICZ13 ZEUS $e^{\pm} p \rightarrow D^{(*)} + \pi^{-} X$				
$5.72 \pm 0.25$	103k	DEL-AMO-SA10P BABR $e^+e^-  o D^{*+}\pi^- X$				
$5.9 \begin{array}{c} +3.0 \\ -1.7 \end{array} \begin{array}{c} +2.4 \\ -1.0 \end{array}$		CHEKANOV 09 ZEUS $e^{\pm} p \rightarrow D^{*+} \pi^{-} X$				
• • We do not use the following data for averages, fits, limits, etc. • •						
$3.30\pm0.48$ $3.8\ \pm0.6\ \pm0.8$	210k	$^2$ AAIJ 13CC LHCB $pp  o D^{*+}\pi^- X$ 3 AUBERT 09Y BABR $B^+  o D_1^0 \ell^+ \nu_\ell$				
$2.74 ^{igoplus 1.40}_{-0.93}$		4 AVERY 94C CLE2 $e^+e^- \to D^{*+}\pi^- X$				

 $<sup>^1</sup>$  From the combined fit of the  $M(D^+\pi^-)$  and  $M(D^{*+}\pi^-)$  distributions. and  ${\rm A}_{D_2}$  fixed to the theoretical prediction of -1. A pure D-wave not excluded although some S-wave mixing possible.

### $D_1(2420)^0$ REFERENCES

<sup>&</sup>lt;sup>2</sup> Systematic uncertainty not estimated. Resonance parameters fixed.

<sup>&</sup>lt;sup>3</sup> Assuming  $\Gamma(\Upsilon(4S) \to B^+B^-) / \Gamma(\Upsilon(4S) \to B^0\overline{B}{}^0) = 1.065 \pm 0.026$  and equal partial widths and helicity angle distributions for charged and neutral  $D_1$  mesons.

<sup>&</sup>lt;sup>4</sup> Systematic uncertainties not estimated.