$$I^{G}(J^{PC}) = 0^{+}(0^{-+})$$

#### OMITTED FROM SUMMARY TABLE

Quantum numbers shown are quark-model predictions.

### $\eta_b(2S)$ MASS

9999.0 $\pm 3.5^{+2.8}_{-1.9}$  $^{1}\,\mathrm{MIZUK}$ 26k BELL

• • We do not use the following data for averages, fits, limits, etc.

9974.6 $\pm$ 2.3 $\pm$ 2.1 11  $\pm$  4 2,3 DOBBS

12

 $\Upsilon(2S) 
ightarrow \gamma$  hadrons

### $m_{\Upsilon(2S)}-m_{\eta_b(2S)}$

VALUE (MeV) TECN COMMENT  $24.3\pm3.5^{+2.8}_{-1.9}$ <sup>4</sup> MIZUK 26k 12 BELL

• • • We do not use the following data for averages, fits, limits, etc. •

 $11 \pm 4$  5,6 DOBBS

12

 $\Upsilon(2S) 
ightarrow \gamma$  hadrons

Created: 5/30/2017 17:21

# $\eta_b(2S)$ WIDTH

VALUE (MeV)	CL%	DOCUMENT ID		TECN	COMMENT
<24	90	MIZUK	12	BELL	$e^+e^-  ightarrow \gamma \pi^+\pi^-$ hadrons

## $\eta_b(2S)$ DECAY MODES

	Mode	Fraction $(\Gamma_i/\Gamma)$
Γ <sub>1</sub>	hadrons	seen

 $<sup>^{1}</sup>$  Assuming  $\Gamma_{\eta_{b}(2S)}=$  4.9 MeV. Not independent of the corresponding mass difference measurement.

2 Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

 $<sup>^3</sup>$ Assuming  $\Gamma_{\eta_h(2S)}=5$  MeV. Not independent of the corresponding mass difference

 $<sup>^4</sup>$  Assuming  $\Gamma_{\eta_b(2S)}=$  4.9 MeV. Not independent of the corresponding mass measure-

Obtained by analyzing CLEO III data but not authored by the CLEO Collaboration.

<sup>&</sup>lt;sup>6</sup> Assuming  $\Gamma_{\eta_h(2S)}=5$  MeV. Not independent of the corresponding mass measurement.

## $\eta_b(2S)$ BRANCHING RATIOS

 $\Gamma(\text{hadrons})/\Gamma_{\text{total}}$   $\Gamma_1/\Gamma$ 

seen  $^{7}$  DOBBS  $^{12}$   $\gamma(2S) 
ightarrow \gamma$  hadrons

 $^{7}\,\mathrm{Obtained}$  by analyzing CLEO III data but not authored by the CLEO Collaboration.

## $\eta_b(2S)$ REFERENCES

DOBBS 12 PRL 109 082001 S. Dot MIZUK 12 PRL 109 232002 R. Miz

S. Dobbs *et al.* R. Mizuk *et al.* 

(BELLE Collab.)

Created: 5/30/2017 17:21