

Possible Choices of Complex Co-ordinates for a Surface


Moduli Spaces of Riemann Surfaces

Zhaoshen Zhai


Graduate mentor: Kaleb Ruscitti

April 23, 2023


How many complex coordinates on \mathbb{R}^2 ?




How many complex coordinates on \mathbb{R}^2 ?



How many complex coordinates on \mathbb{R}^2 ?



How many complex coordinates on \mathbb{R}^2 ?



Refined question


How many complex coordinates on \mathbb{R}^2 ?

Refined question


~~How many complex coordinates on \mathbb{R}^2 ?~~

How many complex coordinates on \mathbb{R}^2 *up to biholomorphism*?

\mathbb{C} and \mathbb{H}



\mathbb{C} and \mathbb{H}



Riemann surfaces


What about for manifolds?

Riemann surfaces

What about for manifolds?

Definition

A Riemann surface is a connected 1-dimensional complex manifold.



Compact Riemann surfaces

Theorem (Classification)

Every compact Riemann surface is classified by its genus.

Compact Riemann surfaces

Theorem (Classification)

Every compact Riemann surface is classified by its genus.



Topological torus


Theorem

Every torus is a quotient $\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2)$ for some linearly independent $\omega_1, \omega_2 \in \mathbb{C}$.

Topological torus

Theorem


Every torus is a quotient $\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2)$ for some linearly independent $\omega_1, \omega_2 \in \mathbb{C}$.



Topological torus

Theorem

Every torus is a quotient $\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2)$ for some linearly independent $\omega_1, \omega_2 \in \mathbb{C}$.



Theorem

There is, up to homeomorphism, only one topological torus.


Questions

- *Up to biholomorphism, in how many ways can we give the torus complex coordinates?*

Questions

- *Up to biholomorphism, in how many ways can we give the torus complex coordinates?*
- *What is the moduli space of the torus?*

Complex tori



Complex tori

Theorem

Every complex tori is the quotient $X_\tau := \mathbb{C}/\Gamma$ where $\Gamma = \mathbb{Z} \oplus \mathbb{Z}\tau$ for some $\tau \in \mathbb{H}$.


Theorem

Every complex tori is the quotient $X_\tau := \mathbb{C}/\Gamma$ where $\Gamma = \mathbb{Z} \oplus \mathbb{Z}\tau$ for some $\tau \in \mathbb{H}$. Indeed, for all linearly independent $\omega_1, \omega_2 \in \mathbb{C}$,


$$\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2) \cong X_\tau$$

for $\tau := \omega_2/\omega_1 \in \mathbb{H}$.


Complex tori



Complex tori



Complex tori



Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Complex tori

Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Theorem

For any $\tau, \tau' \in \mathbb{H}$, the tori X_τ and $X_{\tau'}$ are biholomorphic iff there exists some $\gamma \in \mathrm{PSL}_2(\mathbb{Z})$ such that $\tau' = \gamma(\tau)$.

Complex tori

Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Theorem

For any $\tau, \tau' \in \mathbb{H}$, the tori X_τ and $X_{\tau'}$ are biholomorphic iff there exists some $\gamma \in \mathrm{PSL}_2(\mathbb{Z})$ such that $\tau' = \gamma(\tau)$.

Corollary

The moduli space of complex tori is $\mathbb{H} / \mathrm{PSL}_2(\mathbb{Z})$.

Theorem

There is a unique choice of complex coordinates on the sphere.

Riemann sphere

Theorem

There is a unique choice of complex coordinates on the sphere.

Proof. Hard!

Theorem

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! We used techniques from:

- Covering space theory: We defined the degree of a holomorphic map and proved that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.

Theorem

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! We used techniques from:

- Covering space theory: We defined the degree of a holomorphic map and proved that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.

Theorem

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! We used techniques from:

- Covering space theory: We defined the degree of a holomorphic map and proved that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.
- Sheaf cohomology: uhh something. ■

Thank you!

Thank you!