

Possible Choices of Complex Structures for a Surface


Moduli Spaces of Riemann Surfaces

Zhaoshen Zhai


Graduate mentor: Kaleb Ruscitti

April 28, 2023


Complex structures on \mathbb{R}^2




Complex structures on \mathbb{R}^2



Complex structures on \mathbb{R}^2



Complex structures on \mathbb{R}^2



Refined question


How many complex structures on \mathbb{R}^2 ?

Refined question


~~How many complex structures on \mathbb{R}^2 ?~~

How many complex structures on \mathbb{R}^2 *up to biholomorphism*?

\mathbb{C} and \mathbb{H}




\mathbb{C} and \mathbb{H}



Riemann surfaces

Definition

A Riemann surface is a surface equipped with a complex structure.



Compact surfaces

Theorem (Classification of Surfaces)

For each natural number g , there exists exactly one compact orientable surface up to homeomorphism.

Compact surfaces

Theorem (Classification of Surfaces)

For each natural number g , there exists exactly one compact orientable surface up to homeomorphism.



Topological torus


Theorem

Every torus is a quotient $\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2)$ for some \mathbb{R} -linearly independent $\omega_1, \omega_2 \in \mathbb{C}$.


Topological torus

Theorem

Every torus is a quotient $\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2)$ for some \mathbb{R} -linearly independent $\omega_1, \omega_2 \in \mathbb{C}$.



Complex tori




Theorem

Every complex torus is the quotient $X_\tau := \mathbb{C}/(\mathbb{Z} \oplus \mathbb{Z}\tau)$ for some $\tau \in \mathbb{H}$. Indeed, for all linearly independent $\omega_1, \omega_2 \in \mathbb{C}$,


$$\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2) \cong X_\tau$$

for $\tau := \omega_2/\omega_1 \in \mathbb{H}$.


Complex tori



Complex tori



Complex tori



Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Complex tori

Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Theorem

For any $\tau, \tau' \in \mathbb{H}$, the tori X_τ and $X_{\tau'}$ are biholomorphic iff there exists some $\gamma \in \mathrm{PSL}_2(\mathbb{Z})$ such that $\tau' = \gamma(\tau)$.

Complex tori

Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Theorem

For any $\tau, \tau' \in \mathbb{H}$, the tori X_τ and $X_{\tau'}$ are biholomorphic iff there exists some $\gamma \in \mathrm{PSL}_2(\mathbb{Z})$ such that $\tau' = \gamma(\tau)$.

Corollary

The moduli space of the torus is $\mathbb{H} / \mathrm{PSL}_2(\mathbb{Z})$.

Theorem (Uniformization)

There is a unique choice of complex structure on the sphere.

Riemann sphere

Theorem (Uniformization)

There is a unique choice of complex structure on the sphere.

Proof. Hard!

Riemann sphere

Theorem (Uniformization)

There is a unique choice of complex structure on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex structure. Let X be another sphere.

Theorem (Uniformization)

There is a unique choice of complex structure on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex structure. Let X be another sphere.

- Covering space theory: We showed that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.

Theorem (Uniformization)

There is a unique choice of complex structure on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex structure. Let X be another sphere.

- Covering space theory: We showed that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.

Theorem (Uniformization)

There is a unique choice of complex structure on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex structure. Let X be another sphere.

- Covering space theory: We showed that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.
- Čech cohomology: We used tools from differential forms and cohomology to prove the existence of such a meromorphic function.

Theorem (Uniformization)

There is a unique choice of complex structure on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex structure. Let X be another sphere.

- Covering space theory: We showed that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.
- Čech cohomology: We used tools from differential forms and cohomology to prove the existence of such a meromorphic function. ■

Thank you!

Thank you!