

Possible Choices of Complex Co-ordinates for a Surface


Moduli Spaces of Riemann Surfaces

Zhaoshen Zhai


Graduate mentor: Kaleb Ruscitti

April 26, 2023


How many complex coordinates on \mathbb{R}^2 ?




How many complex coordinates on \mathbb{R}^2 ?



How many complex coordinates on \mathbb{R}^2 ?



How many complex coordinates on \mathbb{R}^2 ?



Refined question


How many complex coordinates on \mathbb{R}^2 ?

Refined question


~~How many complex coordinates on \mathbb{R}^2 ?~~

How many complex coordinates on \mathbb{R}^2 *up to biholomorphism*?

\mathbb{C} and \mathbb{H}




\mathbb{C} and \mathbb{H}




Surfaces

Recall that a surface is a connected 2-dimensional real manifold.



Surfaces

Recall that a surface is a connected 2-dimensional real manifold.



Definition

A Riemann surface is a surface with a choice of complex structure.

Compact surfaces

Theorem (Classification of Surfaces)

For each natural number g , there exists exactly one compact orientable surface up to homeomorphism. We call g the genus of the surface.

Compact surfaces

Theorem (Classification of Surfaces)

For each natural number g , there exists exactly one compact orientable surface up to homeomorphism. We call g the genus of the surface.



Topological torus


Theorem

Every torus is a quotient $\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2)$ for some \mathbb{R} -linearly independent $\omega_1, \omega_2 \in \mathbb{C}$.


Topological torus

Theorem

Every torus is a quotient $\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2)$ for some \mathbb{R} -linearly independent $\omega_1, \omega_2 \in \mathbb{C}$.



Complex tori




Theorem

Every complex tori is the quotient $X_\tau := \mathbb{C}/(\mathbb{Z} \oplus \mathbb{Z}\tau)$ for some $\tau \in \mathbb{H}$. Indeed, for all linearly independent $\omega_1, \omega_2 \in \mathbb{C}$,


$$\mathbb{C}/(\mathbb{Z}\omega_1 \oplus \mathbb{Z}\omega_2) \cong X_\tau$$

for $\tau := \omega_2/\omega_1 \in \mathbb{H}$.


Complex tori



Complex tori



Complex tori



Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Complex tori

Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Theorem

For any $\tau, \tau' \in \mathbb{H}$, the tori X_τ and $X_{\tau'}$ are biholomorphic iff there exists some $\gamma \in \mathrm{PSL}_2(\mathbb{Z})$ such that $\tau' = \gamma(\tau)$.

Complex tori

Definition

The modular group $\mathrm{PSL}_2(\mathbb{Z})$ is the group of functions $\gamma : \mathbb{H} \rightarrow \mathbb{H}$ mapping

$$\tau \mapsto \frac{a\tau + b}{c\tau + d}$$

for some $a, b, c, d \in \mathbb{Z}$ with $ad - bc = 1$.

Theorem

For any $\tau, \tau' \in \mathbb{H}$, the tori X_τ and $X_{\tau'}$ are biholomorphic iff there exists some $\gamma \in \mathrm{PSL}_2(\mathbb{Z})$ such that $\tau' = \gamma(\tau)$.

Corollary

The moduli space of complex tori is $\mathbb{H} / \mathrm{PSL}_2(\mathbb{Z})$.

Riemann sphere

Theorem (Uniformization)

There is a unique choice of complex coordinates on the sphere.

Riemann sphere

Theorem (Uniformization)

There is a unique choice of complex coordinates on the sphere.

Proof. Hard!

Theorem (Uniformization)

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex coordinate. Let X be another sphere.

Theorem (Uniformization)

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex coordinate. Let X be another sphere.

- Covering space theory: We defined the degree of a holomorphic map and proved that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.

Theorem (Uniformization)

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex coordinate. Let X be another sphere.

- Covering space theory: We defined the degree of a holomorphic map and proved that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.

Theorem (Uniformization)

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex coordinate. Let X be another sphere.

- Covering space theory: We defined the degree of a holomorphic map and proved that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.
- Čech cohomology: We used tools from differential forms and cohomology to prove the existence of such a meromorphic function.

Theorem (Uniformization)

There is a unique choice of complex coordinates on the sphere.

Proof. Hard! Fix a sphere $\hat{\mathbb{C}}$, which is equipped with a particular complex coordinate. Let X be another sphere.

- Covering space theory: We defined the degree of a holomorphic map and proved that if X is compact and if there exists a meromorphic function on X with a single simple pole, then $X \cong \hat{\mathbb{C}}$.
- Analytic continuation: We attempted to find such a meromorphic function, but our techniques weren't powerful enough.
- Čech cohomology: We used tools from differential forms and cohomology to prove the existence of such a meromorphic function. ■

Thank you!

Thank you!