

# Tree-like graphings of countable Borel equivalence relations

An exposition to

*Tree-like graphings, wallings, and median graphings of equivalence relations*

by Ruiyuan Chen, Antoine Poulin, Ran Tao, and Anush Tserunyan

Zhaoshen Zhai

October 1, 2024

# Countable Borel equivalence relations

## Definition

A *countable Borel equivalence relation (CBER)* on a standard Borel space  $X$  is a Borel equivalence relation  $E \subseteq X^2$  such that each  $E$ -class is countable.

# Countable Borel equivalence relations

## Definition

A *countable Borel equivalence relation (CBER)* on a standard Borel space  $X$  is a Borel equivalence relation  $E \subseteq X^2$  such that each  $E$ -class is countable.

## Example

Any Borel action  $\Gamma \curvearrowright X$  of a countable (discrete) group on a standard Borel space induces its *orbit equivalence relation*  $E_\Gamma^X$ , which is a CBER.



# Countable Borel equivalence relations

## Definition

A *countable Borel equivalence relation* (CBER) on a standard Borel space  $X$  is a Borel equivalence relation  $E \subseteq X^2$  such that each  $E$ -class is countable.

## Example

Any Borel action  $\Gamma \curvearrowright X$  of a countable (discrete) group on a standard Borel space induces its *orbit equivalence relation*  $E_\Gamma^X$ , which is a CBER.



# Smooth and hyperfinite CBERs

## Example (Smooth)

- Identity relation = on a standard Borel space, say  $\mathbb{R}$  or  $2^{\mathbb{N}}$ .

# Smooth and hyperfinite CBERs

## Example (Smooth)

- Identity relation = on a standard Borel space, say  $\mathbb{R}$  or  $2^{\mathbb{N}}$ .
- $\mathbb{Z}$ -coset equivalence on  $\mathbb{R}$ :  $x E_{\mathbb{Z}}^{\mathbb{R}} y$  iff  $x - y \in \mathbb{Z}$ .

# Smooth and hyperfinite CBERs

## Example (Smooth)

- Identity relation = on a standard Borel space, say  $\mathbb{R}$  or  $2^{\mathbb{N}}$ .
- $\mathbb{Z}$ -coset equivalence on  $\mathbb{R}$ :  $x E_{\mathbb{Z}}^{\mathbb{R}} y$  iff  $x - y \in \mathbb{Z}$ .

## Example (Hyperfinite)

$E_0$  on  $2^{\mathbb{N}}$ , where  $x E_0 y$  iff  $\exists n \in \mathbb{N}, \forall m \geq n : x_m = y_m$ .

# Smooth and hyperfinite CBERs

## Example (Smooth)

- Identity relation  $=$  on a standard Borel space, say  $\mathbb{R}$  or  $2^{\mathbb{N}}$ .
- $\mathbb{Z}$ -coset equivalence on  $\mathbb{R}$ :  $x E_{\mathbb{Z}}^{\mathbb{R}} y$  iff  $x - y \in \mathbb{Z}$ .

## Example (Hyperfinite)

$E_0$  on  $2^{\mathbb{N}}$ , where  $x E_0 y$  iff  $\exists n \in \mathbb{N}, \forall m \geq n : x_m = y_m$ .

This CBER is *hyperfinite*:  $E_0 = \bigcup_n F_n$  for an increasing sequence  $F_0 \subseteq F_1 \cdots$  of finite Borel equivalence relations:



# Smooth and hyperfinite CBERs

## Example (Smooth)

- Identity relation = on a standard Borel space, say  $\mathbb{R}$  or  $2^{\mathbb{N}}$ .
- $\mathbb{Z}$ -coset equivalence on  $\mathbb{R}$ :  $x E_{\mathbb{Z}}^{\mathbb{R}} y$  iff  $x - y \in \mathbb{Z}$ .

## Example (Hyperfinite)

$E_0$  on  $2^{\mathbb{N}}$ , where  $x E_0 y$  iff  $\exists n \in \mathbb{N}, \forall m \geq n : x_m = y_m$ .

This CBER is *hyperfinite*:  $E_0 = \bigcup_n F_n$  for an increasing sequence  $F_0 \subseteq F_1 \cdots$  of finite Borel equivalence relations:

$$x F_n y \quad \leftrightarrow \quad \forall m \geq n : x_m = y_m.$$

# Smooth and hyperfinite CBERs

## Example (Smooth)

- Identity relation = on a standard Borel space, say  $\mathbb{R}$  or  $2^{\mathbb{N}}$ .
- $\mathbb{Z}$ -coset equivalence on  $\mathbb{R}$ :  $x E_{\mathbb{Z}}^{\mathbb{R}} y$  iff  $x - y \in \mathbb{Z}$ .

## Example (Hyperfinite)

$E_0$  on  $2^{\mathbb{N}}$ , where  $x E_0 y$  iff  $\exists n \in \mathbb{N}, \forall m \geq n : x_m = y_m$ .

This CBER is *hyperfinite*:  $E_0 = \bigcup_n F_n$  for an increasing sequence  $F_0 \subseteq F_1 \subseteq \dots$  of finite Borel equivalence relations:

$$x F_n y \quad \leftrightarrow \quad \forall m \geq n : x_m = y_m.$$

## Theorem (Slaman-Steel, Weiss)

Let  $E$  be a CBER on a standard Borel space  $X$ . TFAE:

1.  $E$  is hyperfinite.  $E = \bigcup_n F_n$  where  $F_0 \subseteq F_1 \subseteq \dots$  are FBERs.
2.  $E$  is induced by a Borel  $\mathbb{Z}$ -action.  $E = E_{\mathbb{Z}}^X$  for some  $\mathbb{Z} \curvearrowright X$ .



# Smooth and Hyperfinite CBERs

## Example (Smooth)

- Identity relation = on a standard Borel space, say  $\mathbb{R}$  or  $2^{\mathbb{N}}$ .
- $\mathbb{Z}$ -coset equivalence on  $\mathbb{R}$ :  $x E_{\mathbb{Z}}^{\mathbb{R}} y$  iff  $x - y \in \mathbb{Z}$ .

## Example (Hyperfinite)

$E_0$  on  $2^{\mathbb{N}}$ , where  $x E_0 y$  iff  $\exists n \in \mathbb{N}, \forall m \geq n : x_m = y_m$ .



# Graphing of a CBER





## Definition

A *graphing* of a CBER  $E$  on  $X$  is a Borel graph  $G \subseteq X^2$  whose connected relation is  $E$ , i.e.,  $xEy \leftrightarrow xG \cdots Gy$  for all  $x, y \in X$ .

# Graphing of a CBER

## Definition





A *graphing* of a CBER  $E$  on  $X$  is a Borel graph  $G \subseteq X^2$  whose connected relation is  $E$ , i.e.,  $xEy \leftrightarrow xG \cdots Gy$  for all  $x, y \in X$ .



# Treeings and treeability

## Definition


A *treeing* of a CBER  $E$  is an acyclic graphing, and a CBER  $E$  is said to be *treeable* if it admits a treeing.



# Treeable CBERs

## Example (Free Actions)


Free actions of a free group  $F_r \curvearrowright X$ .



# Treeable CBERs

## Example (Free Actions)

Free actions of a free group  $F_r \curvearrowright X$ .



## Theorem (JKL02)


*Free actions of virtually-free groups are treeable.*



# Treeable CBERs

## Example (Free Actions)

Free actions of a free group  $F_r \curvearrowright X$ .



## Theorem (JKL02)

*Free actions of virtually-free groups are treeable.*


## Theorem (GdlH90)

*Every finitely-generated group whose Cayley graph is a quasi-tree is virtually-free, and hence treeable.*

# Treeable CBERs

## Example (Free Actions)

Free actions of a free group  $F_r \curvearrowright X$ .



## Theorem (JKL02)

*Free actions of virtually-free groups are treeable.*

## Theorem (GdlH90)

*Every finitely-generated group whose Cayley graph is a quasi-tree is virtually-free, and hence treeable.*

## Question (Robin Tucker-Drob; 2015)

Is the class of treeable CBERs robust under quasi-isometries?

# Main result

Theorem (Chen, Poulin, Tao, Tserunyan; 2023+)


*If a CBER  $E$  admits a locally-finite graphing such that each component is a quasi-tree, then  $E$  is treeable.*

# Main result

Theorem (Chen, Poulin, Tao, Tserunyan; 2023+)

*If a CBER  $E$  admits a locally-finite graphing such that each component is a quasi-tree, then  $E$  is treeable.*

Two metric spaces  $X, Y$  are *quasi-isometric* if they are isometric up to a bounded multiplicative and additive error;  $X$  is a *quasi-tree* if it is quasi-isometric to a tree.



# Game plan

Quasi-treeing

Treeing

# Game plan

Quasi-treeing



Quasi-tree

Treeing




# Game plan

Quasi-treeing

Treeing

Quasi-tree

Finitely-sep.  
family of cuts



# Game plan

Quasi-treeing

Treeing

Quasi-tree

Finitely-sep.  
family of cuts





# Game plan


Quasi-treeing

Treeing

Quasi-tree

Finitely-sep.  
family of cuts

Median graph w/  
finite hyperplanes



# Game plan

Quasi-treeing


Treeing

Quasi-tree



Finitely-sep.  
family of cuts

Median graph w/  
finite hyperplanes



'Canonical'  
spanning tree



# Game plan



# Game plan



# Finitely-separating cuts

## Definition

A *cut* in a connected locally-finite graph  $(X, G)$  is a connected co-connected subset  $H \subseteq X$  with finite boundary.



# Finitely-separating cuts

## Definition

A *cut* in a connected locally-finite graph  $(X, G)$  is a connected co-connected subset  $H \subseteq X$  with finite boundary.



Let  $\mathcal{H}$  be a family of cuts such that if  $H \in \mathcal{H}$ , then  $\neg H \in \mathcal{H}$ .

# Finitely-separating cuts

## Definition

A *cut* in a connected locally-finite graph  $(X, G)$  is a connected co-connected subset  $H \subseteq X$  with finite boundary.



Let  $\mathcal{H}$  be a family of cuts such that if  $H \in \mathcal{H}$ , then  $\neg H \in \mathcal{H}$ .


## Definition

Such a family  $\mathcal{H}$  is *finitely-separating* if for each  $x, y \in X$ , there are finitely-many  $H \in \mathcal{H}$  with  $x \in H \not\ni y$ .

# Orientations

## Definition

An *orientation* on  $\mathcal{H}$  is an upward-closed subset  $U \subseteq \mathcal{H}$  containing exactly one of  $H, \neg H$  for every  $H \in \mathcal{H}$ .






# Orientations

## Definition

An *orientation* on  $\mathcal{H}$  is an upward-closed subset  $U \subseteq \mathcal{H}$  containing exactly one of  $H, \neg H$  for every  $H \in \mathcal{H}$ .




We'll only consider the orientations that are *based*, in the sense that each  $H \in U$  contains a minimal  $H_0 \in U$ .

# The dual median graph

## Definition

A *median graph* is a connected graph  $(X, G)$  such that for each  $x, y, z \in X$ , the intersection  $[x, y] \cap [x, z] \cap [y, z]$  is a singleton, called the *median* of  $x, y, z$ , and is denoted by  $\langle x, y, z \rangle$ .



# The dual median graph

## Definition

A *median graph* is a connected graph  $(X, G)$  such that for each  $x, y, z \in X$ , the intersection  $[x, y] \cap [x, z] \cap [y, z]$  is a singleton, called the *median* of  $x, y, z$ , and is denoted by  $\langle x, y, z \rangle$ .



## Theorem (Sageev 95)

If  $\mathcal{H}$  is finitely-separating, then the graph  $\mathcal{M}(\mathcal{H})$ :

- Vertices are based orientations on  $\mathcal{H}$ ;
- Neighbors of  $U$  are  $U \triangle \{H, \neg H\}$  for each minimal  $H \in U \setminus \{-0\}$ ;

is a median graph.

# Ends of graphs

## Definition

The *end compactification* of a connected locally-finite  $(X, G)$  is the Stone space  $\hat{X}$  of the Boolean algebra  $\mathcal{H}_{\partial < \infty}(X)$ , whose non-principal ultrafilters are the *ends* of  $(X, G)$ .



# Ends of graphs

## Definition

The *end compactification* of a connected locally-finite  $(X, G)$  is the Stone space  $\hat{X}$  of the Boolean algebra  $\mathcal{H}_{\partial < \infty}(X)$ , whose non-principal ultrafilters are the *ends* of  $(X, G)$ .



## Definition

A family  $\mathcal{H}$  of cuts is *dense towards ends* of  $X$  if  $\mathcal{H}$  contains a neighborhood basis for every end in  $\hat{X}$ .

# Density towards ends for quasi-trees

## Lemma


*The connected locally-finite graphs in which  $\mathcal{H}_{\text{diam}(\partial) \leq R}$  is dense towards ends for some  $R < \infty$  is invariant under quasi-isometry.*

## Corollary



*If  $(X, G)$  is a locally-finite quasi-tree, then the family*

$$\mathcal{H} := \mathcal{H}_{\text{diam}(\partial) \leq R}(X) \cap \mathcal{H}_{\text{conn}}(X)$$

*of cuts is dense towards ends for some  $R < \infty$ .*



# Wrapping things up...



# The End

Thank you!