

NESTED COLLECTION OF CUTS

ZHAOSHEN ZHAI

Notation 0.1. Let X be a set. For any family $\mathcal{C} \subseteq 2^X$ closed under the complement operation $\neg : 2^X \rightarrow 2^X$, we let $\mathcal{C}^* := \mathcal{C} \setminus \{\emptyset, X\}$ denote the *non-trivial* elements in \mathcal{C} .

1. PRELIMINARIES

1.1. Ultrafilters and their bases. We recall some standard notions, mainly to establish notation.

Definition 1.1. A *filter* on X is a collection $\mathcal{F} \subseteq \mathcal{P}^*(X)$ that is closed-upwards and closed under pairwise intersections, i.e.

- (1) If $A \in \mathcal{F}$ and $B \supseteq A$, then $B \in \mathcal{F}$; and
- (2) If $A_1, A_2 \in \mathcal{F}$, then $A_1 \cap A_2 \in \mathcal{F}$.

Lemma 1.2. For any family $\mathcal{F}_0 \subseteq \mathcal{P}^*(X)$, the following are equivalent.

- (1) For all $F_1, F_2 \in \mathcal{F}_0$, there is some $F \in \mathcal{F}_0$ such that $F \subseteq F_1 \cap F_2$.
- (2) The collection $\mathcal{F} := \{A \subseteq X : A \supseteq F \text{ for some } F \in \mathcal{F}_0\} \subseteq \mathcal{P}^*(X)$ is a filter on X .

Proof. If (1) holds, then \mathcal{F} is upward closed since if $A \supseteq F$ for some $F \in \mathcal{F}_0$, then $B \supseteq F$ for any $B \supseteq A$. Moreover, if $A_i \supseteq F_i$ for $i = 1, 2$, then $A_1 \cap A_2 \supseteq F_1 \cap F_2 \supseteq F$ for some $F \in \mathcal{F}_0$ furnished by (1).

Conversely, let $F_1, F_2 \in \mathcal{F}_0$. Then $F_1, F_2 \in \mathcal{F}$, so $F_1 \cap F_2 \in \mathcal{F}$ by (2). Thus $F_1 \cap F_2 \supseteq F$ for some $F \in \mathcal{F}_0$, by definition of \mathcal{F} , as desired. ■

Definition 1.3. A *basis for a filter* on X is a family $\mathcal{F}_0 \subseteq \mathcal{P}^*(X)$ satisfying the equivalent definitions above, and we call the collection \mathcal{F} in (2) the *filter generated by \mathcal{F}_0* .

Definition 1.4. An *ultrafilter* on X is a \subseteq -maximal filter on X .

1.2. Nested cuts and \mathcal{T} .

Definition 1.5. A *cut* in X is a connected co-connected set $C \subseteq X$ with finite vertex boundary.

Definition 1.6. A family \mathcal{C} of cuts in X is *nested* if every pair $C, C' \in \mathcal{C}$ has an empty corner; i.e. if one of $C \cap C', \neg C \cap C', C \cap \neg C', \neg C \cap \neg C'$ is empty.

2. THE GRAPH $\mathcal{T}_{\mathcal{C}}$

Theorem 2.1. Let \mathcal{C} be a family of nested cuts in X . The graph $\mathcal{T}_{\mathcal{C}}$, whose

- Vertices are finitely-based ultrafilters on \mathcal{C} ;
- Neighbors of

is acyclic. Furthermore, if *something*, then $\mathcal{T}_{\mathcal{C}}$ is a tree.

Proof. h ■

3. WHAT GOES WRONG IF \mathcal{C} IS NON-NESTED?