WOI- The maximum enhopy principle Jaynes (1957)

Imagine, 7m don't kom fi'i,
yn only know the awage of some funchin
f(xi), that is

then what can we say about 1 P. 4? and the awage of any other factor & (w)? The maximum - entropy purishe: tall 10:14 that maximizes alts enhapsy gien the constrains L=-ZiPilogPi- \(\(\fi\) -yt (z', t(k;) b', -1) y'yt en selænde un pibliere FL = 0 -> solus fr Pi that

FPi maving antimaximiles entropy

$$\frac{\delta L}{\delta P_{i}} = -\log P_{i} - P_{i} \cdot \frac{1}{P_{i}} - \lambda - \lambda f f(\kappa_{i})$$

$$\log P_{i} = -(1 + \lambda + \lambda f f(\kappa_{i}))$$

$$P_{i} = e$$

$$\frac{\lambda}{\delta P_{i}} = \frac{-(1 + \lambda + \lambda f f(\kappa_{i}))}{e^{-(1 + \lambda + \lambda f f(\kappa_{i}))}}$$

$$\frac{\lambda}{\delta P_{i}} = \frac{-(1 + \lambda + \lambda f f(\kappa_{i}))}{e^{-(1 + \lambda + \lambda f f(\kappa_{i}))}}$$

$$\frac{\lambda}{\delta P_{i}} = \frac{-(1 + \lambda + \lambda f f(\kappa_{i}))}{e^{-(1 + \lambda + \lambda f f(\kappa_{i}))}}$$

$$\frac{\lambda}{\delta P_{i}} = \frac{-(1 + \lambda + \lambda f f(\kappa_{i}))}{e^{-(1 + \lambda + \lambda f f(\kappa_{i}))}}$$

$$\frac{\lambda}{\delta P_{i}} = \frac{-(1 + \lambda + \lambda f f(\kappa_{i}))}{e^{-(1 + \lambda + \lambda f f(\kappa_{i}))}}$$

$$\frac{\lambda}{\delta P_{i}} = \frac{-\lambda f f(\kappa_{i})}{e^{-(1 + \lambda + \lambda f f(\kappa_{i}))}}$$

$$\sum_{i} f(x_{i}) P_{i} = f = \frac{\sum_{i} f(x_{i}) e}{\sum_{i} e^{x_{i}} f(x_{i})}$$

1)
$$f = 0$$
 $P_C = \frac{1}{n}$ He original High.

If you knot no thy asme nothy , all events equally likely

$$H = -\sum_{i} \frac{1}{n} \log n$$

$$\frac{1}{2} = \log n$$

$$b_{x}(x) = \frac{\sum_{i} e_{x_{i}} y_{i}t}{-x_{i}y_{i}t}$$

an exponential distribution

ci) if
$$\neg m$$
 ken the $\neg m$

$$f(x) = (x - m)^{2}$$

$$f'(x - m)^{2} \Rightarrow \frac{1}{e^{k}f(n-m)^{2}} \Rightarrow a \text{ Normal dishilation}$$

| • | | |
|---|--|--|
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |