Descriptive Set Theory

MATH 592 · Scribed Course Notes · Part 2: Definable Subsets of Polish Spaces

Definitions Results
Examples Notes
green text marks a term or notation at the point where it is defined

Contents

1 Course Overview **1**

2 σ-Algebras and Borel Sets **1**
2.1 σ-Algebras, Measurable Spaces and Borel Sets 2
2.2 Stratification of Borel Sets into a Hierarchy 3
2.3 The Classes $\mathcal{B}$, $\mathbf{\Sigma}^0_\xi$, $\mathbf{\Pi}^0_\xi$, $\mathbf{\Delta}^0_\xi$ 7
2.4 Universal Sets and Diagonalization 9

3 Change of Topology **13**
3.1 Turning Borel Sets into Clopen Sets 13
3.2 Making Closed-Graph Functions Continuous for Free 16

4 Analytic Sets **18**
4.1 Definition and Closure Properties 18
4.2 Universal Sets for Analytic / Coanalytic Sets 19
4.3 Analytic Separation 20
4.4 Souslin's Operation $\mathcal{A}$ 22

5 Uniformization Theorems **26**
5.1 Projections and Compactness 26
5.2 The Luzin–Souslin Theorem 27
5.3 The Luzin–Novikov Theorem and Countable Uniformization 28
5.4 Countable Borel Equivalence Relations 29
5.5 Further Uniformization Theorems 32

6 Standard Borel Spaces **33**
6.1 The Borel Isomorphism Theorem 33
6.2 The Effros Borel Space 35

Source manuscript page 1

1 Course Overview

Professor: Anush Tserunyan
Scribe: Yitian Zhang
Email: yit dot zha3 at gmail dot com

Note 1. The abbreviations used in the manuscript have been expanded in this edition.

Note 2. Text set in blue (or red) reproduces the scribe's own annotations, written in blue (or red) ink in the manuscript.

Note 3. The marker "Source manuscript page n" indicates where page n of the handwritten manuscript (Part 2) begins.

Note 4. Following the colour convention of the lecturer's notes, a term or a piece of notation is set in green at the point where it is defined, e.g. "a set is analytic if ..." or "$\mathcal{B}(X) := \dots$". Italics are reserved for ordinary emphasis.

Note 5. Results are numbered within sections, and a reference such as Theorem 3.1 is a clickable link to the result it cites; the number is set in this colour so that links are recognizable. Section references like §2.4 are links too.

Note 6. Results from Part 1 are cited by their number in the Part 1 edition of these notes, e.g. Part 1, Theorem 2.9 (Alexandrov's theorem). These are links into the file `Math592_Part1_Polish_Spaces.pdf`, which work when the two PDF files are kept in the same folder. Homework problems ("HW 3, Q2") are not part of the notes and are cited by name only.

Note 7. The manuscript ends with the title of the next section, *Projective Hierarchy* (manuscript p. 62), which has no content yet; it will be added once it is written.

Relevant readings as prerequisites.

- Analysis 1: § Topology and definitions.
- Analysis 2: § Metric spaces.
- Folland, *Real Analysis*, Chapter 4.

Source manuscript page 2

Introduction. In Part 1, we discussed the properties of Polish spaces and their characterizations. In this part, we are mainly concerned with the properties of subsets of Polish spaces, and of functions between Polish spaces, putting our focus on the so-called "definability".

We begin our discussion with Borel measurability and Borel (measurable) sets and functions.

Source manuscript page 3

2 σ-Algebras and Borel Sets

2.1 σ-Algebras, Measurable Spaces and Borel Sets

Definition 2.1

Recall that an algebra on a given set X is a collection of subsets $\mathcal{A} \subseteq \mathcal{P}(X)$ which contains $\varnothing$ and is closed under complements and finite unions (hence also finite intersections). A σ-algebra is an algebra which is closed under countable unions (hence also countable intersections). □

Remark 2.2

Arbitrary intersections of σ-algebras on X are still σ-algebras, so for a given collection $\mathcal{E} \subseteq \mathcal{P}(X)$, it makes sense to define the σ-algebra generated by $\mathcal{E}$ as the intersection of all σ-algebras containing $\mathcal{E}$. In other words, this is the $\subseteq$-smallest σ-algebra containing $\mathcal{E}$, denoted $\langle \mathcal{E} \rangle_\sigma$, and $\langle \mathcal{E} \rangle$ denotes the algebra generated by $\mathcal{E}$.

For $\mathcal{E} \subseteq \mathcal{P}(X)$, denote $\neg\mathcal{E} := \{X \setminus A : A \in \mathcal{E}\}$, the dual collection of $\mathcal{E}$. □

Proposition 2.3

Let X be a set and $\varnothing \in \mathcal{E} \subseteq \mathcal{P}(X)$. Then $\langle \mathcal{E} \rangle_\sigma$ is the $\subseteq$-smallest collection $\mathcal{S}$ which contains $\mathcal{E} \sqcup \neg\mathcal{E}$ and is closed under countable unions and countable intersections.

Proof. Put $\widetilde{\mathcal{S}} := \{A \in \mathcal{S} : A, A^c \in \mathcal{S}\} \subseteq \mathcal{S}$. We show $\widetilde{\mathcal{S}}$ is a σ-algebra containing $\mathcal{E}$, hence $\widetilde{\mathcal{S}} \supseteq \langle \mathcal{E} \rangle_\sigma$, and $\widetilde{\mathcal{S}} \subseteq \mathcal{S} \subseteq \langle \mathcal{E} \rangle_\sigma$ (the latter since $\langle \mathcal{E} \rangle_\sigma$ is itself a collection with the two properties, and $\mathcal{S}$ is the smallest one), hence $\widetilde{\mathcal{S}} = \mathcal{S} = \langle \mathcal{E} \rangle_\sigma$.

Observe that $\mathcal{E} \subseteq \widetilde{\mathcal{S}}$ since $A \in \mathcal{E} \implies A^c \in \neg\mathcal{E}$, and $\mathcal{S} \supseteq \mathcal{E} \sqcup \neg\mathcal{E} \implies A, A^c \in \mathcal{S} \implies A \in \widetilde{\mathcal{S}}$. Moreover, $\widetilde{\mathcal{S}}$ is a σ-algebra: For $A \in \widetilde{\mathcal{S}}$, $A^c \in \widetilde{\mathcal{S}}$ by definition, and

Source manuscript page 4

for $(A_n)_{n\in\mathbb{N}} \subseteq \widetilde{\mathcal{S}}$, $\bigcup_{n\in\mathbb{N}} A_n \in \mathcal{S}$ since $\mathcal{S}$ is closed under countable unions, and $\left(\bigcup_{n\in\mathbb{N}} A_n\right)^c = \bigcap_{n\in\mathbb{N}} A_n^c \in \mathcal{S}$ since $\mathcal{S}$ is closed under countable intersections; thus $\bigcup_{n\in\mathbb{N}} A_n \in \widetilde{\mathcal{S}}$. Lastly $\varnothing \in \widetilde{\mathcal{S}}$. Hence $\mathcal{E} \subseteq \widetilde{\mathcal{S}} \implies \langle \mathcal{E} \rangle_\sigma \subseteq \widetilde{\mathcal{S}}$. □

Definition 2.4

A measurable space is a set X equipped with a σ-algebra $\mathcal{I}$, i.e. a tuple $(X, \mathcal{I})$. For measurable spaces $(X, \mathcal{I})$, $(Y, \mathcal{J})$, a function $f : X \to Y$ is called measurable or $(\mathcal{I}, \mathcal{J})$-measurable if f^{-1} takes sets in $\mathcal{J}$ to sets in $\mathcal{I}$, i.e. $f^{-1}(B) \in \mathcal{I}$, $\forall B \in \mathcal{J}$.

When X is a topological space, the σ-algebra generated by the collection of all open sets is called the Borel σ-algebra of X, denoted $\mathcal{B}(X)$, and the sets in $\mathcal{B}(X)$ are called Borel sets.

For a function $f : X \to Y$ from a measurable space $(X, \mathcal{I})$ to a topological space Y, we say that f is measurable or $\mathcal{I}$-measurable if f is $(\mathcal{I}, \mathcal{B}(Y))$-measurable. Lastly, if X, Y are topological spaces, then a function $f : X \to Y$ is called Borel if it is $(\mathcal{B}(X), \mathcal{B}(Y))$-measurable. □

Note 2.5

If we ask for $(\mathcal{I}, \mathcal{J})$-measurability instead of $(\mathcal{I}, \mathcal{B}(Y))$-measurability (for a σ-algebra $\mathcal{J}$ on Y larger than $\mathcal{B}(Y)$), then the class of functions will be much smaller and less useful. □

Proposition 2.6

Continuous functions are Borel. More generally, let

Source manuscript page 5

$(X, \mathcal{I})$, $(Y, \mathcal{J})$ be measurable spaces, $\mathcal{E} \subseteq \mathcal{J}$ generating $\mathcal{J}$, i.e. $\langle \mathcal{E} \rangle_\sigma = \mathcal{J}$. Then if $f : X \to Y$ satisfies $f^{-1}(B) \in \mathcal{I}$, $\forall B \in \mathcal{E}$, then f is $(\mathcal{I}, \mathcal{J})$-measurable.

Proof. Note that the second part implies the first part. Define $\mathcal{J}_0 := \{B \in \mathcal{J} : f^{-1}(B) \in \mathcal{I}\}$. Then $\mathcal{J}_0 \subseteq \mathcal{J}$ while $\mathcal{J}_0 \supseteq \mathcal{E}$. Furthermore, $\mathcal{J}_0$ is a σ-algebra since $\varnothing = f^{-1}(\varnothing) \in \mathcal{I}$, and for $B, B_0, B_1, \ldots \in \mathcal{J}_0$, $f^{-1}(B^c) = f^{-1}(B)^c \in \mathcal{I}$, $f^{-1}(\bigcup_n B_n) = \bigcup_n f^{-1}(B_n) \in \mathcal{I}$, i.e. f^{-1} commutes with these operations and $\mathcal{I}$ is closed under them. Hence $\mathcal{J}_0 = \mathcal{J} = \langle \mathcal{E} \rangle_\sigma$. □

2.2 Stratification of Borel Sets into a Hierarchy

Definition 2.7

Let X be a topological space. We define the hierarchy of Borel sets level-by-level, via transfinite recursion. Let ω_1 denote the first uncountable ordinal. For each $1 \leq \xi < \omega_1$, define the collections of subsets of X as follows:

$$\mathbf{\Sigma}^0_1(X) := \{U \subseteq X : U \text{ open}\}, \qquad \mathbf{\Pi}^0_\xi(X) := \neg\mathbf{\Sigma}^0_\xi(X),$$
$$\mathbf{\Sigma}^0_\xi(X) := \Big\{ \bigcup_{n\in\mathbb{N}} A_n : A_n \in \mathbf{\Pi}^0_{\xi_n}(X),\ \xi_n < \xi,\ \forall n \in \mathbb{N} \Big\} \quad (2 \leq \xi < \omega_1).$$


In addition, we define the so-called ambiguous classes:

$$\mathbf{\Delta}^0_\xi(X) := \mathbf{\Sigma}^0_\xi(X) \cap \mathbf{\Pi}^0_\xi(X). \quad \square$$

Observation. Notice that by definition, for $2 \leq \xi < \eta$, $\mathbf{\Sigma}^0_\xi(X) \subseteq \mathbf{\Sigma}^0_\eta(X)$, since the collection we can pick the A_n's from is enlarged from ξ to η, and $\mathbf{\Pi}^0_\xi(X) = \neg\mathbf{\Sigma}^0_\xi(X) \subseteq \neg\mathbf{\Sigma}^0_\eta(X) = \mathbf{\Pi}^0_\eta(X)$.

Source manuscript page 6

Hence we have the following diagram:



(Blue arrows: each $\mathbf{\Pi}^0_\xi$ is the dual of $\mathbf{\Sigma}^0_\xi$, and $\mathbf{\Sigma}^0_{\xi+1}$ consists of the countable unions of $\mathbf{\Pi}^0_\xi$ sets.) Here ω denotes the first infinite ordinal, i.e. $\mathbb{N}$ (ordered as usual). $\mathbf{\Sigma}^0_\omega(X)$ contains unions $\bigcup_{n\in\mathbb{N}} A_n$ where $A_n \in \mathbf{\Pi}^0_n(X)$, and there is no reason why $\bigcup_{n\in\mathbb{N}} A_n$ should appear in $\mathbf{\Sigma}^0_m(X)$, for some $m \in \mathbb{N}$. Thus to obtain the full Borel sets, we must continue transfinitely via the definition above (up to ω_1, as we will see).

In particular, $\mathbf{\Sigma}^0_1(X) \subseteq \mathbf{\Sigma}^0_2(X)$ is not true in general topological spaces, as open sets are not necessarily F_σ; symmetrically, $\mathbf{\Pi}^0_1(X) \subseteq \mathbf{\Pi}^0_2(X)$ is not true in general, as closed sets are not necessarily G_δ. □

Proposition 2.8

Let X be a metrizable space.

(a) $\mathbf{\Sigma}^0_\eta(X) \cup \mathbf{\Pi}^0_\eta(X) \subseteq \mathbf{\Delta}^0_\xi(X)$, $\forall\, 1 \leq \eta < \xi < \omega_1$.
(b) $\mathcal{B}(X) = \biguplus_{1\leq\xi<\omega_1} \mathbf{\Sigma}^0_\xi(X) = \biguplus_{1\leq\xi<\omega_1} \mathbf{\Pi}^0_\xi(X) = \biguplus_{1\leq\xi<\omega_1} \mathbf{\Delta}^0_\xi(X)$.

Proof. (a) Take $A \in \mathbf{\Sigma}^0_\eta(X)$. Then $A \in \mathbf{\Sigma}^0_\xi(X)$ by our observation above. In particular, $\mathbf{\Sigma}^0_1(X) \subseteq \mathbf{\Sigma}^0_2(X)$ since open sets are F_σ in metrizable spaces (lecture 1, Part 1, Lemma 2.8). $A \in \mathbf{\Pi}^0_\xi(X)$ by taking $A_0 = A \in \mathbf{\Sigma}^0_\eta(X)$, $A_n = X \in \mathbf{\Sigma}^0_1(X)$ for $n \geq 1$; then by definition, $A = \bigcap_{n\in\mathbb{N}} A_n \in \mathbf{\Pi}^0_\xi(X)$.

Source manuscript page 7

Hence $A \in \mathbf{\Sigma}^0_\xi(X) \cap \mathbf{\Pi}^0_\xi(X) = \mathbf{\Delta}^0_\xi(X)$. Lastly, the dual simply takes the complement of the sets in the collection, thus

$$\begin{aligned}\mathbf{\Pi}^0_\eta(X) = \neg\mathbf{\Sigma}^0_\eta(X) &\subseteq \neg(\mathbf{\Sigma}^0_\xi(X) \cap \mathbf{\Pi}^0_\xi(X)) \\ &= (\neg\mathbf{\Sigma}^0_\xi(X)) \cap (\neg\mathbf{\Pi}^0_\xi(X)) = \mathbf{\Pi}^0_\xi(X) \cap \mathbf{\Sigma}^0_\xi(X) \\ &= \mathbf{\Delta}^0_\xi(X).\end{aligned}$$

Hence $\mathbf{\Sigma}^0_\eta(X) \cup \mathbf{\Pi}^0_\eta(X) \subseteq \mathbf{\Delta}^0_\xi(X)$.

(b) Notice that (a) implies that we have an intertwined chain of classes, and consequently

$$\bigcup_{\xi<\omega_1} \mathbf{\Sigma}^0_\xi(X) = \bigcup_{\xi<\omega_1} \mathbf{\Pi}^0_\xi(X) = \bigcup_{\xi<\omega_1} \mathbf{\Delta}^0_\xi(X) :$$

By definition of $\mathbf{\Delta}^0_\xi(X)$, $\bigcup_{\xi<\omega_1} \mathbf{\Sigma}^0_\xi(X) \supseteq \bigcup_{\xi<\omega_1} \mathbf{\Delta}^0_\xi(X)$ and $\bigcup_{\xi<\omega_1} \mathbf{\Pi}^0_\xi(X) \supseteq \bigcup_{\xi<\omega_1} \mathbf{\Delta}^0_\xi(X)$. On the other hand, by (a),


$$\begin{aligned}\bigcup_{\xi<\omega_1} \mathbf{\Delta}^0_\xi(X) \supseteq \bigcup_{1<\xi<\omega_1} \mathbf{\Delta}^0_\xi(X) &\supseteq \bigcup_{\eta<\omega_1} (\mathbf{\Sigma}^0_\eta(X) \cup \mathbf{\Pi}^0_\eta(X)) \\ &= \bigcup_{\xi<\omega_1} \mathbf{\Sigma}^0_\xi(X) \;\cup\; \bigcup_{\xi<\omega_1} \mathbf{\Pi}^0_\xi(X).\end{aligned}$$

It is left to show that $\mathcal{B}(X) =$ any of the unions. Clearly $\mathbf{\Sigma}^0_\xi(X) \subseteq \mathcal{B}(X)$ since at every inductive step we only take complements and countable unions. And $\mathbf{\Sigma}^0_1(X)$ generates $\mathcal{B}(X)$. Hence it remains to show the unions are σ-algebras. Indeed, closure under complement follows from $\neg\mathbf{\Delta}^0_\xi(X) = \mathbf{\Delta}^0_\xi(X)$, and closure under countable union follows from each $\mathbf{\Sigma}^0_\xi(X)$ being closed under countable union (each set in $\mathbf{\Sigma}^0_\xi(X)$ is a countable union of $A_n \in \mathbf{\Pi}^0_{\xi_n}(X)$,

Source manuscript page 8

and their countable union is a countable union of countable unions, hence still a countable union), together with the fact that countably many sets $A_n \in \mathbf{\Sigma}^0_{\xi_n}(X)$ all lie in one $\mathbf{\Sigma}^0_\xi(X)$, namely for $\xi := \sup_n \xi_n < \omega_1$ (a countable supremum of countable ordinals is countable). □

Observation. Thus for metrizable spaces, we have the following picture:



From HW 1, every second countable space has $\leq |2^{\mathbb{N}}| = \mathfrak{c}$ many open sets, hence also closed sets. Taking countable unions of sets from a collection of cardinality $\leq \mathfrak{c}$ still results in $\leq |2^{\mathbb{N}}|$ many sets since $(2^{\mathbb{N}})^{\mathbb{N}} \cong 2^{\mathbb{N}\times\mathbb{N}} \cong 2^{\mathbb{N}}$. Thus by transfinite induction on ξ, $|\mathbf{\Sigma}^0_\xi(X)| \leq \mathfrak{c}$ and $|\mathbf{\Pi}^0_\xi(X)| \leq \mathfrak{c}$. Finally

$$|\mathcal{B}(X)| = \Big|\bigcup_{1\leq\xi<\omega_1} \mathbf{\Sigma}^0_\xi(X)\Big| \leq |\omega_1| \times |2^{\mathbb{N}}| \leq |2^{\mathbb{N}} \times 2^{\mathbb{N}}| = |2^{\mathbb{N}}|,$$

with the last equality given by the interleaving map $((a_n)_n, (b_n)_n) \mapsto (a_1, b_1, a_2, b_2, \dots)$. Hence there are at most continuum-many Borel sets in a second countable space, and continuum is preserved along the zigzag chain above. In contrast, $\mathcal{L}$:= the Lebesgue measurable subsets of $\mathbb{R}$ has cardinality at least $2^{\mathfrak{c}} = 2^{|2^{\mathbb{N}}|}$,

Source manuscript page 9

since the $\frac{1}{3}$-Cantor set has cardinality continuum with Lebesgue measure 0, thus all of its subsets are Lebesgue measurable with measure 0. □

Proposition 2.9

Let X be a topological space and $1 \leq \xi < \omega_1$, i.e. ξ a countable ordinal.

(a) $\mathbf{\Sigma}^0_\xi(X)$ is closed under taking countable unions; $\mathbf{\Pi}^0_\xi(X)$ is closed under taking countable intersections; $\mathbf{\Delta}^0_\xi(X)$ is closed under taking complements.

(b) If X is metrizable, then all three classes are closed under taking finite unions and intersections.

Proof. (a) $\mathbf{\Sigma}^0_\xi(X)$ is closed under taking countable unions since for $\xi = 1$, a countable union of open sets is still open; for $\xi \geq 2$, a countable union of countable unions is still a countable union. $\mathbf{\Pi}^0_\xi(X)$ follows by taking complements with $\mathbf{\Sigma}^0_\xi(X)$, and

$$\neg\mathbf{\Delta}^0_\xi(X) = \neg(\mathbf{\Sigma}^0_\xi(X) \cap \mathbf{\Pi}^0_\xi(X)) = \neg\mathbf{\Sigma}^0_\xi(X) \cap \neg\mathbf{\Pi}^0_\xi(X) = \mathbf{\Pi}^0_\xi(X) \cap \mathbf{\Sigma}^0_\xi(X) = \mathbf{\Delta}^0_\xi(X).$$

(b) It suffices to show $\mathbf{\Sigma}^0_\xi(X)$ is closed under finite intersections and then everything else follows. Take $\bigcup_{n\in\mathbb{N}} A_n, \bigcup_{n\in\mathbb{N}} B_n \in \mathbf{\Sigma}^0_\xi(X)$. Then $\big(\bigcup_{n\in\mathbb{N}} A_n\big) \cap \big(\bigcup_{n\in\mathbb{N}} B_n\big) = \bigcup_{n,m\in\mathbb{N}}(A_n \cap B_m)$, where $A_n \in \mathbf{\Pi}^0_{\xi_n}(X)$,

$B_m \in \mathbf{\Pi}^0_{\eta_m}(X)$, $\xi_n, \eta_m < \xi$. Hence $A_n, B_m \in \mathbf{\Pi}^0_\eta(X)$, $\eta = \max(\xi_n, \eta_m) < \xi$, since $\mathbf{\Pi}^0_\zeta(X) \subseteq \mathbf{\Pi}^0_\eta(X)$ for $\zeta < \eta$ (Proposition 2.8(a)), and $A_n \cap B_m \in \mathbf{\Pi}^0_\eta(X)$ by (a). Thus $\bigcup_{n,m\in\mathbb{N}}(A_n \cap B_m) \in \mathbf{\Sigma}^0_\xi(X)$ by definition. □

Source manuscript page 10

Example 2.10

Recall we proved for X a compact metric space, Y a separable complete metric space, that $C(X,Y)$ is Polish with respect to the uniform metric (HW 1). We show the set of continuously differentiable functions on $[0,1]$, $C^1([0,1])$, is $\mathbf{\Pi}^0_3$ as a subset of $C([0,1]) = C([0,1],\mathbb{R})$, i.e. $C^1([0,1]) \in \mathbf{\Pi}^0_3(C([0,1]))$, i.e. having a definition of the form $\forall^{\mathbb{N}}\exists^{\mathbb{N}}\forall^{\mathbb{N}}$.

We would reduce the complexity of the definition by reducing the complexity of quantifiers (e.g. $\forall \varepsilon \in \mathbb{R}^+ \rightsquigarrow \forall \varepsilon \in \mathbb{Q}^+$) and applying theorems.

For $f \in C([0,1])$, $f \in C^1([0,1]) \iff \forall \varepsilon \in \mathbb{Q}^+$, $\exists$ finitely many rational open intervals $I_0, \dots, I_{n-1}$ covering $[0,1]$ such that $\forall\, 0 \le j < n$, $f \in A_{I_j,\varepsilon}$, where

$$A_{I_j,\varepsilon} := \left\{ f \in C([0,1]) : \forall\, a \neq b,\ c \neq d \text{ in } I_j,\ \left| \frac{f(a)-f(b)}{a-b} - \frac{f(c)-f(d)}{c-d} \right| \le \varepsilon \right\}. \qquad (*)$$

[Exercise. Hint: Recall that $g(x)$ is continuous at x_0 iff for all $x_n \xrightarrow{n\to\infty} x_0$, $g(x_n) \to g(x_0)$, and noticing that $(*)$ implies Cauchyness and convergence of g at once, we get differentiability at $x \in [0,1]$. Moreover, $(*)$ implies continuity of f' as well.]

Now fixing I_j with $a, b, c, d \in I_j$, the map

$$F_{a,b,c,d} : C([0,1]) \to [0,\infty), \qquad f \mapsto \left| \frac{f(a)-f(b)}{a-b} - \frac{f(c)-f(d)}{c-d} \right|$$

Source manuscript page 11

is easily continuous since every $f \in C([0,1])$ is uniformly continuous. Then

$$A_{I_j,\varepsilon} = \underbrace{\bigcap_{\substack{a,b,c,d\in I_j \\ a\neq b,\ c\neq d}} \underbrace{F^{-1}_{a,b,c,d}([0,\varepsilon])}_{\text{continuous preimage of closed sets = closed}}}_{\bigcap \text{ closed=closed}}$$

is closed in $C([0,1])$. Thus

$$f \in C^1([0,1]) \iff \underbrace{\underbrace{\forall \varepsilon \in \mathbb{Q}^+,\ \exists\, (I_0,\dots,I_{n-1}) \in \text{rational open intervals}^{<\mathbb{N}},\ \underbrace{\forall j < n,\ f \in A_{I_j,\varepsilon}}_{\mathbf{\Pi}^0_1=\text{closed}}}_{\mathbf{\Sigma}^0_2=F_\sigma}}_{\mathbf{\Pi}^0_3=\neg\mathbf{\Sigma}^0_3}$$

(Alternatively, $\mathbf{\Pi}^0_3$ is closed under countable intersections and $\mathbf{\Sigma}^0_2 \subseteq \mathbf{\Delta}^0_3 \subseteq \mathbf{\Pi}^0_3$.) Explicitly, if $\mathcal{I} := \{(p,q) \cap [0,1] : p < q \in \mathbb{Q}\}$, which is countable (! $\mathbb{N}^{<\mathbb{N}}$ is countable, $\mathbb{N}^{\mathbb{N}}$ is uncountable, $= |2^{\mathbb{N}}| = \mathfrak{c}$), then

$$C^1([0,1]) = \bigcap_{\varepsilon\in\mathbb{Q}^+} \bigcup_{\substack{(I_0,I_1,\dots,I_{n-1})\in\mathcal{I}^{<\mathbb{N}} \\ \bigcup_j I_j = [0,1]}} \bigcap_{0\le j<n} A_{I_j,\varepsilon}. \quad \square$$

2.3 The Classes $\mathcal{B}$, $\mathbf{\Sigma}^0_\xi$, $\mathbf{\Pi}^0_\xi$, $\mathbf{\Delta}^0_\xi$

Definition 2.11

By $\mathcal{B}$, $\mathbf{\Sigma}^0_\xi$, $\mathbf{\Pi}^0_\xi$, $\mathbf{\Delta}^0_\xi$, we denote the class of the corresponding sets in all topological spaces, and for each such space X, we write $\mathcal{B}(X)$, $\mathbf{\Sigma}^0_\xi(X)$, $\mathbf{\Pi}^0_\xi(X)$, $\mathbf{\Delta}^0_\xi(X)$ to mean the corresponding set of subsets of X. □

Source manuscript page 12


Notation. For a set $A \subseteq X \times Y$, we denote by

$$A_x := \{y \in Y : (x,y) \in A\}, \quad \text{for } x \in X, \qquad \text{and} \qquad A^y := \{x \in X : (x,y) \in A\}, \quad \text{for } y \in Y,$$

the vertical and horizontal fibers over the point $x \in X$ and $y \in Y$ respectively. Similarly, for a function $f : X \times Y \to Z$, denote by

$$f_x : Y \to Z,\ y \mapsto f(x,y) \qquad \text{and} \qquad f^y : X \to Z,\ x \mapsto f(x,y)$$

the vertical and horizontal fibers over the point $x \in X$ and $y \in Y$, respectively. □



Proposition 2.12

Let Γ be any one of the classes $\mathbf{\Sigma}^0_\xi$, $\mathbf{\Pi}^0_\xi$, $\mathbf{\Delta}^0_\xi$, $\xi < \omega_1$. Then Γ is closed under

(a) continuous preimages, i.e. if X, Y are topological spaces, $f : X \to Y$ is continuous and $B \in \Gamma(Y)$, then $f^{-1}(B) \in \Gamma(X)$;

(b) putting countably many sets as fibers over $\mathbb{N}$, i.e. if X is a topological space, $A_n \in \Gamma(X)$, then the set $B \subseteq \mathbb{N} \times X$ defined by $B_n := A_n$ is in $\Gamma(\mathbb{N} \times X)$, where B_n is the fiber of B over $n \in \mathbb{N}$, $\mathbb{N} \times X$ is equipped with the product topology, and $\mathbb{N}$ is equipped with the discrete topology.

Proof. (a) Note that the statement with $\mathbf{\Delta}^0_\xi$ follows from $\mathbf{\Sigma}^0_\xi$ and $\mathbf{\Pi}^0_\xi$ from the fact that taking intersections commutes with taking preimages. We show the statement for $\mathbf{\Sigma}^0_\xi$ and $\mathbf{\Pi}^0_\xi$ simultaneously by transfinite induction on ξ.

Base case: $\xi = 1$. Continuous preimages of open sets ($\mathbf{\Sigma}^0_1(Y)$) are

Source manuscript page 13

open ($\mathbf{\Sigma}^0_1(X)$). $\mathbf{\Pi}^0_1$ follows by taking the dual collection.

Inductive step: Our definition of $\mathbf{\Sigma}^0_\xi$ and $\mathbf{\Pi}^0_\xi$ does not split into successor / limit cases, thus we need not split into cases in our transfinite induction proof. Suppose $\mathbf{\Sigma}^0_\eta$ and $\mathbf{\Pi}^0_\eta$ are closed under continuous preimages, for all $\eta < \xi$. Then for $\bigcup_n A_n \in \mathbf{\Sigma}^0_\xi(Y)$, with $A_n \in \mathbf{\Pi}^0_{\xi_n}(Y)$, $\xi_n < \xi$, $\forall n$, we have $f^{-1}(\bigcup_n A_n) = \bigcup_n f^{-1}(A_n)$ with each $f^{-1}(A_n) \in \mathbf{\Pi}^0_{\xi_n}(X)$, $\xi_n < \xi$, by the induction hypothesis. Thus by definition, $f^{-1}(\bigcup_n A_n) = \bigcup_n f^{-1}(A_n) \in \mathbf{\Sigma}^0_\xi(X)$. (And $\mathbf{\Pi}^0_\xi$ follows by taking the dual collection.)

(b) Induct transfinitely on ξ in the same way as above. Alternatively, consider $\pi : \mathbb{N} \times X \to X$ the projection map, which is continuous by definition of the product topology. Then for $\bigcup_n A_n \in \mathbf{\Sigma}^0_\xi(X)$,

$$B = \bigcup_n (\{n\} \times B_n) = \bigcup_n \left[(\{n\} \times X) \cap \pi^{-1}(A_n) \right]$$

is a countable union of (clopen ($\mathbf{\Delta}^0_1(X) \subseteq \mathbf{\Sigma}^0_\xi(X)$) sets) $\cap$ ($\mathbf{\Sigma}^0_\xi(\mathbb{N} \times X)$ sets). We show that (clopen set) $\cap$ ($\mathbf{\Sigma}^0_\xi$ set) is $\mathbf{\Sigma}^0_\xi$. Indeed, clopen sets are $\mathbf{\Delta}^0_1$, hence $\mathbf{\Pi}^0_1$ and $\mathbf{\Sigma}^0_1$, hence $\mathbf{\Pi}^0_2$ since $\mathbf{\Pi}^0_1 \subseteq \mathbf{\Sigma}^0_2 \implies \mathbf{\Sigma}^0_1 = \neg\mathbf{\Pi}^0_1 \subseteq \neg\mathbf{\Sigma}^0_2 = \mathbf{\Pi}^0_2$, hence $\mathbf{\Pi}^0_\eta \; \forall \eta < \omega_1$ since $\mathbf{\Pi}^0_2 \subseteq \mathbf{\Pi}^0_3 \subseteq \cdots$ in any topological space. Let C be clopen and $\bigcup_n D_n \in \mathbf{\Sigma}^0_\xi$ with $D_n \in \mathbf{\Pi}^0_{\xi_n}$, $\xi_n < \xi \; \forall n$. Then $C \cap (\bigcup_n D_n) = \bigcup_n (C \cap D_n)$, where each $C \cap D_n$ is an intersection in $\mathbf{\Pi}^0_{\xi_n}$. Hence $C \cap D_n \in \mathbf{\Pi}^0_{\xi_n}$ since the $\mathbf{\Pi}^0_\eta$ classes are closed under countable intersections. Thus $\bigcup_n (C \cap D_n) \in \mathbf{\Sigma}^0_\xi$. Hence $B = \bigcup_n \left[(\{n\} \times X) \cap \pi^{-1}(A_n) \right] \in \mathbf{\Sigma}^0_\xi(\mathbb{N} \times X)$.

Source manuscript page 14

For $\mathbf{\Pi}^0_\xi$, notice that $(m, x) \in B = \bigcup_n \{n\} \times B_n \iff \forall n \in \mathbb{N}$, $n \neq m$ or $x \in A_n$, thus

$$B = \underbrace{\bigcap_{n \in \mathbb{N}} \underbrace{\left[\underbrace{((\mathbb{N} \setminus \{n\}) \times X)}_{\mathbf{\Delta}^0_1 \subseteq \mathbf{\Pi}^0_\xi} \cup \underbrace{\pi^{-1}(A_n)}_{\mathbf{\Pi}^0_\xi} \right]}_{\mathbf{\Pi}^0_\xi \; (*)}}_{\mathbf{\Pi}^0_\xi \text{ (by the closure property)}}.$$

To show $(*)$, first show that every set in $\mathbf{\Pi}^0_\xi$ takes the form $\bigcap_n E_n$ with $E_n \in \mathbf{\Sigma}^0_{\xi_n}$, $\xi_n < \xi$, then rewrite the union term as some countable $\bigcap$ in $\mathbf{\Pi}^0_\xi$ (distribute the clopen set into the intersection; a clopen set union a $\mathbf{\Sigma}^0_{\xi_n}$ set is $\mathbf{\Sigma}^0_{\xi_n}$ by the dual of what we just showed). □

Definition 2.13

Let Γ be one of the classes. Denote the dual class by $\neg\Gamma$, i.e. if X is a topological space, then

$$\neg\Gamma(X) := \{A \subseteq X : A^c \in \Gamma(X)\}, \qquad \text{thus } \mathbf{\Sigma}^0_\xi = \neg\mathbf{\Pi}^0_\xi.$$

Furthermore, for topological spaces X, Y and $A \subseteq X \times Y$, denote $\mathrm{proj}_X(A) := \{x \in X : \exists\, y \in Y \text{ such that } (x, y) \in A\}$, and put

$$\exists^Y \Gamma(X) := \{\mathrm{proj}_X(A) : A \in \Gamma(X \times Y)\} \subseteq \mathcal{P}(X),$$
$$\forall^Y \Gamma(X) := \neg\exists^Y \neg\Gamma(X) := \neg(\exists^Y(\neg\Gamma)(X)) \subseteq \mathcal{P}(X).$$

I.e.,

$$\begin{aligned}
S \in \exists^Y \Gamma(X) &\iff \exists\, A \in \Gamma(X \times Y)\ \forall x \in X,\ [x \in S \iff \exists\, y \in Y\ (x,y) \in A],\\
S \in \forall^Y \Gamma(X) &\iff S^c \in (\exists^Y(\neg\Gamma))(X)\\
&\iff \exists\, A \in \neg\Gamma(X \times Y)\ \forall x \in X\ [x \in S^c \iff \exists\, y \in Y\ (x,y) \in A]\\
&\iff \exists\, C \in \Gamma(X \times Y)\ \forall x \in X\ [x \in S \iff \forall y \in Y\ (x,y) \notin A] \quad (C = A^c)\\
&\iff \exists\, C \in \Gamma(X \times Y)\ \forall x \in X\ [x \in S \iff \forall y \in Y\ (x,y) \in C].
\end{aligned}$$

Source manuscript page 15

By $\neg\exists^Y\neg\Gamma(X)$, we mean: apply the operator $\exists^Y$ to the class $\neg\Gamma$ first, then apply $\exists^Y\neg\Gamma$ to X, and finally take the dual collection $\neg(\exists^Y\neg\Gamma(X))$. □

Corollary 2.14

$\forall\, 1 \le \xi < \omega_1$, for X metrizable, $\mathbf{\Sigma}^0_{\xi+1}(X) = \exists^{\mathbb{N}}\mathbf{\Pi}^0_\xi(X)$. Thus also $\mathbf{\Pi}^0_{\xi+1}(X) = \forall^{\mathbb{N}}\mathbf{\Sigma}^0_\xi(X)$.

Proof. $\exists^{\mathbb{N}}\mathbf{\Pi}^0_\xi(X) \subseteq \mathbf{\Sigma}^0_{\xi+1}(X)$: Take $S \in \exists^{\mathbb{N}}\mathbf{\Pi}^0_\xi(X)$, with $A \in \mathbf{\Pi}^0_\xi(\mathbb{N} \times X)$ such that $\mathrm{proj}_X(A) = S$. Then observe that $\mathrm{proj}_X(A) = \bigcup_{n\in\mathbb{N}} A_n$, with each fiber A_n = the continuous preimage of A by $\iota_n : x \mapsto (n,x)$. By Proposition 2.12(a), each A_n is $\mathbf{\Pi}^0_\xi(X)$. Hence $S \in \mathbf{\Sigma}^0_{\xi+1}(X)$.

$\mathbf{\Sigma}^0_{\xi+1}(X) \subseteq \exists^{\mathbb{N}}\mathbf{\Pi}^0_\xi(X)$: Take $S = \bigcup_{n\in\mathbb{N}} A_n$, with $A_n \in \mathbf{\Pi}^0_{\xi_n}(X)$, $\xi_n < \xi + 1$, i.e. $\xi_n \le \xi$. Thus $A_n \in \mathbf{\Pi}^0_\xi(X)$, $\forall n$ (using metrizability so that $\mathbf{\Pi}^0_1 \subseteq \mathbf{\Pi}^0_2$). By Proposition 2.12(b), A with the A_n as fibers over n is $\mathbf{\Pi}^0_\xi(\mathbb{N} \times X)$. Thus $S = \mathrm{proj}_X(A)$ is $\exists^{\mathbb{N}}\mathbf{\Pi}^0_\xi(X)$ by definition.

Lastly $\mathbf{\Pi}^0_{\xi+1}(X) = \neg\mathbf{\Sigma}^0_{\xi+1}(X) = \neg\exists^{\mathbb{N}}\mathbf{\Pi}^0_\xi(X) = \neg\exists^{\mathbb{N}}\neg\mathbf{\Sigma}^0_\xi(X) = \forall^{\mathbb{N}}\mathbf{\Sigma}^0_\xi(X)$. □

Remark 2.15

Thus $\bigcup_{1\le n<\omega} \mathbf{\Sigma}^0_n$ are all the sets obtained from open sets by $\neg$ and $\exists^{\mathbb{N}}$ (finite $\cap$ and $\cup$ can be encoded by $\neg$ and $\exists^{\mathbb{N}}$; in contrast, infinite $\cap$ and $\cup$ could escape to $\mathbf{\Sigma}^0_\omega$ outside of $\bigcup_{1\le n<\omega} \mathbf{\Sigma}^0_n$).

Source manuscript page 16

Because of this, the superscript "0" refers to the fact that we quantify over $\mathbb{N}$ as opposed to $2^{\mathbb{N}}$, i.e. $\mathcal{P}(\mathbb{N})$. $\mathbf{\Sigma}^1_n$, $\mathbf{\Pi}^1_n$ would then be used when we allow quantification over $2^{\mathbb{N}}$ or $\mathbb{N}^{\mathbb{N}}$ or $\mathbb{R}$ or any uncountable Polish space. □

2.4 Universal Sets and Diagonalization


Aside. A very recent work done by Evan Leach (MIT, UCLA, Sep. 2026) presented a combinatorial proof using infinite circuits of what we are going to prove next using diagonalization (the second corollary below, Corollary 2.23).

Lemma 2.16 (Diagonalization Lemma (Cantor))

For a set X and $R \subseteq X \times X$, the antidiagonal set

$$\text{AntiDiag}(R) := \{x \in X : (x, x) \notin R\}$$

is not a vertical or horizontal fiber of R, i.e. $\text{AntiDiag}(R) \neq R_x$ and $\text{AntiDiag}(R) \neq R^x$, $\forall x \in X$.



Proof. Notice that for every $x \in X$, $x \in \text{AntiDiag}(R) \iff (x, x) \notin R \iff x \notin R_x$ (and likewise $\iff x \notin R^x$), thus $\text{AntiDiag}(R) \neq R_x$ (and $\neq R^x$), for any $x \in X$. □

Corollary 2.17

$\mathbb{R}$ and $2^{\mathbb{N}}$ are uncountable.

We illustrate that the lemma above is really generalizing

Source manuscript page 17

the elementary digit-flipping proof of $\mathbb{R}$'s uncountability.

Proof. Suppose $(x_n)_{n\in\mathbb{N}} \subseteq 2^{\mathbb{N}}$, i.e. each x_i is a 0-1 sequence. We can view it as $R \subseteq \mathbb{N} \times \mathbb{N}$ where $(m, n) \in R \iff x_m(n) = 1$. Hence the fiber $R_m = \{n : x_m(n) = 1\}$ is just the set of entries of the m-th sequence having a "1". Consider $\text{AntiDiag}(R) = \{n : x_n(n) = 0\}$. Flipping the digits, we get $y(n) := 1 - x_n(n)$ (the characteristic function of $\text{AntiDiag}(R)$). By Cantor's lemma, $\text{AntiDiag}(R) \neq R_m$, $\forall m$, thus $y \neq x_m$ for any m. So $y \in 2^{\mathbb{N}} \setminus (x_n)_{n\in\mathbb{N}}$. One can inject $2^{\mathbb{N}} \hookrightarrow \mathbb{R}$ via the decimal representation of reals, hence $|\mathbb{N}| < |2^{\mathbb{N}}| \leq |\mathbb{R}|$. □

Exercise 2.18

Use the lemma and show $|X| < |\mathcal{P}(X)|$, i.e. no surjection $X \twoheadrightarrow \mathcal{P}(X)$ exists. (Hint: Suppose such $X \twoheadrightarrow \mathcal{P}(X)$ exists, then view it as a correspondence / subset of $X \times X$. What are the vertical fibres and what is the antidiagonal in this case? Correspondence = set-valued function.) □

Definition 2.19

Let Γ be a class of sets in topological spaces. Let X, Y be topological spaces. We say that a set $U \subseteq Y \times X$ parametrizes $\Gamma(X)$ if $\{U_y : y \in Y\} = \Gamma(X)$. Think of Y as a parameter set, and decoding along Y gives $\Gamma(X)$. If moreover U itself is in Γ, i.e. $U \in \Gamma(Y \times X)$, then we call U a Y-universal set for $\Gamma(X)$. □

Source manuscript page 18

Observation. Let Γ be any class of sets in topological spaces (i.e. Γ is an arbitrary class we can define over a topological space) which is closed under taking continuous preimages. Then Γ is closed under taking fibers, since if $R \in \Gamma(X \times Y)$ and $x \in X$, then R_x is the preimage of R under the continuous function $Y \to X \times Y$, $y \mapsto (x, y)$.

Note 2.20

We have used this observation once already, when we proved Corollary 2.14. □

Theorem 2.21

Let X be second countable metrizable. Then for $1 \leq \xi < \omega_1$, there is a $2^{\mathbb{N}}$-universal set for $\mathbf{\Sigma}^0_\xi(X)$ and for $\mathbf{\Pi}^0_\xi(X)$.

Proof. Transfinite induction on ξ.

$\xi = 1$. Fix a countable basis $(W_n)_{n\in\mathbb{N}}$ for X. Then define $U \subseteq 2^{\mathbb{N}} \times X$: for each $z \in 2^{\mathbb{N}}$,

$$U_z := \bigcup \{W_n : n \in \mathbb{N} \text{ and } z(n) = 1\}.$$

It is clear that $U = \bigcup_{z\in 2^{\mathbb{N}}} \{z\} \times U_z$ parametrizes $\mathbf{\Sigma}^0_1(X)$. Furthermore, U is $\mathbf{\Sigma}^0_1(2^{\mathbb{N}} \times X)$, i.e. open, because

$$\begin{aligned}(z, x) \in U &\iff x \in \bigcup \{W_n : n \in \mathbb{N},\ z(n) = 1\} \\ &\iff \underbrace{\underbrace{\underbrace{\exists n \in \mathbb{N}}_{\text{(countable) union}} \Big(\underbrace{\underbrace{z(n) = 1}_{\substack{\text{clopen} \\ (\{z : z(n)=1\} \times X) \\ \text{clopen (Part 1, trees)}}} \wedge \underbrace{x \in W_n}_{\substack{\text{open} \\ (2^{\mathbb{N}} \times W_n)}}}_{\text{open}} \Big)}_{\text{open}}\end{aligned}$$

(the cylinder sets $\{z : z(n) = 1\}$ are clopen by the note on cylinder sets in Part 1).

Inductive step. Suppose there is a $2^{\mathbb{N}}$-universal set $U^{(\eta)}$ for $\mathbf{\Sigma}^0_\eta(X)$, for each

Source manuscript page 19

$\eta < \xi$. Then $V^{(\eta)} := \left(U^{(\eta)}\right)^c$ is a $2^{\mathbb{N}}$-universal set for $\mathbf{\Pi}^0_\eta(X)$ because fibers commute with complements from the definition (exercise). It remains to build a universal set for $\mathbf{\Sigma}^0_\xi(X)$.

Fix $\eta_n < \xi$ with $\eta_n \leq \eta_{n+1}$, $\forall n \in \mathbb{N}$, such that $\sup\{\eta_n + 1 : n \in \mathbb{N}\} = \xi$. Take $B = \bigcup_k C_k \in \mathbf{\Sigma}^0_\xi(X)$, with each $C_k \in \mathbf{\Pi}^0_{\zeta_k}$, $\zeta_k < \xi$, $\forall k$. We rewrite B as follows: for each $\zeta < \xi = \sup_n(\eta_n + 1)$, there must be

some N with $\zeta \in \eta_N + 1$, i.e. $\zeta < \eta_N + 1$, i.e. $\zeta \leq \eta_N$. Then since the sequence is increasing, $\forall n \geq N$, $\eta_n \geq \eta_N \geq \zeta$. Now give each C_k its own slot: for each k, denote the N for ζ_k by N_k. Inductively set $n_k := \max(N_k, n_{k-1} + 1)$ so that $n_0 < n_1 < n_2 < \cdots$ and $\eta_{n_k} \geq \eta_{N_k} \geq \zeta_k$ from our choice of N. Put $B_{n_k} := C_k$, and for the rest, the unused slots, $B_n := \varnothing$. Observe that we rewrote $B = \bigcup_n B_n$ with $B_n \in \mathbf{\Pi}^0_{\eta_n}$, and particularly $B_{n_k} = C_k \in \mathbf{\Pi}^0_{\zeta_k} \subseteq \mathbf{\Pi}^0_{\eta_{n_k}}$.

Now to encode B, we need to encode all B_n into one code. We use $2^{\mathbb{N}} \cong (2^{\mathbb{N}})^{\mathbb{N}}$ (see the proof of Part 1, Theorem 4.12), so it's enough to show the existence of a $(2^{\mathbb{N}})^{\mathbb{N}}$-universal set for $\mathbf{\Sigma}^0_\xi(X)$. To this end, define $U \subseteq (2^{\mathbb{N}})^{\mathbb{N}} \times X$ by setting, for each $z \in (2^{\mathbb{N}})^{\mathbb{N}}$,

$$U_z := \bigcup_{n \in \mathbb{N}} \left(V^{(\eta_n)}\right)_{z(n)}.$$

From the definition, $U = \bigcup_{z \in (2^{\mathbb{N}})^{\mathbb{N}}} \{z\} \times U_z$ parametrizes $\mathbf{\Sigma}^0_\xi(X)$. Furthermore, U is itself

Source manuscript page 20

in $\mathbf{\Sigma}^0_\xi((2^{\mathbb{N}})^{\mathbb{N}} \times X)$ since

$$(z,x) \in U \iff x \in \bigcup_{n \in \mathbb{N}} \left(V^{(\eta_n)}\right)_{z(n)} \iff \underbrace{\underbrace{\exists n \in \mathbb{N}}_{\text{countable } \cup}\,,\ \underbrace{(z(n),x) \in V^{(\eta_n)}}_{\mathbf{\Pi}^0_{\eta_n}}}_{\mathbf{\Sigma}^0_\xi}$$

(the set of (z,x) with $(z(n),x) \in V^{(\eta_n)}$ is the preimage of $V^{(\eta_n)}$ under the continuous map $(z,x) \mapsto (z(n),x)$, hence $\mathbf{\Pi}^0_{\eta_n}$ by Proposition 2.12(a)). Universality follows. □

Corollary 2.22

Let X be second countable metrizable, Y an uncountable Polish space. For each $1 \leq \xi < \omega_1$, there is a Y-universal set for $\mathbf{\Sigma}^0_\xi(X)$, as well as one for $\mathbf{\Pi}^0_\xi(X)$.

Proof. We build Y-universal sets for $\mathbf{\Pi}^0_\xi(X)$, then take complements to get Y-universal sets for $\mathbf{\Sigma}^0_\xi(X)$. By the perfect set property for Polish spaces (i.e. the Cantor–Bendixson theorem, Part 1, Theorems 5.6 and 5.5), since Y is uncountable, there is a homeomorphic copy $C \subseteq Y$ of $2^{\mathbb{N}}$. Thus by Theorem 2.21, there is a C-universal set $U \in \mathbf{\Pi}^0_\xi(C \times X)$ for $\mathbf{\Pi}^0_\xi(X)$. Clearly U parametrizes $\mathbf{\Pi}^0_\xi(X)$; it remains to observe

$$\mathbf{\Pi}^0_\xi(C \times X) = \mathbf{\Pi}^0_\xi(Y \times X)\big|_{C \times X} := \{A \cap (C \times X) : A \in \mathbf{\Pi}^0_\xi(Y \times X)\}$$

(HW 3, Q2). But because C is closed, for any $A \in \mathbf{\Pi}^0_\xi(Y \times X)$, $A \cap (C \times X) \in \mathbf{\Pi}^0_\xi(Y \times X)$, since $C \times X$ is closed ($\mathbf{\Pi}^0_1(Y \times X)$), thus $\mathbf{\Pi}^0_\xi(Y \times X)$, and $\mathbf{\Pi}^0_\xi$ is closed under countable $\cap$. Hence $U \in \mathbf{\Pi}^0_\xi(C \times X) = \mathbf{\Pi}^0_\xi(Y \times X)|_{C \times X} \subseteq \mathbf{\Pi}^0_\xi(Y \times X)$, so U is Y-universal for $\mathbf{\Pi}^0_\xi(X)$. □

Corollary 2.23

For every uncountable Polish space X and every $1 \leq \xi < \omega_1$, $\mathbf{\Sigma}^0_\xi(X) \neq \mathbf{\Pi}^0_\xi(X)$. Consequently, we get the so-called strictness of the Borel hierarchy in uncountable Polish spaces,

Source manuscript page 21

that

$$\mathbf{\Delta}^0_1(X) \subsetneq \begin{matrix}\mathbf{\Sigma}^0_1(X)\\ \mathbf{\Pi}^0_1(X)\end{matrix} \subsetneq \mathbf{\Delta}^0_2(X) \subsetneq \begin{matrix}\mathbf{\Sigma}^0_2(X)\\ \mathbf{\Pi}^0_2(X)\end{matrix} \subsetneq \cdots$$

Proof. Let U be an X-universal set for $\mathbf{\Sigma}^0_\xi(X)$, and $A := \mathrm{AntiDiag}(U)$. Then by definition $A = \delta^{-1}(U^c)$ with $\delta : X \to X \times X$ given by $\delta(x) = (x, x)$. Moreover, $A \in \mathbf{\Pi}^0_\xi(X)$ since $U^c \in \mathbf{\Pi}^0_\xi(X \times X)$, δ is continuous, and $\mathbf{\Pi}^0_\xi$ is closed under continuous preimages (Proposition 2.12(a)). But A is not in $\mathbf{\Sigma}^0_\xi(X)$ because A is not a fiber of U (Lemma 2.16), yet U parametrizes $\mathbf{\Sigma}^0_\xi(X)$. □

Source manuscript page 22

3 Change of Topology

3.1 Turning Borel Sets into Clopen Sets

Notation. To be precise, we denote a Polish space X by $(X, \mathcal{T})$, where $\mathcal{T}$ is the topology of X. We have the following most useful tool with Borel sets.

Theorem 3.1

Let $(X, \mathcal{T})$ be a Polish space. For each Borel set $A \subseteq X$, there is a finer Polish topology $\mathcal{T}_A$ with respect to which A is clopen and $\mathcal{B}(\mathcal{T}_A) = \mathcal{B}(\mathcal{T})$.

We prove two lemmas first.

Lemma 3.2

Let $(X, \mathcal{T})$ be a Polish space, $F \subseteq X$ a closed set. Then the topology $\mathcal{T}_F$ generated by $\mathcal{T} \cup \{F\}$ is Polish, with F clopen in $\mathcal{T}_F$ and $\mathcal{B}(\mathcal{T}_F) = \mathcal{B}(\mathcal{T})$.

Proof. We just need to show $\mathcal{T}_F$ is Polish. Observe that $\mathcal{T}_F$ is nothing but the disjoint union topology of F and $X \setminus F$ with their relative topologies respectively, both of which are Polish since F and $X \setminus F$ are both G_δ subsets of the Polish space $(X, \mathcal{T})$ (Part 1, Theorem 2.9), i.e. the map

F	$X \setminus F$

$$(X, \mathcal{T}_F) \to \{0\} \times F \sqcup \{1\} \times (X \setminus F), \qquad x \mapsto \begin{cases}(0, x) & \text{if } x \in F;\\ (1, x) & \text{otherwise}\end{cases}$$

is (trivially) a homeomorphism. But we showed on day 1 (Part 1, Proposition 2.4(b)) that a disjoint union of Polish spaces is Polish (with respect to $d((n, x), (m, y)) := d_n(x, y)$ if $n = m$; 1 otherwise). Hence $(X, \mathcal{T}_F)$

Source manuscript page 23

is Polish. In particular, $\mathcal{B}(X, \mathcal{T}_F) = \mathcal{B}(X, \mathcal{T})$ since F is closed, hence Borel in $\mathcal{T}$. □

Lemma 3.3

Let $(X, \mathcal{T})$ be a Polish space, $(\mathcal{T}_n)_{n\in\mathbb{N}}$ a sequence of Polish topologies on X refining $\mathcal{T}$ with $\mathcal{T}_n \subseteq \mathcal{B}(\mathcal{T})$ (i.e. $\mathcal{B}(\mathcal{T}_n) = \mathcal{B}(\mathcal{T})$). Then the topology $\mathcal{T}_\infty$ generated by $\bigcup_{n\in\mathbb{N}} \mathcal{T}_n$ is Polish with $\mathcal{B}(\mathcal{T}_\infty) = \mathcal{B}(\mathcal{T})$.

Proof. Take a countable basis $\mathcal{U}_n$ for each $\mathcal{T}_n$. Then $\mathcal{T}_\infty$ is generated by the finite intersections of sets from $\bigcup_n \mathcal{U}_n$. So every open set in $\mathcal{T}_\infty$ is some countable union (by second countability of $\mathcal{T}_\infty$) of finite intersections of sets from $\bigcup_n \mathcal{U}_n$, hence $\mathcal{B}(\mathcal{T})$. Thus $\mathcal{T}_\infty \subseteq \mathcal{B}(\mathcal{T}) \implies \mathcal{B}(\mathcal{T}_\infty) \subseteq \mathcal{B}(\mathcal{T})$ since $\mathcal{B}(\mathcal{T}_\infty)$ is the smallest σ-algebra containing $\mathcal{T}_\infty$ (and $\supseteq$ holds since $\mathcal{T} \subseteq \mathcal{T}_\infty$).

It is left to show that $(X, \mathcal{T}_\infty)$ admits a complete compatible metric. We show that $(X, \mathcal{T}_\infty)$ is homeomorphic to the diagonal $\Delta := \{(x)_{n\in\mathbb{N}} : x \in X\} \subseteq X^{\mathbb{N}}$ equipped with the product topology, via

$$\delta : (X, \mathcal{T}_\infty) \to \Big(X^{\mathbb{N}}, \prod_{n\in\mathbb{N}} \mathcal{T}_n\Big), \qquad x \mapsto (x)_n = (x, x, \dots).$$

Clearly δ is a bijection onto Δ. δ is continuous since each coordinate map $\delta_n := \mathrm{proj}_n \circ \delta : (X, \mathcal{T}_\infty) \to (X, \mathcal{T}_n)$ is continuous: δ_n is simply the identity map from a finer topology to a coarser topology, hence continuous. Then it is easy to check that the coordinate maps δ_n being

Source manuscript page 24

continuous $\implies$ δ continuous. On the other hand, to show $\delta^{-1} : \Delta \to (X, \mathcal{T}_\infty)$ is continuous, it suffices to show that preimages of open sets in $\mathcal{T}_n$ by δ^{-1} are open, since $\mathcal{T}_\infty$ is generated by $\bigcup_n \mathcal{T}_n$, and preimages commute with $\cap$ and $\cup$. But for any $n \in \mathbb{N}$, $\delta^{-1} = \mathrm{proj}_n|_\Delta$, hence for $U \in \mathcal{T}_n$, $(\delta^{-1})^{-1}(U) = (\mathrm{proj}_n|_\Delta)^{-1}(U)$ is open in the product topology since the proj_n are continuous by definition of the product topology.

Therefore $(X, \mathcal{T}_\infty) \xrightarrow{\delta} \Delta \subseteq X^{\mathbb{N}}$ is a homeomorphism. Furthermore, Δ is closed since each $\mathcal{T}_n$ is Hausdorff (exercise: show the complement is open). Hence Δ is Polish since it is a G_δ subset of a Polish space (a countable product of Polish spaces is Polish, Part 1, Proposition 2.4(c), and closed sets are G_δ, Part 1, Theorem 2.9). Thus $(X, \mathcal{T}_\infty)$ is Polish. □

Proof of Theorem 3.1 (Clopenification). We do the same σ-algebra trick. Let

$$\mathcal{S} := \{A \in \mathcal{B}(\mathcal{T}) : \exists \text{ Polish } \mathcal{T}_A \supseteq \mathcal{T} \text{ with } A \text{ clopen in } \mathcal{T}_A \text{ and } \mathcal{B}(\mathcal{T}_A) = \mathcal{B}(\mathcal{T})\}.$$

$\mathcal{S} \supseteq$ all closed sets by Lemma 3.2. $\mathcal{S}$ is closed under complements since clopenness is. Finally, for $A_n \in \mathcal{S}$, there are Polish $\mathcal{T}_n \supseteq \mathcal{T}$, $\mathcal{B}(\mathcal{T}_n) = \mathcal{B}(\mathcal{T})$, with A_n clopen in $\mathcal{T}_n$; by Lemma 3.3, $\mathcal{T}_\infty$ is Polish with $\bigcup_n A_n$ open in $\mathcal{T}_\infty$. Lastly apply Lemma 3.2 again (to the closed set $\big(\bigcup_n A_n\big)^c$) so that $\bigcup_n A_n$ is clopen in $\widetilde{\mathcal{T}}_\infty \supseteq \mathcal{T}_\infty$ with $\mathcal{B}(\widetilde{\mathcal{T}}_\infty) = \mathcal{B}(\mathcal{T}_\infty) = \mathcal{B}(\mathcal{T})$. Hence $\bigcup_n A_n \in \mathcal{S}$, i.e. $\mathcal{S}$ is

Source manuscript page 25

closed under countable $\cup$, hence a σ-algebra. But open sets ($= \mathcal{T}$) are in $\mathcal{S}$ (since closed sets are) and $\mathcal{S} \subseteq \mathcal{B}(\mathcal{T}) \implies \langle\mathcal{T}\rangle_\sigma = \mathcal{B}(\mathcal{T}) = \mathcal{S}$. □

Caution. We proved that a closed (thus the same for an open) set can be made clopen by simply taking the topology $\mathcal{T}_F$ generated by $\mathcal{T} \cup \{F\}$. However, it is not true that if B is a Borel set in a Polish space $(X, \mathcal{T})$, then the topology generated by $\mathcal{T} \cup \{B, B^c\}$ is Polish. To obtain a Polish topology in

which B is clopen, one has to clopenify all (countably many) sets that go into the definition of B from open / closed sets (what we did in Lemma 3.3). □

Corollary 3.4

Let $\mathcal{U}$ be a countable collection of Borel sets in a Polish space $(X, \mathcal{T})$. Then there is a Polish topology $\mathcal{T}' \supseteq \mathcal{T}$ in which every set in $\mathcal{U}$ is clopen and $\mathcal{B}(\mathcal{T}') = \mathcal{B}(\mathcal{T})$.

Proof. For each $U \in \mathcal{U}$, let $\mathcal{T}_U \supseteq \mathcal{T}$ be a Polish topology in which U is clopen (Theorem 3.1). Then by Lemma 3.3 above, the topology generated by $\bigcup_{U\in\mathcal{U}} \mathcal{T}_U$ is Polish with the desired property. □

Definition 3.5

Recall that a topological space X has the perfect set property (PSP) if X is either countable or there is a nonempty perfect subset (i.e. perfect closed) of X.

Source manuscript page 26

If X is Polish, then X has the PSP iff X is countable or $2^{\mathbb{N}}$ topologically embeds into X: ($\Rightarrow$) is by the perfect set theorem from Part 1, §5 (Part 1, Theorem 5.4). ($\Leftarrow$) since $2^{\mathbb{N}}$ is compact, thus compact in the Hausdorff space X, hence closed in X (Part 1, Theorem 4.7(b)). Thus $2^{\mathbb{N}}$ is a nonempty perfect (closed) subset of X. □

We now give the corollary that not only Polish spaces and Polish subspaces (i.e. G_δ subsets of Polish spaces) have the PSP, but the Borel sets also have it.

Corollary 3.6

Every Borel set B in a Polish space $(X, \mathcal{T})$ has the perfect set property.

Proof. Take a Polish topology $\mathcal{T}' \supseteq \mathcal{T}$ on X in which B is clopen, hence Polish in the subspace topology of $\mathcal{T}'$. By the Cantor–Bendixson theorem (Part 1, Theorem 5.5), B has the PSP, so if B is uncountable, then there is a continuous embedding $f : 2^{\mathbb{N}} \hookrightarrow (B, \mathcal{T}'|_B)$. Then trivially, f is still continuous as a map to $(B, \mathcal{T}|_B)$ and still injective, hence an embedding into $(B, \mathcal{T}|_B)$ because $2^{\mathbb{N}}$ is compact and $\mathcal{T}|_B$ is Hausdorff (Part 1, Theorem 4.7(e)). □

Corollary 3.7 (Making Borel functions continuous)

Let $(X, \mathcal{T})$ be a Polish space and Y be a second countable topological space. Let $f : X \to Y$ be a Borel (measurable) function. Then there is a Polish topology $\mathcal{T}_f \supseteq \mathcal{T}$ such that $f : (X, \mathcal{T}_f) \to Y$ is continuous and $\mathcal{B}(\mathcal{T}_f) = \mathcal{B}(\mathcal{T})$.

Source manuscript page 27

Proof. Let $\mathcal{U}$ be a countable basis for Y, and $\mathcal{T}_f \supseteq \mathcal{T}$ a Polish topology with $\mathcal{B}(\mathcal{T}_f) = \mathcal{B}(\mathcal{T})$ in which every set $f^{-1}(U)$ is clopen, for all $U \in \mathcal{U}$ (Corollary 3.4). Hence f is continuous from $(X, \mathcal{T}_f)$ to Y. □

Corollary 3.8

Let $(X, \mathcal{T})$ be a Polish space, $B \subseteq X$ a Borel subset. Then there is a continuous bijection $f : F \to B$ for some closed $F \subseteq \mathbb{N}^{\mathbb{N}}$. In particular, if $B \neq \varnothing$, then there is a continuous surjection $\mathbb{N}^{\mathbb{N}} \twoheadrightarrow B$.

Proof. Let $\mathcal{T}' \supseteq \mathcal{T}$ be a Polish topology with B clopen. Hence $(B, \mathcal{T}'|_B)$ is Polish, so there is a continuous bijection $f : F \xrightarrow{\sim} (B, \mathcal{T}'|_B)$, where $F \subseteq \mathbb{N}^{\mathbb{N}}$ is closed (Part 1, Theorem 6.14). If $B \neq \varnothing$, then there is a continuous surjection $g : \mathbb{N}^{\mathbb{N}} \twoheadrightarrow (B, \mathcal{T}'|_B)$, composing f with a retraction $\mathbb{N}^{\mathbb{N}} \twoheadrightarrow F$ (Part 1, Proposition 3.17). But then f, g are trivially continuous with respect to the coarser topology $\mathcal{T}|_B$. □

Proposition 3.9

Let $(X, \mathcal{T})$ be a Polish space. Then there exists a zero-dimensional Polish topology $\mathcal{T}_0 \supseteq \mathcal{T}$ such that $\mathcal{B}(\mathcal{T}_0) = \mathcal{B}(\mathcal{T})$.

Proof. HW 3. □

Proposition 3.10

Let Γ be a countable group, and consider a Borel action of Γ on a Polish space $(X, \mathcal{T})$, i.e. every $g \in \Gamma$ acts as a Borel automorphism of X. Then there exists a Polish topology $\mathcal{T}_0 \supseteq \mathcal{T}$ with $\mathcal{B}(\mathcal{T}_0) = \mathcal{B}(\mathcal{T})$ that makes the action of Γ continuous. Moreover, $\mathcal{T}_0$ can be taken to be zero-dimensional.

Source manuscript page 28

Proof. HW 3. □

3.2 Making Closed-Graph Functions Continuous for Free

Definition 3.11

Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces, $f : X \to Y$. The topology generated by f over $\mathcal{T}_X$ is the coarsest topology $\mathcal{T}_{X,f} \supseteq \mathcal{T}_X$ which makes f continuous. □

Observation. $\mathcal{T}_{X,f}$ is the topology generated by $\mathcal{T}_X \cup \{f^{-1}(V) : V \in \mathcal{T}_Y\}$. Every $U \in \mathcal{T}_{X,f}$ is an arbitrary union of sets of the form $U \cap f^{-1}(V)$, $U \in \mathcal{T}_X$, $V \in \mathcal{T}_Y$.

Lemma 3.12

Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be topological spaces and $f : X \to Y$. Then we have two topologies on $X \times Y$: $\mathcal{T}_X \times \mathcal{T}_Y$ and $\mathcal{T}_{X,f} \times \mathcal{T}_Y$. On the graph $G := \mathrm{graph}(f) = \{(x, f(x)) : x \in X\} \subseteq X \times Y$, the relative topologies with respect to $\mathcal{T}_X \times \mathcal{T}_Y$ and $\mathcal{T}_{X,f} \times \mathcal{T}_Y$ coincide.


Proof. Notice that the basic open sets in $\mathcal{T}_{X,f} \times \mathcal{T}_Y|_G$ have the form

$$[(U \cap f^{-1}(V)) \times W] \cap G = [(U \times W) \cap G] \cap [(f^{-1}(V) \times W) \cap G].$$

But $U \times W$ is open in $\mathcal{T}_X \times \mathcal{T}_Y$, thus $(U \times W) \cap G$ is open in $\mathcal{T}_X \times \mathcal{T}_Y|_G$. It is left to show that $(f^{-1}(V) \times W) \cap G$ is open in $\mathcal{T}_X \times \mathcal{T}_Y|_G$. But this is simply because

$$(f^{-1}(V) \times W) \cap G = [X \times (V \cap W)] \cap G,$$

which is open in $\mathcal{T}_X \times \mathcal{T}_Y|_G$ since $X \in \mathcal{T}_X$ and $V \cap W \in \mathcal{T}_Y$.



□

Lemma 3.13 (Continuity via graph)

Let X, Y be topological spaces. A

Source manuscript page 29

function $f : X \to Y$ is continuous

$$\iff \pi_f : X \to \text{graph}(f),\ x \mapsto (x, f(x)) \text{ is a homeomorphism} \iff \pi_f \text{ is continuous.}$$

Proof. If π_f is continuous, then so is f since $f = \text{proj}_Y \circ \pi_f$. On the other hand, if f is continuous, then so is π_f since each coordinate function $x \mapsto x$, $x \mapsto f(x)$ is continuous, hence continuity of π_f by definition of the product topology. Lastly, we get π_f^{-1} continuous for free since $\pi_f^{-1} = \text{proj}_X|_{\text{graph}(f)}$. □

Proposition 3.14 (Ruiyuan Chen)

Let $(X, \mathcal{T}_X)$, $(Y, \mathcal{T}_Y)$ be Polish spaces, $f : X \to Y$. If $\text{graph}(f)$ is closed in $\mathcal{T}_X \times \mathcal{T}_Y$ (for instance, if f is continuous with respect to some coarser Hausdorff topology on Y), then $\mathcal{T}_{X,f}$ is Polish.

Proof. $f : (X, \mathcal{T}_{X,f}) \to Y$ is continuous by definition of $\mathcal{T}_{X,f}$. Set $G := \text{graph}(f)$. Then $\pi_f : (X, \mathcal{T}_{X,f}) \to (G, \mathcal{T}_{X,f} \times \mathcal{T}_Y|_G)$ is a homeomorphism (Lemma 3.13). Then $(G, \mathcal{T}_X \times \mathcal{T}_Y|_G)$ is Polish, being a closed subset of the Polish space $(X \times Y, \mathcal{T}_X \times \mathcal{T}_Y)$, and $\mathcal{T}_X \times \mathcal{T}_Y|_G = \mathcal{T}_{X,f} \times \mathcal{T}_Y|_G$ by Lemma 3.12. Thus $(X, \mathcal{T}_{X,f}) \cong (G, \mathcal{T}_X \times \mathcal{T}_Y|_G) = (G, \mathcal{T}_{X,f} \times \mathcal{T}_Y|_G)$ is Polish. □

Source manuscript page 30

4 Analytic Sets

4.1 Definition and Closure Properties

We know that the class of Borel sets is closed under continuous / Borel preimages. Lebesgue had a false proof that it is closed under continuous images. It was discovered by Souslin years later, who proved that these are new sets and studied them with his PhD advisor Luzin.

Definition 4.1

A subset A of a Polish space X is called analytic if it is a continuous image of a Borel subset of some Polish space, i.e. if there are a Polish space Y, a Borel subset $B \subseteq Y$, and a continuous function $f : Y \to X$ such that $A = f(B)$. □

Proposition 4.2 (Equivalence of analyticity)

Let X be Polish, $\varnothing \neq A \subseteq X$. Then the following are equivalent:

(1) A is analytic (the definition above).
(2) There are a Polish space Y and a continuous $f : Y \to X$ such that $f(Y) = A$.
(3) There is a continuous $f : \mathbb{N}^{\mathbb{N}} \to X$ such that $f(\mathbb{N}^{\mathbb{N}}) = A$.
⋆(4) There is a closed $F \subseteq X \times \mathbb{N}^{\mathbb{N}}$ such that $\mathrm{proj}_X(F) = A$.
(5) There are a Polish Y and a Borel $B \subseteq X \times Y$ such that $\mathrm{proj}_X(B) = A$.
(6) There are a Polish Y, a Borel $B \subseteq Y$ and a Borel (measurable) $f : Y \to X$ such that $f(B) = A$.

Proof. Immediately, (4) ⇒ (5) ⇒ (1) and (1) ⇒ (6). (1) ⇒ (2) follows by a change of topology to make B clopen (Theorem 3.1), hence Polish with a finer topology $\mathcal{T}_B$, but then f is just more continuous. (2) ⇒ (3) since there is a continuous surjection $\mathbb{N}^{\mathbb{N}} \twoheadrightarrow Y$ (Part 1, Theorem 6.14), thus we can

Source manuscript page 31

compose it with f. (3) ⇒ (4) follows by taking

$$F = \mathrm{graph}(f)^{-1} := \{(f(x), x) : x \in \mathbb{N}^{\mathbb{N}}\}$$

since the graph of a continuous function (into a Hausdorff space) is closed, and $A = f(\mathbb{N}^{\mathbb{N}}) = \mathrm{proj}_X(F)$. (6) ⇒ (1) since we can make a Borel measurable function into a continuous function by Corollary 3.7. □

Notation. We denote by $\mathbf{\Sigma}^1_1$ the class of all analytic subsets of Polish spaces, so given a Polish space X, the set of analytic subsets of X is denoted by $\mathbf{\Sigma}^1_1(X)$. Its dual class is denoted by $\mathbf{\Pi}^1_1 := \neg\mathbf{\Sigma}^1_1$, with

$$\mathbf{\Pi}^1_1(X) = \neg\mathbf{\Sigma}^1_1(X) := \{A^c : A \in \mathbf{\Sigma}^1_1(X)\},$$

its elements called coanalytic sets. We also have the ambiguous class $\mathbf{\Delta}^1_1 := \mathbf{\Pi}^1_1 \cap \mathbf{\Sigma}^1_1$.

By (4) above, we have that $\mathbf{\Sigma}^1_1 = \exists^{\mathbb{N}^{\mathbb{N}}}\mathbf{\Pi}^0_1$, i.e. $\mathbf{\Sigma}^1_1(X) = \exists^{\mathbb{N}^{\mathbb{N}}}\mathbf{\Pi}^0_1(X)$, as we discussed in Remark 2.15. Taking its dual, we get $\mathbf{\Pi}^1_1(X) = \forall^{\mathbb{N}^{\mathbb{N}}}\mathbf{\Sigma}^0_1(X)$. This is what will allow us to build a universal set for $\mathbf{\Sigma}^1_1$, hence an analytic set which is not Borel, via Cantor's diagonalization. □

Observation. $\mathcal{B}(X) \subseteq \mathbf{\Delta}^1_1(X)$ since by (6), taking $f = \mathrm{id}$, all Borel sets are analytic; but $\mathcal{B}(X)$ is closed under taking complements, hence they are also coanalytic. □

Proposition 4.3

The class $\mathbf{\Sigma}^1_1$ is closed under

Source manuscript page 32

(i) continuous images and preimages (in fact, Borel images and preimages);
(ii) countable unions and countable intersections.

Proof. (i) Let X, Y be Polish spaces, $g : X \to Y$ continuous, $\varnothing \neq A \subseteq X$ an analytic subset. Then there is a continuous $f : \mathbb{N}^{\mathbb{N}} \to X$ with $f(\mathbb{N}^{\mathbb{N}}) = A$. Thus $g(A) \subseteq Y$ is witnessed by $g \circ f : \mathbb{N}^{\mathbb{N}} \to Y$.

Now suppose $B \subseteq Y$ is analytic. Then there is a closed $F \subseteq Y \times \mathbb{N}^{\mathbb{N}}$ such that $B = \mathrm{proj}_Y(F)$. Define $\tilde{g} : X \times \mathbb{N}^{\mathbb{N}} \to Y \times \mathbb{N}^{\mathbb{N}}$ as $\tilde{g}((x, n)) := (g(x), n)$, which is continuous since its component maps are. Hence $\tilde{g}^{-1}(F)$ is closed. Moreover, observe that

$$\begin{aligned} x \in g^{-1}(B) &\iff g(x) \in B = \mathrm{proj}_Y(F) \iff \exists\, n \in \mathbb{N}^{\mathbb{N}} \text{ such that } (g(x), n) \in F \\ &\iff \exists\, n \in \mathbb{N}^{\mathbb{N}} \text{ such that } (x, n) \in \tilde{g}^{-1}(F) \iff x \in \mathrm{proj}_X\left(\tilde{g}^{-1}(F)\right). \end{aligned}$$

Thus $g^{-1}(B) = \mathrm{proj}_X\left(\tilde{g}^{-1}(F)\right)$, with the latter an analytic set by condition (4) of Proposition 4.2 and the fact that $\tilde{g}^{-1}(F)$ is closed (since $\tilde{g}$ is continuous). Borel functions follow from the fact that they can be made continuous in a finer Polish topology on X (Corollary 3.7).

(ii) Countable intersections: Let X be a Polish space, $A_n \subseteq X$ analytic. Then there are closed $C_n \subseteq X \times \mathbb{N}^{\mathbb{N}}$ such that $A_n = \mathrm{proj}_X(C_n)$, i.e. for each $x \in X$, $x \in A_n \iff \exists\, y \in \mathbb{N}^{\mathbb{N}}$, $(x, y) \in C_n$. Let $A := \bigcap_{n \in \mathbb{N}} A_n$. Then for $x \in X$,

$$x \in A \iff \forall n \in \mathbb{N},\ \exists\, y_n \in \mathbb{N}^{\mathbb{N}},\ (x, y_n) \in C_n \iff \exists\, y \in (\mathbb{N}^{\mathbb{N}})^{\mathbb{N}}\ \forall n \in \mathbb{N},\ (x, y(n)) \in C_n.$$

Source manuscript page 33

Hence with $\widetilde{C}_n := \{(x, y) \in X \times (\mathbb{N}^{\mathbb{N}})^{\mathbb{N}} : (x, y(n)) \in C_n\}$, we get the following:

(a) $A = \mathrm{proj}_X\left(\bigcap_n \widetilde{C}_n\right)$ (trivial).
(b) The $\widetilde{C}_n$ are closed: Observe that the map $\pi_n : X \times (\mathbb{N}^{\mathbb{N}})^{\mathbb{N}} \to X \times \mathbb{N}^{\mathbb{N}}$, $(x, y) \mapsto (x, y(n))$, is continuous since each component map is a projection, and $\widetilde{C}_n = \pi_n^{-1}(C_n)$ with C_n closed, hence the $\widetilde{C}_n$ are closed.

Thus $\bigcap_n \widetilde{C}_n$ is closed, thus A is analytic by (a) and the fact that $\mathbb{N}^{\mathbb{N}} \cong \mathbb{N}^{\mathbb{N} \times \mathbb{N}} \cong (\mathbb{N}^{\mathbb{N}})^{\mathbb{N}}$. We leave the countable union as an exercise (HW 3). □

4.2 Universal Sets for Analytic / Coanalytic Sets

In the same way we proved the strictness of the Borel hierarchy in uncountable Polish spaces with universal sets and Cantor's diagonalization, we show the analog for analytic / coanalytic sets, that $\mathbf{\Pi}^1_1 \neq \mathbf{\Sigma}^1_1$.

Theorem 4.4 (Souslin)

For every uncountable Polish Y and an arbitrary Polish X, there is a Y-universal set for $\mathbf{\Sigma}^1_1(X)$. The same holds for $\mathbf{\Pi}^1_1(X)$.

Proof. We use $\mathbf{\Sigma}^1_1 = \exists^{\mathbb{N}^{\mathbb{N}}}\mathbf{\Pi}^0_1$. Let $F \subseteq Y \times (X \times \mathbb{N}^{\mathbb{N}})$ be a Y-universal set for $\mathbf{\Pi}^0_1(X \times \mathbb{N}^{\mathbb{N}})$ (Corollary 2.22). Let $U := \mathrm{proj}_{Y\times X}(F)$. Then we can verify that U is Y-universal for $\mathbf{\Sigma}^1_1(X)$

Source manuscript page 34

because fibers commute with projections:

$$\{U_y : y \in Y\} = \{\mathrm{proj}_X(F_y) : y \in Y\} = \{\mathrm{proj}_X(C) : C \in \mathbf{\Pi}^0_1(X \times \mathbb{N}^{\mathbb{N}})\} = \mathbf{\Sigma}^1_1(X)$$

by definition, and $U = \mathrm{proj}_{Y\times X}(F) \in \mathbf{\Sigma}^1_1(Y \times X)$ by definition as well. □

Corollary 4.5 (Souslin)


For each uncountable Polish space X, $\mathbf{\Sigma}^1_1(X) \neq \mathbf{\Pi}^1_1(X)$. In particular, $\mathcal{B}(X) \subseteq \mathbf{\Delta}^1_1(X) \subsetneq \mathbf{\Sigma}^1_1(X), \mathbf{\Pi}^1_1(X)$.

Proof. Let $U \subseteq X \times X$ be an X-universal set for $\mathbf{\Sigma}^1_1(X)$. Then $A := \mathrm{AntiDiag}(U) = \{x \in X : (x, x) \notin U\}$ is coanalytic because $A = \delta^{-1}(U^c) = \delta^{-1}(U)^c$, where $\delta : X \to X^2$, $x \mapsto (x, x)$, is continuous and $\mathbf{\Sigma}^1_1$ is closed under continuous preimages. But A is not analytic because $A \neq U_x$, $\forall x \in X$ (Lemma 2.16), while $\{U_x : x \in X\} = \mathbf{\Sigma}^1_1(X)$. □

4.3 Analytic Separation

Theorem 4.6 (Luzin)

Let X be Polish and $A, B \subseteq X$ be disjoint analytic sets. Then A, B are Borel-separable, i.e. there is a Borel $C \subseteq X$ such that $C \supseteq A$ and $C^c \supseteq B$.




Proof. For a set $P \subseteq X$, observe that the sets which are Borel-separable from P form a σ-ideal, i.e. if each Q_n is Borel-separable from P by a Borel set B_n, then P is Borel-separable from $\bigcup_n Q_n$ by $\bigcup_n B_n$: $\bigcup_n B_n \supseteq \bigcup_n Q_n$ and $(\bigcup_n B_n) \cap P = \bigcup_n (B_n \cap P) = \varnothing$. Hence moreover, if

Source manuscript page 35

P_n, Q_m are Borel-separable for all $m, n \in \mathbb{N}$, then $\bigcup_n P_n$ and $\bigcup_m Q_m$ are Borel-separable: separate P_n from $\bigcup_m Q_m$ for each n, then separate $\bigcup_n P_n$ from $\bigcup_m Q_m$.

Now fix nonempty disjoint analytic sets A, B (if one is empty, then the case is trivial). Then there are continuous surjections $f : \mathbb{N}^{\mathbb{N}} \twoheadrightarrow A$ and $g : \mathbb{N}^{\mathbb{N}} \twoheadrightarrow B$. For $s \in \mathbb{N}^{<\mathbb{N}}$, put $A_s := f([s]_{\mathbb{N}})$, $B_s := g([s]_{\mathbb{N}})$, so $A_s = \bigcup_{n\in\mathbb{N}} A_{sn}$ and $B_s = \bigcup_{n\in\mathbb{N}} B_{sn}$, and $A = A_\varnothing$ and $B = B_\varnothing$. If A, B are not separable, then since $A = A_\varnothing = \bigcup_n A_n$, $B = B_\varnothing = \bigcup_m B_m$, there are $n_0, m_0 \in \mathbb{N}$ such that A_{n_0} and B_{m_0} are not Borel-separable. Further, since $A_s = \bigcup_n A_{sn}$, $B_s = \bigcup_n B_{sn}$, there are $n_1, m_1 \in \mathbb{N}$ such that $A_{n_0 n_1}$ and $B_{m_0 m_1}$ are not Borel-separable. Hence by induction, there are infinite branches $x, y \in \mathbb{N}^{\mathbb{N}}$ such that for every $\ell \in \mathbb{N}$, $A_{x|_\ell}$, $B_{y|_\ell}$ are not Borel-separable.



But $f(x)$ and $g(y)$ are distinct because $f(x) \in A$ and $g(y) \in B$, hence separable by open sets by Hausdorffness, i.e. there are $U, V \subseteq X$ disjoint open such that $U \ni f(x)$ and $V \ni g(y)$. By

Source manuscript page 36

the continuity of f and g and the fact that $([x|_\ell])_\ell$ and $([y|_\ell])_\ell$ form neighborhood bases at x, y respectively, there is $\ell \in \mathbb{N}$ such that $[x|_\ell] \subseteq f^{-1}(U)$, $[y|_\ell] \subseteq g^{-1}(V)$. But then $A_{x|_\ell} = f([x|_\ell])$ and $B_{y|_\ell} = g([y|_\ell])$ are Borel-separable by either U or V. $\Rightarrow\!\!\!\!\!\!\not\Leftarrow$ □

Corollary 4.7 (Souslin)

Let X be Polish. Then $\mathcal{B}(X) = \mathbf{\Delta}^1_1(X)$.

Proof. It is left to prove $\mathbf{\Delta}^1_1(X) \subseteq \mathcal{B}(X)$ (the other inclusion is the observation above). Take $A \in \mathbf{\Delta}^1_1(X)$. Then both A and A^c are analytic. Hence by analytic separation (Theorem 4.6), there is $B \in \mathcal{B}(X)$ such that $B \supseteq A$, $B^c \supseteq A^c$. Thus $B = A$ and $A \in \mathcal{B}(X)$. □

Remark 4.8

This gives a powerful method for proving that a set is Borel, i.e. we just check analyticity and coanalyticity. □

Corollary 4.9

Let X, Y be Polish, $f : X \to Y$. The following are equivalent:

(1) f is Borel, i.e. f-preimages of Borel (equivalently open) sets are Borel.
(2) $\mathrm{graph}(f) \in \mathcal{B}(X \times Y)$, i.e. the graph is Borel.
(3) $\mathrm{graph}(f) \in \mathbf{\Sigma}^1_1(X \times Y)$, i.e. the graph is analytic.

Proof. (1) ⇒ (2) Fix a countable basis $\{V_n\}_{n\in\mathbb{N}}$ for Y and note that $\forall (x,y)\in X\times Y$,

$$\begin{aligned}(x,y)\in \operatorname{graph}(f) &\iff f(x)=y \iff \forall n\in\mathbb{N}\ (y\in V_n \implies x\in f^{-1}(V_n))\\ &\iff \underbrace{\forall n\in\mathbb{N}}_{\text{countable }\cap}\ \big(\ \underbrace{y\notin V_n}_{\text{closed }(V_n^c)}\ \text{ or }\ \underbrace{x\in f^{-1}(V_n)}_{\text{Borel}}\big).\end{aligned}$$


(2) ⇒ (3) is trivial.

(3) ⇒ (1) Assume $\operatorname{graph}(f)$ is analytic and fix a Borel $B\subseteq$

Source manuscript page 37

Y to show that $f^{-1}(B)$ is Borel. For each $x\in X$, we have

$$\begin{aligned}x\in f^{-1}(B) &\iff \exists y\in Y\ (f(x)=y \text{ and } y\in B)\\ &\iff \exists y\in Y\ \big[(x,y)\in\operatorname{graph}(f)\ \wedge\ (x,y)\in X\times B\big].\end{aligned}$$



Thus

$$f^{-1}(B)=\underbrace{\operatorname{proj}_X}_{\text{continuous}}\Big[\ \underbrace{\underbrace{\operatorname{graph}(f)}_{\mathbf{\Sigma}^1_1}\cap\underbrace{(X\times B)}_{=\ \underbrace{\operatorname{proj}_Y^{-1}}_{\text{continuous}}\ \underbrace{(B)}_{\text{Borel}}}}_{\substack{\text{Borel, since } \mathcal{B} \text{ is closed}\\ \text{under continuous preimages}\\ \mathbf{\Sigma}^1_1 \text{ by closure under (countable) finite intersections}}}\ \Big]$$

is analytic, since $\mathbf{\Sigma}^1_1$ is closed under continuous images (Proposition 4.3). But then $f^{-1}(B)^c=f^{-1}(B^c)$ is analytic, so $f^{-1}(B)\in\mathbf{\Delta}^1_1(X)=\mathcal{B}(X)$ is Borel by Souslin's theorem (Corollary 4.7). □

Corollary 4.10

Let X be Polish, $\{A_n\}_{n\in\mathbb{N}}$ a family of pairwise disjoint analytic subsets of X. Then there are pairwise disjoint Borel sets $\{B_n\}_{n\in\mathbb{N}}$ such that $B_n\supseteq A_n$, $\forall n\in\mathbb{N}$.

Proof. Exercise. Hint: Use finite intersections to construct Borel separations for each A_n, then disjointify them. □

4.4 Souslin's Operation $\mathcal{A}$

We now introduce a powerful machinery that equivalently defines the analytic sets. It further allows us to prove that all analytic sets are universally measurable, i.e. measurable with respect to an arbitrary complete σ-finite

Source manuscript page 38

measure μ that measures all Borel sets, i.e. $\mathcal{M} \supseteq \mathcal{B}(X)$, where $\mathcal{M}$ denotes the μ-measurable sets.

Definition 4.11

For a set X and a pruned tree $T \subseteq \mathbb{N}^{<\mathbb{N}}$, we call any sequence $(P_s)_{s\in T}$ of subsets of X a Souslin scheme on X. Call it monotone if $P_t \subseteq P_s$ for all $t \supseteq s$. Call it proper if it is monotone and $P_{si} \cap P_{sj} = \varnothing$, $\forall\, si, sj \in T$, $i \neq j \in \mathbb{N}$. □

Definition 4.12

Souslin's operation $\mathcal{A}$ applied to a Souslin scheme $(P_s)_{s\in T}$ on X results in the set

$$\mathcal{A}(P_s)_{s\in T} := \bigcup_{y\in[T]} \bigcap_{n\in\mathbb{N}} P_{y|_n}. \quad \square$$

Observation. Given a Souslin scheme $(P_s)_{s\in T}$, we can extend T to $\mathbb{N}^{<\mathbb{N}}$ by just setting $P_s := \varnothing$ for $s \in \mathbb{N}^{<\mathbb{N}} \setminus T$. Note that we are not assuming anything about the scheme nor about $\mathcal{A}(P_s)_{s\in T}$. Hence, we might take WLOG our $T = \mathbb{N}^{<\mathbb{N}}$. □

Proposition 4.13

If $(P_s)_{s\in T}$ is a proper Souslin scheme, then

$$\mathcal{A}(P_s)_{s\in T} = \bigcap_{n\in\mathbb{N}} \bigcup_{\substack{s\in T\\ |s|=n}} P_s.$$

Proof. $\subseteq$: If $x \in \mathcal{A}(P_s)_{s\in T}$, there is $y \in [T]$ such that $\forall n$, $x \in P_{y|_n}$. Then $\forall n$, $x \in \bigcup_{s\in T,|s|=n} P_s$.

$\supseteq$: If $x \in \bigcap_{n\in\mathbb{N}} \bigcup_{s\in T,|s|=n} P_s$, then $\forall n$, $\exists!\, s_n \in T$ with $|s_n| = n$ such that $x \in P_{s_n}$, by properness. By monotonicity, $t := s_{n+1}|_n \subseteq s_{n+1}$,

Source manuscript page 39

hence $P_t \supseteq P_{s_{n+1}} \ni x$, thus by uniqueness, $t = s_n$, and $s_n \subseteq s_{n+1}$. Thus defining $y := \bigcup_{n\in\mathbb{N}} s_n \in [T]$ we get $x \in \bigcap_{n\in\mathbb{N}} P_{y|_n} \subseteq \mathcal{A}(P_s)_{s\in T}$. □

Lemma 4.14

If a Souslin scheme $(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$ satisfies $P_s = \bigcup_{i\in\mathbb{N}} P_{si}$, then $\mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}} = P_\varnothing$.

Proof. By definition, $\mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}} \subseteq P_\varnothing$. Let $x \in P_\varnothing$; note that there is $y_0 \in \mathbb{N}$ such that $x \in P_{y_0}$, then there is $y_1 \in \mathbb{N}$ such that $x \in P_{y_0y_1}$, ... By induction we get $y \in \mathbb{N}^{\mathbb{N}}$ such that $x \in P_{y|_n}$, $\forall n \in \mathbb{N}$, so $x \in \mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$. □

Definition 4.15

Let Γ be a class of sets in metrizable topological spaces. For each such space X, denote by $\mathcal{A}\Gamma(X)$ the collection of sets of the form $\mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$ where $P_s \in \Gamma(X)$. The following shows that the operation $\mathcal{A}$ can be viewed as a projection. □

Observation. For a Souslin scheme $(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$ on a set X, $\mathcal{A}(P_s)_s = \mathrm{proj}_X(P)$, where the set $P \subseteq X\times\mathbb{N}^{\mathbb{N}}$ is defined by: for each $(x,y) \in X \times \mathbb{N}^{\mathbb{N}}$,

$$\begin{aligned}(x,y)\in P \;:\iff\; & \forall n\in\mathbb{N},\; x\in P_{y|_n}\\ \iff\; & \forall n\in\mathbb{N}\;\exists s\in\mathbb{N}^{<\mathbb{N}}\;(y|_n = s \;\wedge\; x\in P_s)\\ \iff\; & \forall n\in\mathbb{N}\;\forall s\in\mathbb{N}^{<\mathbb{N}}\;(y|_n = s \implies x\in P_s).\end{aligned}$$

Moreover, if Γ is a class of sets containing all clopen sets and closed under finite unions and countable intersections (e.g. $\Gamma = \mathbf{\Pi}^0_1$ or $\mathbf{\Sigma}^1_1$), then $\mathcal{A}\Gamma \subseteq \exists^{\mathbb{N}^{\mathbb{N}}}\Gamma$ because

$$(x,y)\in P \iff \underbrace{\forall n\in\mathbb{N}}_{\text{countable}\cap}\ \underbrace{\forall s\in\mathbb{N}^{<\mathbb{N}}}_{\text{countable}\cap}\ \underbrace{\big(\underbrace{y|_n\neq s}_{\text{clopen}}\ \vee\ \underbrace{x\in P_s}_{\Gamma}\big)}_{\text{finite}\cup\,\in\,\Gamma}$$

Source manuscript page 40

and $\mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}} = \mathrm{proj}_X(P)$. In particular, $\mathcal{A}\mathbf{\Sigma}^1_1 \subseteq \exists^{\mathbb{N}^{\mathbb{N}}}\mathbf{\Sigma}^1_1 = \mathbf{\Sigma}^1_1$, with the last equality because $\mathbf{\Sigma}^1_1$ is closed under continuous images (proj) and continuous preimages (proj^{-1}). □

We now show $\mathbf{\Sigma}^1_1 \subseteq \mathcal{A}\mathbf{\Sigma}^1_1$ in Polish spaces.

Lemma 4.16

Let X be Polish and $f : \mathbb{N}^{\mathbb{N}} \to X$ be continuous. Then for each $s \in \mathbb{N}^{<\mathbb{N}}$, $P_s := f([s])$ is analytic and

$$f(\mathbb{N}^{\mathbb{N}}) = \mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}} = \mathcal{A}(\overline{P_s})_{s\in\mathbb{N}^{<\mathbb{N}}}.$$

In fact, $(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$ satisfies, for all $s \in \mathbb{N}^{<\mathbb{N}}$:


(i) $P_s \neq \varnothing$;
(ii) $P_s = \bigcup_{i\in\mathbb{N}} P_{si}$ (so that Lemma 4.14 applies);
(iii) for each $y \in \mathbb{N}^{\mathbb{N}}$ and open $U \subseteq X$, if $P_y := \bigcap_{n\in\mathbb{N}} P_{y|_n} \subseteq U$, then there is $n \in \mathbb{N}$ such that $P_{y|_n} \subseteq U$;
(iv) the scheme $(\overline{P_s})_{s\in\mathbb{N}^{<\mathbb{N}}}$ has vanishing diameter with respect to any compatible metric on X.

Proof. We leave the verification of the analyticity of P_s and of (i) and (ii) as exercises. (ii) + Lemma 4.14 $\implies$ $\mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}} = P_\varnothing = f(\mathbb{N}^{\mathbb{N}})$. (iii) is by continuity of f: first notice that $f(y) \in f([y|_n]) = P_{y|_n}$, thus $f(y) \in P_y \subseteq U$. Hence $y \in f^{-1}(U)$, which is open, thus there is $n \in \mathbb{N}$ such that $[y|_n] \subseteq f^{-1}(U)$, i.e. $P_{y|_n} := f([y|_n]) \subseteq U$. (iv) is also by continuity since for $\varepsilon > 0$, there is $n \in \mathbb{N}$ such that $f([y|_n]) \subseteq B_\varepsilon(f(y))$, hence

$$\operatorname{diam}\overline{P_{y|_n}} = \operatorname{diam} P_{y|_n} = \operatorname{diam} f([y|_n]) \leq \operatorname{diam}\left(B_\varepsilon(f(y))\right) \leq 2\varepsilon \to 0.$$

Source manuscript page 41

It is also clear from the definition that $\mathcal{A}(\overline{P_s})_{s\in\mathbb{N}^{<\mathbb{N}}} \supseteq \mathcal{A}(P_s)_s$, hence it remains to show the reverse inclusion. Let $x \in \mathcal{A}(\overline{P_s})_{s\in\mathbb{N}^{<\mathbb{N}}}$, so $x \in \bigcap_{n\in\mathbb{N}} \overline{P_{y|_n}}$ for some $y \in \mathbb{N}^{\mathbb{N}}$. It suffices to show $x = f(y)$, and then $f(y) \in f(\mathbb{N}^{\mathbb{N}}) = P_\varnothing = \mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$. Suppose otherwise; then by Hausdorffness of X, we have disjoint open $U \ni x$, $V \ni f(y)$, thus $x \notin \overline{V}$. By continuity of f, there is $n \in \mathbb{N}$ such that $f([y|_n]) \subseteq V$, hence $x \notin \overline{f([y|_n])} = \overline{P_{y|_n}} \subseteq \overline{V}$, a contradiction.



□

Theorem 4.17 (Characterization of Σ^1_1 via $\mathcal{A}$)

Let X be Polish and $\varnothing \neq A \subseteq X$. The following are equivalent:

(1) A is analytic ($\iff$ there is a continuous $f : \mathbb{N}^{\mathbb{N}} \to X$ such that $f(\mathbb{N}^{\mathbb{N}}) = A$).
(2) $A = \mathcal{A}(F_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$, each $F_s \neq \varnothing$ closed and the scheme $(F_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$ monotone with vanishing diameter.
(3) $A = \mathcal{A}(P_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$, each P_s analytic.

As a result,

$$\mathbf{\Sigma}^1_1 = \mathcal{A}\mathbf{\Sigma}^1_1 = \mathcal{A}\mathbf{\Pi}^0_1.$$

Proof. (1) ⇒ (2): Take $f : \mathbb{N}^{\mathbb{N}} \to X$ continuous with $A = f(\mathbb{N}^{\mathbb{N}})$. Let $F_s := \overline{f([s])}$. Then Lemma 4.16 gives $A = \mathcal{A}(F_s)_{s\in\mathbb{N}^{<\mathbb{N}}}$.

(2) ⇒ (3): Trivial since closed sets are analytic.

(3) ⇒ (1): By the observation above, $\mathcal{A}\mathbf{\Sigma}^1_1 \subseteq \mathbf{\Sigma}^1_1$.

For the last claim, note that $\mathcal{A}\mathbf{\Pi}^0_1 \subseteq \mathcal{A}\mathbf{\Sigma}^1_1$ since closed sets are analytic, while (1) ⇒ (2) gives $\mathbf{\Sigma}^1_1 \subseteq \mathcal{A}\mathbf{\Pi}^0_1$ (with $\varnothing = \mathcal{A}(\varnothing)_{s\in\mathbb{N}^{<\mathbb{N}}}$) and the observation gives $\mathcal{A}\mathbf{\Sigma}^1_1 \subseteq \mathbf{\Sigma}^1_1$. Hence


$$\mathbf{\Sigma}^1_1 \subseteq \mathcal{A}\mathbf{\Pi}^0_1 \subseteq \mathcal{A}\mathbf{\Sigma}^1_1 \subseteq \mathbf{\Sigma}^1_1,$$

so all three classes coincide; in particular $\mathcal{A}\mathbf{\Sigma}^1_1 = \mathcal{A}\mathbf{\Pi}^0_1$. □

Source manuscript page 42

5 Uniformization Theorems

Uniformization theorems are about building nice right inverses to projections, i.e. given $B \subseteq X \times Y$, we'd like a way to uniformly pick a witness $y_x \in Y$ for each $x \in \mathrm{proj}_X(B)$ such that $(x, y_x) \in B$. In particular, given a Borel set $B \subseteq X \times Y$ in a product of Polish spaces, we ask when proj_X admits a Borel right inverse $f : \mathrm{proj}_X(B) \to B$. We will see that $\mathrm{proj}_X(B)$ is Borel in this case as a consequence.



It turns out that this is possible when the projection is "small"-to-one, e.g. one-to-one, countable-to-one, compact-to-one or K_σ (i.e. σ-compact)-to-one.

5.1 Projections and Compactness

Proposition 5.1

(a) Continuous functions map compact sets to compact sets.
(b) Continuous functions map K_σ sets to K_σ sets.
(c) (Tube lemma) For topological spaces X, Y, where Y is compact, $\mathrm{proj}_X : X \times Y \to X$ maps closed sets to closed sets.

Proof. (a), (b) are exercises ((a) is also Part 1, Theorem 4.7(d)).

(c) (HW 3) Let $F \subseteq X \times Y$ be closed. Then for $x \in X$, $x \notin \mathrm{proj}_X(F) \iff \forall y \in Y$, $(x, y) \in F^c$. Since F^c is open, for each $y \in Y$ there are open $U_y \ni x$, $V_y \ni y$ such that $(x, y) \in U_y \times V_y \subseteq F^c$. $\{V_y\}_{y \in Y}$ is an open cover of Y, hence there is a finite subcover $\{V_{y_1}, \ldots, V_{y_n}\}$ of Y. It is left as an exercise to check that $x \in \bigcap_{i=1}^n U_{y_i} \subseteq \mathrm{proj}_X(F)^c$. Hence $\mathrm{proj}_X(F)^c$ is open, thus

Source manuscript page 43

$\mathrm{proj}_X(F)$ is closed. □

Example 5.2

(a) $\mathrm{proj}_0 : \mathbb{R} \times \mathbb{R} \to \mathbb{R}$ does not map closed sets to closed sets. E.g. $B := \mathrm{graph}\left(x \mapsto \frac{1}{x},\ x \neq 0\right)$ is closed in $\mathbb{R} \times \mathbb{R}$, but $\mathrm{proj}_0(B) = \mathbb{R} \setminus \{0\}$ is not closed.
(b) Because $\mathbb{R}$ is σ-compact, so is $\mathbb{R} \times \mathbb{R}$, hence every closed subset of $\mathbb{R} \times \mathbb{R}$ is σ-compact; and proj_0 maps σ-compact sets to σ-compact sets. So proj_0 maps closed sets to σ-compact (in particular F_σ) sets, e.g. $\mathrm{proj}_0(B) = \mathbb{R} \setminus \{0\} = \bigcup_{n \geq 1} \left([-n, -\frac{1}{n}] \cup [\frac{1}{n}, n]\right)$ in (a). □

5.2 The Luzin–Souslin Theorem

Theorem 5.3 (Luzin–Souslin; one-to-one)

Let X, Y be Polish, and $f : X \to Y$ be Borel with $f|_A$ injective for a Borel $A \subseteq X$. Then $f(A)$ is still Borel.

Proof (Ruiyuan Chen). Suppose first $X = A$ and f is continuous, so we need to show that $f(X) \subseteq Y$ is Borel. Let $\mathcal{T}_X, \mathcal{T}_Y$ denote the Polish topologies on X, Y, respectively. Fix a countable basis $\mathcal{U} \subseteq \mathcal{T}_X$ for $\mathcal{T}_X$. For $U \in \mathcal{U}$, by injectivity, $f(U)$ and $f(U^c)$ are disjoint (and analytic), thus by Luzin's analytic separation (Theorem 4.6), there is a Borel set $B_U \subseteq Y$ such that $B_U \supseteq f(U)$ and $B_U^c \supseteq f(U^c)$. Let $\mathcal{T}'_Y \supseteq \mathcal{T}_Y$ be a Polish topology in which all the B_U are clopen, with $\mathcal{B}(\mathcal{T}'_Y) = \mathcal{B}(\mathcal{T}_Y)$ (Corollary 3.4). We now show that $f(X)$ is G_δ in $\mathcal{T}'_Y$; then $f(X) \in \mathcal{B}(\mathcal{T}'_Y) = \mathcal{B}(\mathcal{T}_Y)$.

Notice $\mathrm{graph}(f) \subseteq X \times Y$ is closed in $\mathcal{T}_X \times \mathcal{T}_Y$, hence closed in $\mathcal{T}_X \times \mathcal{T}'_Y$ by refinement. So by making closed-graph functions continuous (Proposition 3.14), the topology $\mathcal{T}'_X := \mathcal{T}_{X,f}$ generated by $f : X \to (Y, \mathcal{T}'_Y)$ over $\mathcal{T}_X$ is still Polish, and $f : (X, \mathcal{T}'_X) \to (Y, \mathcal{T}'_Y)$ is continuous. In fact, f is an open map onto $f(X)$: $\mathcal{T}'_X$ is

Source manuscript page 44

generated by $\mathcal{U} \cup \{f^{-1}(V) : V \in \mathcal{T}'_Y\}$, so every $\mathcal{T}'_X$-open set is a union of sets $U \cap f^{-1}(V)$ with $U \in \mathcal{U}$, $V \in \mathcal{T}'_Y$ (the observation after Definition 3.11). Moreover $f(f^{-1}(V)) = V \cap f(X)$, and $f(U) = B_U \cap f(X)$ by the choice of B_U (and injectivity), thus

$$f(U \cap f^{-1}(V)) = B_U \cap f(X) \cap V$$

is open in $f(X)$. Hence f is an embedding, so $f(X) \cong (X, \mathcal{T}'_X)$ is Polish, therefore G_δ in $(Y, \mathcal{T}'_Y)$ (Part 1, Theorem 2.9). Thus $f(X) \in \mathcal{B}(\mathcal{T}_Y)$.

For general $A \subseteq X$ and Borel f: since the assertion is about $f(A)$ in $\mathcal{T}_Y$, we can safely refine $\mathcal{T}_X$ so that f is continuous and A is clopen (Corollary 3.4, applied to A and the sets $f^{-1}(V)$ for V in a countable basis of Y). Then applying what we just proved to the Polish space A and the continuous injection $f|_A$ gives the desired result. □

Corollary 5.4

(a) If X is a set with Polish topologies $\mathcal{T}_0, \mathcal{T}_1$ on it and $\mathcal{T}_0 \subseteq \mathcal{B}(\mathcal{T}_1)$, then $\mathcal{B}(\mathcal{T}_0) = \mathcal{B}(\mathcal{T}_1)$.

(b) A subset B of a Polish space X is Borel $\iff$ B is a continuous injective image of a closed subset of $\mathbb{N}^{\mathbb{N}}$.

(c) Let X, Y be Polish and $G \subseteq X \times Y$ be a Borel function graph, i.e. $G \in \mathcal{B}(X \times Y)$ such that $\forall x \in X$, $\exists^{\leq 1}$ (there exists at most one) $y \in Y$ with $(x, y) \in G$. In other words, G is the graph of a partial function $f : X \rightharpoonup Y$. Then $\mathrm{dom}(f) \subseteq X$ is Borel.

Proof. (a) Since $\mathcal{T}_0 \subseteq \mathcal{B}(\mathcal{T}_1)$, $\langle \mathcal{T}_0 \rangle_\sigma = \mathcal{B}(\mathcal{T}_0) \subseteq \mathcal{B}(\mathcal{T}_1)$. Consider the injective Borel map $\mathrm{id} : (X, \mathcal{T}_1) \to (X, \mathcal{T}_0)$. Then by Luzin–Souslin, $\forall B \in \mathcal{B}(\mathcal{T}_1)$, $\mathrm{id}(B) = B$ is $\mathcal{T}_0$-Borel. Hence

Source manuscript page 45

$\mathcal{B}(\mathcal{T}_1) = \mathcal{B}(\mathcal{T}_0)$.

(b) By clopenification (Corollary 3.8), ($\Rightarrow$) is known. ($\Leftarrow$) is immediate from the Luzin–Souslin theorem.


(c) $\mathrm{dom}(f) = \mathrm{proj}_X(G)$, which is Borel by Luzin–Souslin since $\mathrm{proj}_X|_G$ is injective. □

5.3 The Luzin–Novikov Theorem and Countable Uniformization

One of the most important theorems in DST is the following.

Theorem 5.5 (Luzin–Novikov)

Let X, Y be Polish spaces and $R \subseteq X \times Y$ be Borel such that every X-fiber of R is countable, i.e. R_x is countable $\forall x \in X$. Then $R = \bigsqcup_{n\in\mathbb{N}} G_n$ for some Borel function graphs G_n; equivalently, $G_n = \mathrm{graph}(\gamma_n)$ for Borel $\gamma_n : B_n \to Y$, with B_n Borel in X.



Proof (Forte Shinko). Deferred to one of the final presentations. □

Corollary 5.6

Countable-to-one Borel images of Borel sets are Borel, i.e. if X, Y are Polish, $B \subseteq X$ is Borel and $f : X \to Y$ is Borel such that $f|_B$ is countable-to-one, then $f(B)$ is Borel.

Proof. Recall that for a map f between Polish spaces, f is Borel $\iff$ $\mathrm{graph}(f)$ is a Borel subset of $X \times Y$ (Corollary 4.9). Hence $C := \{(f(x), x) : x \in B\} \subseteq Y \times X$, the image of $\mathrm{graph}(f) \cap (B \times Y)$ under the swap homeomorphism, is Borel, and its Y-fibers $C_y = \{x \in B : f(x) = y\}$ are countable. So it suffices to show that if $C \subseteq Y \times X$ is Borel with countable

Source manuscript page 46

Y-fibers, then $\mathrm{proj}_Y(C)$ $(= f(B))$ is Borel. Yet by the Luzin–Novikov theorem, $C = \bigsqcup_{n\in\mathbb{N}} C_n$, with each C_n a Borel function graph. But then $\mathrm{proj}_Y(C) = \bigcup_{n\in\mathbb{N}} \mathrm{proj}_Y(C_n)$, with each $\mathrm{proj}_Y(C_n)$ Borel by Luzin–Souslin (Corollary 5.4(c)). □

Corollary 5.7 (Countable uniformization; an equivalent form of Luzin–Novikov)

For Polish spaces X, Y, every countable-to-one Borel function $f : X \to Y$ admits a Borel right inverse $g : f(X) \to X$, with $f(X)$ Borel.

Proof. Let $G := \mathrm{graph}(f)^{-1} := \{(y, x) \in Y \times X : (x, y) \in \mathrm{graph}(f)\}$, a Borel set (Corollary 4.9); thus G has countable Y-fibers since f is countable-to-one. By Luzin–Novikov, $G = \bigsqcup_{n\in\mathbb{N}} G_n$, each G_n a

Borel function graph. Define

$$\begin{aligned} k : f(X) &\to \mathbb{N}, & y &\mapsto \min\{n \in \mathbb{N} \mid y \in \operatorname{proj}_Y G_n\}, \\ g : f(X) &\to X, & y &\mapsto \text{the unique } x \in X \text{ such that } (y,x) \in G_{k(y)}. \end{aligned}$$

Then k is Borel: it suffices to check this with the singletons. For every $n \in \mathbb{N}$, $k^{-1}(\{n\}) = \operatorname{proj}_Y G_n \setminus \bigcup_{i<n} \operatorname{proj}_Y(G_i)$, with each $\operatorname{proj}_Y G_i$ Borel by Luzin–Souslin (Corollary 5.4(c)). Moreover, $f(X) = \bigcup_n \operatorname{proj}_Y G_n$ is Borel.

Lastly, we show g is Borel. Pick $U \subseteq X$ open, then

$$\begin{aligned} y \in g^{-1}(U) &\iff g(y) \in U \\ &\iff \exists\, x \in U \text{ such that } (y,x) \in G_{k(y)} \\ &\iff \exists\, n \in \mathbb{N} \text{ such that } y \in k^{-1}(\{n\}) \ \wedge\ \exists\, x \in U \text{ such that } (y,x) \in G_n \\ &\iff \underbrace{\exists\, n \in \mathbb{N}}_{\text{countable } \cup} \text{ such that } \underbrace{y \in k^{-1}(\{n\})}_{\text{Borel}} \ \wedge\ \underbrace{y \in \operatorname{proj}_Y\left(G_n \cap (Y \times U)\right)}_{\text{Borel}}, \end{aligned}$$

Source manuscript page 47

where $\operatorname{proj}_Y\left(G_n \cap (Y \times U)\right)$ is Borel by Corollary 5.4(c), as $G_n \cap (Y \times U)$ is still a Borel function graph. Thus $f \circ g = \mathrm{id}_{f(X)}$ with g Borel. □

5.4 Countable Borel Equivalence Relations

The following notion is why DST is applicable in dynamics and measured group theory.

Definition 5.8

A countable Borel equivalence relation (CBER) is an equivalence relation E on a Polish space X such that E is Borel as a subset of $X \times X$ and each E-class is countable. □

Example 5.9

If Γ is a countable group and $\Gamma \curvearrowright X$ a Borel action on a Polish space X, i.e. each $\gamma \in \Gamma$ acts as a Borel automorphism of X, then the orbit equivalence relation E_Γ of this action is a CBER, since Γ is countable and by definition of a group action, for $x, y \in X$,

$$x\, E_\Gamma\, y \iff \underbrace{\exists\, \gamma \in \Gamma}_{\text{countable } \cup} \ \underbrace{(\gamma x = y)}_{\text{Borel}},$$

where $\{(x,y) : \gamma x = y\} = \operatorname{graph}(\gamma)$ is Borel, being the graph of the Borel function $\gamma : X \to X$ (Corollary 4.9). For example, take the shift action $\Gamma \curvearrowright 2^\Gamma$, where

$$(\gamma \cdot x)(\delta) := x(\delta\gamma), \qquad \delta, \gamma \in \Gamma,\ x \in 2^\Gamma. \quad \square$$

The following theorem says these are the only examples of CBERs.

Corollary 5.10 (Feldman–Moore theorem)

Every CBER $E \subseteq X^2$ on a Polish space X is a countable union of graphs of Borel involutions on X, i.e. $E = \bigcup_{n\in\mathbb{N}} \operatorname{graph}(\gamma_n)$, where each $\gamma_n : X \to X$ is

Source manuscript page 48

a Borel involution, i.e. $\gamma_n^{-1} = \gamma_n$. In particular, E is the orbit equivalence relation of a Borel action of a countable group Γ generated by involutions. In fact,

$$E \setminus \Delta(X) = \bigsqcup_{n\in\mathbb{N}} (\operatorname{graph}(\gamma_n) \setminus \Delta(X)), \qquad \text{where } \Delta(X) := \{(x,x) \mid x \in X\}. \tag{$\dagger$}$$

Proof. We do the classical proof; a proof with Borel edge-colouring is carried out in HW 4.

Convention. For $R \subseteq X^2$, we denote $R^{-1} := \{(y,x) : (x,y) \in R\}$. We view a Borel (partial) function as both a function and its graph, interchangeably: a Borel function graph $f \subseteq X^2$ is at the same time the partial function $f : X \rightharpoonup X$ with domain $\operatorname{dom}(f) := \operatorname{proj}_0(f)$ and range $\operatorname{ran}(f) := \operatorname{proj}_1(f)$, where $\operatorname{proj}_0, \operatorname{proj}_1 : X^2 \to X$ are the projections onto the first and the second coordinate; so $(x,y) \in f$ means $f(x) = y$. When the f_n are written in a union (or an intersection, or a difference), that means taking the union of their graphs in X^2. Sometimes, when we need to emphasize that we are thinking of the graph, we write $\operatorname{graph}(f_n)$. In particular, f^{-1} always denotes the inverse relation; when f is injective, it is (the graph of) the inverse function.

By Corollary 5.4(c), $\operatorname{dom}(f)$ is Borel and f is a Borel function on it. Its range $\operatorname{ran}(f)$ is analytic in general, but it is Borel if f is injective, by Luzin–Souslin again (Theorem 5.3).

Step 1 (injective pieces). By Luzin–Novikov (Theorem 5.5; each fiber E_x is an E-class, hence countable), we can write E as a disjoint union of Borel partial functions, $E = \bigsqcup_{n\in\mathbb{N}} f_n$. We will use these partial functions to build a set of Borel involutions of X. Notice

$$E = E^{-1} = \bigsqcup_{m\in\mathbb{N}} f_m^{-1}$$

since E is symmetric, where f_m^{-1} is the inverse relation of f_m (the inverse function need not exist yet). One can verify that the partial function $f_n \cap f_m^{-1}$ is injective and Borel: if $(x,y),(x',y) \in f_n \cap f_m^{-1}$, then $(y,x),(y,x') \in f_m$, so $x = f_m(y) = x'$. Since $E = E \cap E^{-1} = \bigsqcup_{n,m\in\mathbb{N}}(f_n \cap f_m^{-1})$, WLOG we

Source manuscript page 49

replace $\{f_n\}_n$ by $\{f_n \cap f_m^{-1}\}_{n,m}$ (re-indexed by $\mathbb{N}$) and assume the f_n's are injective with $E = \bigsqcup_{n\in\mathbb{N}} f_n$.

Step 2 (disjoint domains and ranges). Next, we want to extend each f_n to a Borel involution of X. If $\operatorname{dom}(f_n) \cap \operatorname{ran}(f_n) = \varnothing$, then this can be easily done:

$$\tilde{f}_n(x) := \begin{cases} f_n(x) & \text{if } x \in \operatorname{dom}(f_n), \\ f_n^{-1}(x) & \text{if } x \in \operatorname{ran}(f_n), \\ x & \text{otherwise,} \end{cases} \qquad \text{for } n \in \mathbb{N}. \tag{$*$}$$

Then this is clearly an involution, and it is Borel since f_n, f_n^{-1} and $x \mapsto x$ are Borel with Borel domains.

We now make $\operatorname{dom}(f_n)$ and $\operatorname{ran}(f_n)$ disjoint as follows. Pick a countable basis $\mathcal{W} = (W_k)_{k\in\mathbb{N}}$ of X; thus $\mathcal{W}^2 := (W_k \times W_l)_{k,l\in\mathbb{N}}$ is a countable basis of X^2. Let $(x,y) \in X^2$ with $x \neq y$. Then

there are disjoint basic open sets $U_{(x,y)} \ni x$, $V_{(x,y)} \ni y$ in $\mathcal{W}$, hence $U_{(x,y)} \times V_{(x,y)} \in \mathcal{W}^2$ and $(x,y) \in U_{(x,y)} \times V_{(x,y)} \subseteq X^2 \setminus \Delta(X)$ by disjointness. Hence

$$X^2 \setminus \Delta(X) = \bigcup_{(x,y) \notin \Delta(X)} \underbrace{U_{(x,y)} \times V_{(x,y)}}_{\in \mathcal{W}^2}.$$

But $\mathcal{W}^2$ is countable, thus the union is countable: $X^2 \setminus \Delta(X) = \bigcup_{m \in \mathbb{N}} U_m \times V_m$ with $U_m \cap V_m = \varnothing$. Hence

$$E = \Delta(X) \cup \Big(E \cap \bigcup_m (U_m \times V_m)\Big) = \Delta(X) \cup \bigcup_{n,m \in \mathbb{N}} (f_n \cap (U_m \times V_m)).$$


Every graph in the latter union has its domain ($\subseteq U_m$) and range ($\subseteq V_m$) disjoint. Again, replacing $\{f_n\}_{n \in \mathbb{N}}$ with $\{f_n \cap (U_m \times V_m)\}_{n,m \in \mathbb{N}}$, we assume each Borel injective f_n has disjoint domain and range, and now

$$E = \Delta(X) \sqcup \bigcup_{n \in \mathbb{N}} f_n.$$

(The union of the f_n's need not be disjoint any more; the union with $\Delta(X)$ is, since every f_n misses the diagonal.)

Source manuscript page 50

Step 3 (the involutions; $E = E_\Gamma$).



Extend each f_n into a Borel involution $\tilde{f}_n =: \gamma_n$ as in $(*)$. Let Γ be the group generated by $(\gamma_n)_n$ under composition. Then the natural action $\Gamma \curvearrowright X$ is Borel. Furthermore, since E is reflexive and symmetric, $\operatorname{graph}(\gamma_n) = f_n \cup f_n^{-1} \cup \{(x,x) : x \notin \operatorname{dom}(f_n) \cup \operatorname{ran}(f_n)\} \subseteq E$, so by our construction

$$E = \Delta(X) \cup \bigcup_{n \in \mathbb{N}} \operatorname{graph}(f_n) = \Delta(X) \cup \bigcup_{n \in \mathbb{N}} \operatorname{graph}(\gamma_n),$$
$$\text{and} \quad E_\Gamma = \bigcup_{\gamma \in \Gamma} \{(x, \gamma(x)) : x \in X\} = \bigcup_{\gamma \in \Gamma} \operatorname{graph}(\gamma).$$

Lastly, we show $E = E_\Gamma$. Clearly $E \subseteq E_\Gamma$ since $(\gamma_n)_n$ generates Γ and $\Delta(X) = \operatorname{graph}(\mathrm{id}_X)$ for $\mathrm{id}_X \in \Gamma$. Conversely, take an element $\gamma = \gamma_{n_1} \circ \gamma_{n_2} \circ \cdots \circ \gamma_{n_k} \in \Gamma$ (every element has this form, as each generator is its own inverse). Then for any $x \in X$, γ acts on x by

$$x \xmapsto{\gamma_{n_k}} \gamma_{n_k}(x) \longmapsto \cdots \xmapsto{\gamma_{n_1}} \gamma(x).$$

But each step is an equivalence in E since $\operatorname{graph}(\gamma_n) \subseteq E$ for the generators, thus by transitivity of E, $x \mathrel{E} \gamma(x)$. Thus every pair $(x, \gamma(x)) \in E_\Gamma$ is in E, hence $E_\Gamma = E$.

Step 4 (the disjoint version (†)). It remains to show (†). However, with our construction of the γ_n and the symmetry of E together, we can easily see that $\bigcup_n (\operatorname{graph}(\gamma_n) \setminus \Delta(X))$ can never be a disjoint union (unless $E = \Delta(X)$): if $(x,y) \in f_n$, then $(y,x) \in E \setminus \Delta(X)$ lies in some f_m, with $m \neq n$

as $y \notin \mathrm{dom}(f_n)$; so $(y,x) \in \mathrm{graph}(\gamma_n) \cap \mathrm{graph}(\gamma_m)$. We fix it as follows. Before turning f_n into γ_n, we have $E = \Delta(X) \sqcup \bigcup_n f_n$ (Step 2) with each f_n a Borel partial function that is injective and $\mathrm{dom}(f_n) \cap \mathrm{ran}(f_n) = \varnothing$. Thus $E \setminus \Delta(X) = \bigcup_{n\in\mathbb{N}} f_n$. We now do the same trick as in countable uniformization (Corollary 5.7): set

$$c : E \setminus \Delta(X) \to \mathbb{N}, \qquad (x,y) \mapsto \text{the least } n \in \mathbb{N} \text{ such that } (x,y) \in f_n.$$

Source manuscript page 51

c is Borel since $c^{-1}(\{n\}) = f_n \setminus \bigcup_{i<n} f_i$. Symmetrize c by $k(x,y) := \min\{c(x,y), c(y,x)\}$. The map $(x,y) \mapsto c(y,x)$ is c composed with the swap homeomorphism, hence Borel; thus k is Borel. Hence each class $M_n := k^{-1}(\{n\})$ is a symmetric Borel subset of $E \setminus \Delta(X)$. Moreover, for all $x \in X$, there are no $y \neq z$ with $(x,y),(x,z) \in M_n$: if not, then by definition of k and M_n,

$$n = \left\{\begin{matrix} c(x,y) \\ \text{or} \\ c(y,x) \end{matrix}\right\} = \left\{\begin{matrix} c(x,z) \\ \text{or} \\ c(z,x) \end{matrix}\right\}.$$

All 4 cases are impossible:

- $c(x,y) = c(x,z) = n$ means $(x,y),(x,z) \in \mathrm{graph}(f_n)$, i.e. $f_n(x) = y$ and $f_n(x) = z$ with $y \neq z$, impossible;
- $(y,x),(z,x) \in \mathrm{graph}(f_n)$ is impossible since f_n is injective;
- $(y,x),(x,z) \in \mathrm{graph}(f_n)$ or $(x,y),(z,x) \in \mathrm{graph}(f_n)$ are impossible since $\mathrm{dom}(f_n) \cap \mathrm{ran}(f_n) = \varnothing$.

Lastly, we redefine the involutions: let $\gamma_n(x) :=$ the unique y with $(x,y) \in M_n$ if it exists, else x. It is an involution by symmetry of M_n, and Borel since

$$\mathrm{graph}(\gamma_n) = M_n \cup \{(x,x) : x \notin \mathrm{proj}_0(M_n)\}$$

is Borel in X^2 ($\mathrm{proj}_0(M_n)$ is Borel by Corollary 5.4(c)). Now $\mathrm{graph}(\gamma_n) \setminus \Delta(X) = M_n$; $E_\Gamma = E$ for the group Γ generated by these γ_n, in the same way as before (Step 3: $\mathrm{graph}(\gamma_n) \subseteq E$ and $E = \Delta(X) \cup \bigsqcup_n M_n$); and

$$E \setminus \Delta(X) = \bigcup_{n\in\mathbb{N}} (\mathrm{graph}(f_n) \setminus \Delta(X)) = \bigsqcup_{n\in\mathbb{N}} M_n = \bigsqcup_{n\in\mathbb{N}} (\mathrm{graph}(\gamma_n) \setminus \Delta(X)). \qquad \square$$

5.5 Further Uniformization Theorems

Lastly, we have the most powerful uniformization theorems available, which we won't prove.

Theorem 5.11 (Arsenin–Kunugui)

Let X, Y be Polish and $B \subseteq X \times Y$ be

Source manuscript page 52

Borel with every X-fiber B_x being K_σ. Then

(1) $\mathrm{proj}_X(B)$ is Borel (analog of Luzin–Souslin, Theorem 5.3);
(2) $\mathrm{proj}_X|_B$ has a Borel right inverse $\mathrm{proj}_X(B) \to B$ (analog of countable uniformization, Corollary 5.7).

Theorem 5.12 (Saint Raymond)

Let X, Y be Polish and $B \subseteq X \times Y$ be Borel with every X-fiber B_x being K_σ. Then $B = \bigcup_{n\in\mathbb{N}} B_n$, where each B_n is Borel and every X-fiber $(B_n)_x$ is compact (analog of the Luzin–Novikov theorem, Theorem 5.5).

Source manuscript page 53

6 Standard Borel Spaces

6.1 The Borel Isomorphism Theorem

We will prove that any two uncountable Polish spaces are Borel isomorphic.

Corollary 6.1 (Borel Cantor–Schröder–Bernstein)

Let X, Y be Polish and $f : X \hookrightarrow Y$, $g : Y \hookrightarrow X$ be Borel injections. Then X, Y are Borel isomorphic.

Proof. The usual proof for CSB (monotone map $\to$ fixed set $\to$ bijection) fails to show Borel-ness. We implement a fix to it instead of doing the usual Knaster–Tarski construction.

For $A \subseteq X$, put $\varphi(A) := X \setminus g(Y \setminus f(A))$. Then $\varphi : \mathcal{P}(X) \to \mathcal{P}(X)$ is monotone, i.e. for $A \subseteq B \subseteq X$, $\varphi(A) \subseteq \varphi(B)$.

Furthermore, φ preserves Borel sets: let A be Borel. Then $f(A)$ is Borel by Luzin–Souslin (Theorem 5.3), and so is $Y \setminus f(A)$. Thus $g(Y \setminus f(A))$ is Borel by Luzin–Souslin again, and so is its complement $\varphi(A)$.

Lastly, φ commutes with countable intersections: first notice that $f(\bigcap_n A_n) = \bigcap_n f(A_n)$ (exercise: use injectivity of f). Then, since images commute with unions,

$$\begin{aligned}\varphi\Big(\bigcap_n A_n\Big) &= X \setminus g\Big(Y \setminus f\Big(\bigcap_n A_n\Big)\Big) = X \setminus g\Big(Y \setminus \bigcap_n f(A_n)\Big)\\ &= X \setminus g\Big(\bigcup_n (Y \setminus f(A_n))\Big) = X \setminus \bigcup_n g(Y \setminus f(A_n))\\ &= \bigcap_n \Big(X \setminus g(Y \setminus f(A_n))\Big) = \bigcap_n \varphi(A_n).\end{aligned}$$

Let $X_n := \varphi^n(X)$. Then $(X_n)_n$ is decreasing: $X_1 \subseteq X = X_0$, and then by monotonicity, $X_2 = \varphi(X_1) \subseteq \varphi(X_0) = X_1$, and the

Source manuscript page 54

rest follows by induction. Since φ preserves Borel sets, all the X_n are Borel by induction. Put $A_0 := \bigcap_n X_n$, which is Borel as a countable intersection of Borel sets.

Now observe that A_0 is a fixed point: since φ commutes with countable intersections,

$$\varphi(A_0) = \bigcap_n \varphi(X_n) = \bigcap_n X_{n+1} = A_0$$

($\bigcap_n X_n = \bigcap_n X_{n+1}$ since the X_n decrease). What remains is the same construction as in the usual CSB: define $h : X \to Y$ by

$$h(x) := \begin{cases} f(x), & x \in A_0, \\ g^{-1}(x), & x \in X \setminus A_0. \end{cases}$$

This is clearly well-defined and bijective.

Let $C \subseteq Y$ be Borel. Then notice that

$$h^{-1}(C) = (A_0 \cap f^{-1}(C)) \cup g(C \setminus f(A_0)),$$

where A_0 is Borel as discussed, $f^{-1}(C)$ is Borel by assumption, and $f(A_0)$ is Borel by Luzin–Souslin, hence $g(C \setminus f(A_0))$ is Borel by Luzin–Souslin again. Thus $h^{-1}(C)$ is Borel, hence h is Borel. Given a Borel bijection, its inverse is immediately Borel by Luzin–Souslin again. Hence X is Borel isomorphic to Y by h. □

Theorem 6.2 (Borel Isomorphism Theorem)

Any two uncountable Polish spaces are Borel isomorphic.

Proof. Let X be an uncountable Polish space. By Cantor–Bendixson (Part 1, Theorems 5.5 and 5.6), there is a topological embedding $f : 2^{\mathbb{N}} \hookrightarrow X$; and by binary representation (Part 1, Proposition 2.6), there is an injection $b : X \hookrightarrow 2^{\mathbb{N}}$ such that for the

Source manuscript page 55

clopen sets $[n \mapsto 1] := \{x \in 2^{\mathbb{N}} : x(n) = 1\}$, $b^{-1}([n \mapsto 1])$ is open. It is obvious that for the basic open cylinder sets U in $2^{\mathbb{N}}$, $b^{-1}(U)$ is Borel, hence b is Borel. Thus by Borel CSB (Corollary 6.1), X is Borel isomorphic to $2^{\mathbb{N}}$. □

Definition 6.3

A measurable space $(X, \mathcal{S})$ is called a standard Borel space if $\mathcal{S} = \mathcal{B}(\mathcal{T})$ for some Polish topology $\mathcal{T}$ on X. □

Corollary 6.4

There is a unique uncountable standard Borel space up to Borel isomorphism, namely $2^{\mathbb{N}}$.

Proof. Apply the Borel Isomorphism Theorem (Theorem 6.2). □

Example 6.5

(a) For a Polish space X, $(X, \mathcal{B}(X))$ is standard Borel.

(b) Let X be Polish and $B \subseteq X$ be Borel. Then $(B, \mathcal{B}(X)|_B)$ is standard Borel, where $\Gamma(X)|_B := \{C \cap B : C \in \Gamma(X)\}$. In particular, since B is Borel,

$$\mathcal{B}(X)|_B = \{C \cap B : C \subseteq X \text{ Borel}\} = \{C \in \mathcal{B}(X) : C \subseteq B\}. \quad \square$$

Proof of (b). Let $\mathcal{T}$ be the Polish topology on X and $\mathcal{T}' \supseteq \mathcal{T}$ be a Polish topology making B clopen with $\mathcal{B}(\mathcal{T}') = \mathcal{B}(\mathcal{T})$ (Theorem 3.1). Then $(B, \mathcal{T}'|_B)$ is Polish, so $(B, \mathcal{B}(\mathcal{T}'|_B))$ is standard Borel. But $\mathcal{B}(\mathcal{T}'|_B) = \mathcal{B}(\mathcal{T}')|_B = \mathcal{B}(\mathcal{T})|_B$. □

Proposition 6.6

Example 6.5(b) characterizes Borel sets, i.e. if X is Polish, $A \subseteq X$ and $(A, \mathcal{B}(X)|_A)$ is standard Borel, then $A \subseteq X$ is Borel.

Proof. Note that this together with Example 6.5(b) gives HW 4, Q6.

Source manuscript page 56

By standard-Borel-ness, there is a Polish topology $\mathcal{T}$ on A such that $\mathcal{B}(\mathcal{T}) = \mathcal{B}(X)|_A$. Define $\iota : (A, \mathcal{T}) \hookrightarrow X$ by the identity map. For Borel $C \subseteq X$,

$$\iota^{-1}(C) = C \cap A \in \mathcal{B}(X)|_A = \mathcal{B}(\mathcal{T}).$$

Thus ι is a Borel injection, and A is Borel in $\mathcal{B}(X)|_A = \mathcal{B}(\mathcal{T})$. Hence $\iota(A) = A$ is Borel in X by the Luzin–Souslin theorem (Theorem 5.3). □

6.2 The Effros Borel Space

Definition 6.7

For a Polish space X, let $\mathcal{F}(X)$ denote the set of all closed subsets of X. The Effros Borel space of X is the measurable space $(\mathcal{F}(X), \mathcal{E})$, where $\mathcal{E}$ is the σ-algebra generated by the sets of the form

$$[U] := \{F \in \mathcal{F}(X) : F \cap U \neq \varnothing\}, \quad \text{for all open } U \subseteq X. \quad \square$$

Theorem 6.8

For every Polish space X, its Effros Borel space is standard Borel.

Proof. We use binary representation again. Let $(U_n)_{n\in\mathbb{N}}$ be a basis of nonempty open sets for the Polish topology on X, and define $c : \mathcal{F}(X) \to 2^{\mathbb{N}}$ by $F \mapsto (F \cap U_n \neq \varnothing)_{n\in\mathbb{N}}$, i.e. the n-th bit is 1 if $F \cap U_n \neq \varnothing$, else 0.

This map is injective: if $F_0 \neq F_1$ are closed sets in X, then at least one of $F_0 \setminus F_1$ and $F_1 \setminus F_0$ is nonempty, say $F_0 \setminus F_1 \neq \varnothing$. Then for all $x \in F_0 \setminus F_1$, there is a U_n such that $x \in U_n \subseteq F_1^c$, i.e. $U_n \cap F_1 = \varnothing$. Hence $c(F_0)(n) = 1$ and $c(F_1)(n) = 0$, i.e. $c(F_0) \neq$

Source manuscript page 57

$c(F_1)$.

We verify that c is Borel. Notice that the cylinders $([n \mapsto 1])_{n\in\mathbb{N}}$ generate $\mathcal{B}(2^{\mathbb{N}})$. Hence it suffices to check that $c^{-1}([n \mapsto 1])$ is Borel for every $n \in \mathbb{N}$. But

$$c^{-1}([n \mapsto 1]) = \{F \subseteq X : F \text{ closed}, \ F \cap U_n \neq \varnothing\} = [U_n],$$

which is Effros Borel by definition.

Furthermore, c maps Effros Borel sets to Borel sets relative to the image $c(\mathcal{F}(X))$: the $[U_n]$'s generate the Effros Borel σ-algebra since for every open $U \subseteq X$, $[U] = \bigcup_{U_n \subseteq U} [U_n]$. The map c commutes with arbitrary unions, and it commutes with taking complements relative to $c(\mathcal{F}(X))$. Lastly, notice that $c([U_n]) = [n \mapsto 1] \cap c(\mathcal{F}(X))$. Hence c maps the generating sets into the Borel σ-algebra $\mathcal{B}(2^{\mathbb{N}})|_{c(\mathcal{F}(X))}$, thus c maps all the Effros Borel sets to Borel sets in $\mathcal{B}(2^{\mathbb{N}})|_{c(\mathcal{F}(X))}$, as c commutes with complements and intersections.

Denote the image $c(\mathcal{F}(X)) =: \mathcal{C}$. We have shown that $(\mathcal{F}(X), \mathcal{E})$ is Borel isomorphic to $(\mathcal{C}, \mathcal{B}(2^{\mathbb{N}})|_{\mathcal{C}})$. If $\mathcal{C}$ is a Borel subset of $2^{\mathbb{N}}$, then by Example 6.5(b), $(\mathcal{C}, \mathcal{B}(2^{\mathbb{N}})|_{\mathcal{C}})$ is standard Borel, say $\mathcal{B}(2^{\mathbb{N}})|_{\mathcal{C}} = \mathcal{B}(\mathcal{T}_{\mathcal{C}})$ for a Polish topology $\mathcal{T}_{\mathcal{C}}$ on $\mathcal{C}$; and with $\mathcal{T} := \{c^{-1}(V) : V \in \mathcal{T}_{\mathcal{C}}\}$, $(\mathcal{F}(X), \mathcal{T})$ is homeomorphic to $(\mathcal{C}, \mathcal{T}_{\mathcal{C}})$, hence Polish, and $\mathcal{B}(\mathcal{T}) = c^{-1}(\mathcal{B}(2^{\mathbb{N}})|_{\mathcal{C}}) = \mathcal{E}$, since c is a

Source manuscript page 58

Borel isomorphism.

Hence it remains to show $\mathcal{C} \in \mathcal{B}(2^{\mathbb{N}})$. To do so, it is HW 4, Q7 to check the following. Fix a complete compatible metric d on X. Then for all $y \in 2^{\mathbb{N}}$,

$$
\begin{aligned}
y \in c(\mathcal{F}(X)) &\iff y = c(F) \text{ for some } F \in \mathcal{F}(X) \\
&\overset{\circledast}{\iff} (\forall n, m \in \mathbb{N} \text{ with } U_n \subseteq U_m,\ y(n) = 1 \Rightarrow y(m) = 1) \wedge \qquad (\dagger) \\
&\quad \left[\forall U_n\ \forall \varepsilon \in \mathbb{Q}^+,\ y(n) = 1 \Rightarrow (\exists\, \overline{U_m} \subseteq U_n \text{ with } \operatorname{diam}(U_m) < \varepsilon,\ y(m) = 1)\right] \qquad (*) \\
&\iff \underbrace{\left[\forall n, m \in \mathbb{N} \text{ such that } U_n \subseteq U_m,\ \underbrace{(y(n) = 0 \ \vee\ y(m) = 1)}_{\text{clopen}}\right]}_{\text{closed}} \wedge \\
&\quad \underbrace{\left[\forall U_n\ \forall \varepsilon \in \mathbb{Q}^+,\ \underbrace{y(n) = 0}_{\text{clopen}} \vee \underbrace{(\exists\, \overline{U_m} \subseteq U_n \text{ such that } \operatorname{diam}(U_m) < \varepsilon \wedge \underbrace{y(m) = 1}_{\text{clopen}})}_{\text{open}}\right]}_{G_\delta}.
\end{aligned}
$$


So $\mathcal{C}$ is the intersection of a closed set and a G_δ set, hence G_δ, in particular Borel. (So $\mathcal{C}$ is even Polish in the subspace topology by Part 1, Theorem 2.9, and we may take $\mathcal{T}_{\mathcal{C}}$ to be the subspace topology, i.e. $\mathcal{T} = \{c^{-1}(V \cap \mathcal{C}) : V \subseteq 2^{\mathbb{N}} \text{ open}\}$.)

We show the key equivalence $\circledast$.

($\Rightarrow$) If $y = c(F)$, then if $U_n \subseteq U_m$, $y(n) = 1 \Rightarrow y(m) = 1$ because $\varnothing \neq U_n \cap F \subseteq U_m \cap F$. For U_n and $\varepsilon \in \mathbb{Q}^+$, if $y(n) = 1$, then $U_n \cap F \neq \varnothing$. Let $x \in U_n \cap F$. Then there is $0 < \delta < \varepsilon/2$ with $B_\delta(x) \subseteq U_n$. Then, same as in the proof of the Baire category theorem (Part 1, Theorem 8.14), there is a U_m such that

$$x \in U_m \subseteq \overline{U_m} \subseteq B_{\delta/2}[x] \subseteq B_\delta(x) \subseteq U_n,$$

and $\operatorname{diam}(U_m) \leq \operatorname{diam}(B_\delta(x)) \leq 2\delta < \varepsilon$, with $y(m) = 1$.



($\Leftarrow$) We want to show that there is a closed F such that $F \cap U_n = \varnothing$ iff $y(n) = 0$. Set

$$F := X \setminus \bigcup_{\substack{n\in\mathbb{N}\\ y(n)=0}} U_n,$$

then F is closed, and $y(n) = 0 \Rightarrow F \cap U_n = \varnothing$. On the other hand, suppose $y(n) = 1$. By $(*)$, we

Source manuscript page 59

construct a sequence

$$U_n \supseteq \overline{U_{m_0}} \supseteq U_{m_0} \supseteq \overline{U_{m_1}} \supseteq U_{m_1} \supseteq \overline{U_{m_2}} \supseteq \cdots,$$

with $y(m_j) = 1$ for all j and $\operatorname{diam}(U_{m_j}) = \operatorname{diam}(\overline{U_{m_j}}) < 1/j$ for $j \in \mathbb{N}^+$. By HW 1, Q2,

$$\bigcap_{j\in\mathbb{N}}^{\downarrow} U_{m_j} = \bigcap_{j\in\mathbb{N}}^{\downarrow} \overline{U_{m_j}} = \{x\}$$

for some $x \in X$. Now $x \in F$: if $x \in U_{\tilde{n}}$ with $y(\tilde{n}) = 0$, then $\bigcap_{j\in\mathbb{N}} U_{m_j} \subseteq U_{\tilde{n}}$ gives $j \in \mathbb{N}$ such that $U_{m_j} \subseteq U_{\tilde{n}}$, but $y(\tilde{n}) = 0$ and $y(m_j) = 1$ contradicts $(\dagger)$. Thus $U_n \cap F \neq \varnothing$ since $x \in U_n \cap F$. □

Example 6.9 (Application)

Recall that every Polish space is homeomorphic to a closed subset of $\mathbb{R}^{\mathbb{N}}$ (Part 1, Theorem 4.11), so it can be regarded as a point in $\mathcal{F}(\mathbb{R}^{\mathbb{N}})$. Thus we can regard $\mathcal{F}(\mathbb{R}^{\mathbb{N}})$ as the standard Borel space of all Polish spaces, and consider homeomorphism between Polish spaces as an equivalence relation on $\mathcal{F}(\mathbb{R}^{\mathbb{N}})$. But it is not an analytic equivalence relation, i.e. it is not analytic as a subset of $\mathcal{F}(\mathbb{R}^{\mathbb{N}})^2$ in the following sense; see Example 6.11. □

Definition 6.10

We call a subset of a standard Borel space $(X, \mathcal{S})$ analytic if it is analytic ($\mathbf{\Sigma}^1_1$, Definition 4.1) with respect to some (equivalently, every) compatible Polish topology, i.e. a Polish topology $\mathcal{T}$ on X with $\mathcal{B}(\mathcal{T}) = \mathcal{S}$: if $\mathcal{T}, \mathcal{T}'$ are both compatible Polish topologies for X, then $\mathrm{id} : (X, \mathcal{T}) \to (X, \mathcal{T}')$ is a Borel isomorphism, and the class $\mathbf{\Sigma}^1_1$ is closed under Borel images and preimages (Proposition 4.3). □

Example 6.11 (Homeomorphism is not analytic)

Every class of an analytic equivalence relation is analytic, so it suffices to find one non-analytic class. We claim that $\{F : F \cong \mathbb{N}\}$ is not analytic. We work in $\mathcal{F}(\mathbb{N}^{\mathbb{N}})$: a closed embedding $e : \mathbb{N}^{\mathbb{N}} \hookrightarrow \mathbb{R}^{\mathbb{N}}$ (Part 1, Theorem 4.11) induces a Borel map $F \mapsto e(F)$, $\mathcal{F}(\mathbb{N}^{\mathbb{N}}) \to \mathcal{F}(\mathbb{R}^{\mathbb{N}})$, preserving homeomorphism type, since the preimage of $[U]$ is $[e^{-1}(U)]$.

Identify trees on $\mathbb{N}$ with points of $2^{\mathbb{N}^{<\mathbb{N}}}$, and let WF be the set of well-founded trees, i.e. the trees W with $[W] = \varnothing$. For a tree W on $\mathbb{N}$, let $W^\star := \{t\star^k : t \in W,\ k \in \mathbb{N}\}$, a pruned tree on $\mathbb{N} \sqcup \{\star\} \cong \mathbb{N}$, and $X_W := [W^\star] = [W] \sqcup \{t\star^\infty : t \in W\}$. Then:

(1) $W \mapsto X_W$ is Borel into the Effros space, since $W^\star$ is pruned and so $X_W \cap [s]_{\mathbb{N}} \neq \varnothing \iff s \in W^\star$, a clopen condition on W.

(2) Each $t\star^\infty$ is isolated in X_W (it is the only point of X_W in $[t\star]_{\mathbb{N}}$), while each $x \in [W]$ is the limit of $x|_n\star^\infty$.

So X_W is discrete iff $W \in \mathrm{WF}$, and after padding W to be infinite without changing well-foundedness (replace W by $\{\varnothing\} \cup \{(n)^\frown t : n \in \mathbb{N},\ t \in W\}$), $X_W \cong \mathbb{N}$ iff $W \in \mathrm{WF}$. Hence WF is a Borel preimage of $\{F : F \cong \mathbb{N}\}$, which therefore cannot be analytic (Proposition 4.3), as WF is $\mathbf{\Pi}^1_1$-complete. □

We can also select points from closed sets in a Borel fashion.

Source manuscript page 60

Definition 6.12

For a Polish space X, call a map $s : \mathcal{F}(X)^* \to X$, where $\mathcal{F}(X)^* := \mathcal{F}(X) \setminus \{\varnothing\}$, a selector if $s(F) \in F$ for each $F \in \mathcal{F}(X)^*$. We say that s is a Borel selector if it is a Borel map. □

Example 6.13

(a) For $X = \mathbb{R}$, the following s is a Borel selector: $s : \mathcal{F}(\mathbb{R})^* \to \mathbb{R}$ by

$$F \mapsto \min\left([-M, M] \cap F\right), \quad \text{where } M := \inf\{N > 0 : [-N, N] \cap F \neq \varnothing\},$$

or

$$F \mapsto \begin{cases} \min\{d \geq 0 : d \in F\}, \\ \max\{d < 0 : d \in F\} & \text{if the min does not exist.} \end{cases}$$

It is an exercise to check that this map is well-defined and Borel.

(b) For $X := \mathbb{N}^\mathbb{N}$, the following is a Borel selector: let $s : \mathcal{F}(\mathbb{N}^\mathbb{N})^* \to \mathbb{N}^\mathbb{N}$ by $F \mapsto$ the leftmost branch in T_F, where $T_F := \{x|_n : x \in F\} \subseteq \mathbb{N}^{<\mathbb{N}}$ is the tree of F. By leftmost, we mean the $x \in [T_F]$ recursively defined by $x(n) :=$ the least $j \in \mathbb{N}$ with $(x|_n)^\frown j \in T_F$, where $x|_n$ is already defined by induction; it exists since T_F is pruned. □

Theorem 6.14 (Borel axiom of choice for closed sets)

For every Polish space X, the Effros Borel space $\mathcal{F}(X)$ admits a Borel selector.

Proof. By HW 2, Q10, there is a continuous open surjection $f : \mathbb{N}^\mathbb{N} \to X$, so $f^{-1}(F)$ is closed for each $F \in \mathcal{F}(X)$. This gives a map $\tilde{f} : \mathcal{F}(X) \to \mathcal{F}(\mathbb{N}^\mathbb{N})$, which is Borel since f is open.

Source manuscript page 61

Hence if $s_{\mathbb{N}^\mathbb{N}} : \mathcal{F}(\mathbb{N}^\mathbb{N})^* \to \mathbb{N}^\mathbb{N}$ is a Borel selector, then $f \circ s_{\mathbb{N}^\mathbb{N}} \circ \tilde{f} : \mathcal{F}(X)^* \to X$ is a Borel selector. The details are filled in HW 4, Q10. □