Defitions and notations

Zhehao Wang

October 29, 2018

0.1 Limit

Let f(x) be a function defined on an interval that contains x = a, except possibly at x = a, then we say that

$$\lim_{x \to a} f(x) = L$$

if for every $\epsilon > 0$ there is some number $\delta > 0$ such that

$$|f(x) - L| < \epsilon$$
 whenever $0 < |x - a| < \delta$

0.2 Gradient

Given $f(x_1, x_n)$ on \mathbf{R}^n

$$\nabla f(a_1,...,a_n) = (\frac{\partial}{\partial x_1}(a_1,...,a_n),...,\frac{\partial}{\partial x_n}(a_1,...,a_n))$$

0.3 Directional derivative

Homework notation

$$f'(x; u) = \lim_{h \to 0} \frac{f(x + hu) - f(x)}{h}$$

Wikipedias notation note that it's f on a vector \vec{x} having its derivative calculated on direction \vec{v} , essentially gradient on direction \vec{v} .

$$\nabla_v f(\vec{x}) = \lim_{h \to 0} \frac{f(\vec{x} + h\vec{v}) - f(\vec{x})}{h}$$

0.4 Vector norm

A norm is a function that assigns a strictly positive length or size to each vector in a vector space (except the zero vector which is assigned a length of 0).

Absolute value norm is a norm on the one-dimensional vector spaces formed by real or complex numbers.

$$||x|| = |x|$$

Euclidean norm on a Euclidean space \mathbb{R}^n is such

$$\|\vec{x}\|_2 = \sqrt{x_1^2 + \ldots + x_n^2}$$

Manhattan or taxicab norm

$$\|\vec{x}\|_1 = \sum_{i=1}^n |x_i|$$

p-norm

$$\|\vec{x}\|_p = (\sum_{i=1}^n |x_i|^p)^{\frac{1}{p}}$$

Note that when p=1, we get Manhattan norm, and when p=2, we get Euclidean norm.

When $p = \infty$

$$\|\vec{x}\|_{\infty} = \max_{i} |x_i|$$

0.5 argmax

Points of the domain of some function at which the function values are maximized.

Given an arbitrary set X, a totally ordered set Y and a function $f: X \to Y$, the arg max over some subset S of X is defined by

$$\underset{x \in S \subseteq X}{\arg\max} f(x) = \{ x \mid x \in S \land \forall y \in S : f(y) \le f(x) \}$$