



# Comprehensive Phenotyping of T Cells Using Flow Cytometry

Charlotte M. Mousset,  Willemijn Hobo,  Rob Woestenenk, Frank Preijers, Harry Dolstra,   
Anniek B. van der Waart\* 

Department of Laboratory Medicine -  
Laboratory of Hematology, Radboud  
Institute of Molecular Life Sciences,  
Radboud University Medical Center,  
Nijmegen, The Netherlands

Received 14 November 2018; Revised 9  
January 2019; Accepted 15  
January 2019

Additional Supporting Information may  
be found in the online version of this  
article.

\*Correspondence to: Anniek B. van der  
Waart, PhD, Geert Grooteplein Zuid  
8, 6525 GA Nijmegen, The Netherlands  
Email: anniek.  
vanderwaart@radboudumc.nl

Published online 4 February 2019 in  
Wiley Online Library  
(wileyonlinelibrary.com)

DOI: 10.1002/cyto.a.23724

© 2019 International Society for  
Advancement of Cytometry

## • Abstract


The T cell compartment can form a powerful defense against extrinsic (e.g., pathogens) and intrinsic danger (e.g., malignant cells). At the same time, specific subsets of T cells control this process to keep the immune system in check and prevent autoimmunity. A wide variety in T cell functionalities exists, which is dependent on the differentiation and maturation state of the T cells. In this review, we report an overview for the identification of CD4<sup>+</sup> T-αβ cells (T-helper (Th)1, Th2, Th9, Th17, Th22, and CD4<sup>+</sup> regulatory T cells), CD8<sup>+</sup> T-αβ cells (cytotoxic T lymphocyte (Tc)1, Tc2, Tc9, Tc17, and CD8<sup>+</sup> regulatory T cells), and their additional effector memory status (naïve, stem cell memory, central memory, effector memory, and effector) using flow cytometry. These different subsets can be discriminated based on selective extracellular markers, in combination with intracellular transcription factor and/or cytokine stainings. Additionally, identification of very small subsets, including antigen-specific T cells, and important technical considerations of flow cytometry are discussed. Together, this overview can be used for comprehensive phenotyping of a T cell subset of interest. © 2019 International Society for Advancement of Cytometry

## • Key terms

T cell; phenotyping; flow cytometry; differentiation; T cell subset

**T** cells appear in different sorts and flavors, with distinct characteristics that can be visualized through flow cytometry. T cells originate from lymphoid progenitors and have the capacity to traffic from the bone marrow to the thymus. Here T-cell development takes place comprising T-cell receptor (TCR)-mediated selection and maturation into naïve T cells (Fig. 1) (1,2). Lymphopoiesis starts with progenitor cells expressing CD2, CD5, CD7, and cytoplasmic CD3, which have the capacity to enter the cortex of the thymus (3,4). At this point, TCRs are formed after rearrangements of α, β, γ, and δ chains resulting in T cells with γδ-chains (T-γδ) and T cells with αβ-chains (T-αβ). These T-γδ and T-αβ cells can be characterized by flow cytometric gating for CD3 and the γδ-TCR or αβ-TCR, respectively (5–8). At this stage, NK-T cells originate from the CD3-expressing precursor T cells by expressing a specific α-chain (Vα14–Jα18) which pairs via a β-chain interaction with glycolipid-CD1d. Upon development NK-T cells acquire expression of the neural cell adhesion molecule (NCAM, CD56) (9). A CD1d tetramer together with CD56 antibody, can be used to study NK-T cells (10). Meanwhile, precursor T cells express CD3 at low levels on their surface and go through a rapid transition, where first CD8 and then CD4 expression is acquired. As expression of CD8 can occur prior to or after rearrangement of the TCRs, T-γδ cells can also separate from the αβ-path after up-regulation of CD8, which results in approximately 30% the T-γδ cells being CD8<sup>+</sup> (8). The CD4<sup>+</sup>CD8<sup>+</sup> double positive (DP) T-αβ cells continue their development by undergoing positive selection via interaction with either peptide:MHC class I complexes resulting in single expression of CD8 or peptide:MHC class II complexes resulting in single expression of CD4 (11). Thereafter, T-αβ migrate from the thymic





**Figure 1.** Discriminating T- $\alpha\beta$  cells from their relatives. T- $\alpha\beta$  cells arise from CD34<sup>+</sup> hematopoietic stem cells which acquire CD2, CD5, and CD7 expression during lymphopoiesis. After migration from the bone marrow to the thymus, CD3 is expressed and T- $\gamma\delta$  cells and NK-T cells deviate from the T- $\alpha\beta$ -path. T- $\gamma\delta$  cells express the  $\gamma\delta$ -TCR and NK-T cells express a specific  $\alpha\beta$ -TCR which pairs with glycolipid-CD1d and they express CD56. The T- $\alpha\beta$  cells gain both CD4 and CD8 and subsequent to positive and negative clonal selection, single positive CD4<sup>+</sup>, and CD8<sup>+</sup> T- $\alpha\beta$  cells are released into the blood. Double negative (DN) T- $\alpha\beta$  cells are negative for both CD4 and CD8.

cortex to the medulla, where they undergo negative clonal selection to remove T cells that have a high affinity interaction for self-antigens. Finally, mature single-positive CD4<sup>+</sup> and CD8<sup>+</sup> T- $\alpha\beta$  cells are released into the blood (12–15). A small percentage (~1%) of the peripheral blood T- $\alpha\beta$  cells does not express CD4 nor CD8, and are known as double negative (DN) T- $\alpha\beta$  cells (16–18).

Mature T- $\alpha\beta$  cell subsets can be further characterized by comprehensive phenotyping, keeping in mind that differentiation often shows gradual changes in expression patterns, and molecules previously described as subset specific (e.g., CD56),

are also expressed by other immune cells (19). In this review we provide an overview of extracellular proteins, intracellular transcription factors, and effector molecules that can be used to characterize different populations of T- $\alpha\beta$  cells.

#### T- $\alpha\beta$ CELL SUBSETS CAN BE DEFINED BY EXTRACELLULAR PROTEINS, TRANSCRIPTION FACTORS, AND CYTOKINE PROFILES

The most common strategy to segregate CD4<sup>+</sup> T- $\alpha\beta$  helper (Th) cell subsets [i.e., Th1, Th2, Th9, Th17, Th22, and


**Table 1.** A comprehensive phenotype of cytokine-based subsets within CD4<sup>+</sup> and CD8<sup>+</sup> T- $\alpha\beta$  cells

	CD4						CD8				
	TH1	TH2	TH9	TH17	TH22	TREG	TC1	TC2	TC9	TC17	TREG
<i>Extracellular</i>											
CCR10					+						
CD25 (IL-2R $\alpha$ )						+					+
CD127 (IL-7R $\alpha$ )						+/-					
CD152 (CTLA4)						+					+
CD161				+						+	
CD183 (CXCR3)	+						+				
CD194 (CCR4)		+		+	+			+		-	
CD196 (CCR6)	-	-		+	+				+	+	
CD294 (CRTH2)		+						+			
IL23R				+						+	
<i>Intracellular</i>											
AHR					+						
Foxp3						+					+
FOXO4					+						
GATA3		+						+			
IRF4		+	+	+			+		+	+	
PU.1			+								
ROR $\gamma$ T				+						+	
T-bet	+						+			-	
<i>Intracellular upon cell stimulation</i>											
IFN $\gamma$	+	-		-	-		+	-			
IL2	+						+				
IL4	-	+		-	-	+	-	+/-			
IL5		+					-	+			
IL9		+	+	+	-				+		
IL10		+			-	+			+		+
IL13		+						+			
IL17				+	-					+	
IL21				+						+	
IL22				+	+						
TNF- $\alpha$	+	+					+				
TGF- $\beta$						+					+
References	(20–22,24)	(20–23,30)	(30)	(8,21,22,28)	(21,25)	(22,26,31)	(29)	(23,29)	(29)	(27–29)	(29)

Boxes marked negative (–) indicate antigens, transcription factors or cytokines which are not expressed or produced by the corresponding subset. An empty box indicates that this is not used for the discrimination of a specific subset.

regulatory T cells (Tregs)] is based on their functionality. In the last decade, new research highlighted that this segregation can be applied to CD8<sup>+</sup> T- $\alpha\beta$  cells as well, separating cytotoxic T lymphocytes (Tc)1, Tc2, Tc9, Tc17, and CD8<sup>+</sup> Tregs. The division is mainly based on extracellular markers, selective intracellular transcription profiles, and the cytokine production profile of the T cells. To define the various T- $\alpha\beta$  subpopulations, a phenotype based on TCR- $\alpha\beta$ , CD4, and CD8 staining should be combined with CD183 (CXCR3; Th1 and Tc1), CD194 (CCR4) and CD294 (CRTH2; both for Th2 and Tc2), CD196 (CCR6) or CD161 (KLRB1; both for Th17 and Tc17), CCR10 (Th22), and CD25 (IL-2R $\alpha$ ) and CD152 (CTLA4; both for CD4<sup>+</sup> and CD8<sup>+</sup> Tregs) (Table 1 and Fig. 2) (20–31). However, so far no distinct extracellular marker for

Th9 and Tc9 cells has been identified. For better discrimination, subsequent intracellular transcription factor profiling can be performed. These include T-bet (Th/c1), GATA3 (Th/c2), PU.1 (Th9), ROR $\gamma$ t (Th/c17), AHR, and/or FOXO4 (Th22) and FoxP3 (CD4/CD8<sup>+</sup> Tregs) (20–30). Selective transcription factors for Tc9 have not yet been found. Nevertheless, the best method to distinguish between the subsets is based on their cytokine secretion profile upon stimulation in the presence of a Golgi system inhibitor, like Brefeldin A or Monensin, to prevent cytokine release. Following subsequent fixation and permeabilization, cells can be stained for intracellular IFN $\gamma$  and IL2 (Th1/Tc1), IL4 and IL5 (Th2/Tc2), IL9 (Th9/Tc9), IL10 (CD4<sup>+</sup>/CD8<sup>+</sup> Treg), IL17 (Th17/Tc17), and IL22 (Th22) (20–30).



**Figure 2.** Identification of CD4<sup>+</sup> and CD8<sup>+</sup> T-αβ subsets. A combination of both surface and intracytoplasmic stainings for cytokines and transcription factors should be used for the correct identification of CD4<sup>+</sup> T-αβ helper (Th) cell subsets Th1, Th2, Th9, Th17, Th22, and Treg, and CD8<sup>+</sup> cytotoxic T-αβ (Tc) cell subsets Tc1, Tc2, Tc9, Tc17, and CD8<sup>+</sup> Tregs. Discriminative surface proteins are visualized on the cell surface, cytokines intracellular as circles and transcription factors intracellular as squares.

### CHARACTERIZATION OF EFFECTOR MEMORY DIFFERENTIATION

T-αβ cell subsets can be further characterized based on their effector memory differentiation status. Either via a linear or partly progressive model, stem cell memory (T<sub>SCM</sub>), central memory (T<sub>CM</sub>), effector memory (T<sub>EM</sub>), and effector T cells (T<sub>EFF</sub>) develop from the naive (T<sub>N</sub>) repertoire upon activation (32–37). Effective T cell activation requires interaction of the TCR-CD3 complex with the peptide:MHC complex on antigen-presenting cells, co-stimulatory signaling and cytokine support. Every effector/memory subset has unique characteristics which provide them with differential migratory capacity, longevity, and functionality. While CD197<sup>+</sup>CD62L<sup>+</sup>

T<sub>N</sub>, T<sub>SCM</sub>, and T<sub>CM</sub> home to lymphoid tissues in search of their specific antigen being presented by antigen-presenting cells, CD197<sup>+</sup>CD62L<sup>+</sup> T<sub>EM</sub> and T<sub>EFF</sub> patrol peripheral tissues and blood and efficiently produce effector molecules upon antigen encounter (36–38). Notably, early memory T cells (T<sub>SCM</sub> and T<sub>CM</sub>) are known for their longevity and high proliferative capacity, which gradually decreases upon differentiation (36). Although the expression of various extracellular markers changes during activation and differentiation (Table 2), one can already discriminate the different CD4<sup>+</sup> and CD8<sup>+</sup> effector memory T cell subsets using a small set of markers, including CD45RA or CD45RO in combination with CD197 (CCR7) and CD95 (Fig. 3A). Additionally, often the

**Table 2.** A comprehensive phenotype of memory differentiation subsets within CD4<sup>+</sup> and CD8<sup>+</sup> T- $\alpha\beta$  cells


	CD4				CD8				REFERENCES
	T <sub>N</sub>	T <sub>SCM</sub>	T <sub>CM</sub>	T <sub>EM/EFF</sub>	T <sub>N</sub>	T <sub>SCM</sub>	T <sub>CM</sub>	T <sub>EM/EFF</sub>	
<i>Extracellular</i>									
CD11a (LFA-1)	−/+	+	+	+	−	+	+	++	(44,45)
CD27	+++	+++	++	+/−	+++	+++	++	+/−	(40,44,46–51)
CD28	++	+++	+++	+/−	++	+++	+++	+/−	(40,44,46–48)
CD38	−	+	+	+	+++	++	++	+	(37,44,52)
<b>CD45RA</b>	+	+	−	−	+	+	−	+/−	(46–51,53,54)
<b>CD45RO</b>	−	−	+	+/−	−	−	+	+/−	(44,46–49,51,53,54)
CD57	−*	−*	+	++	−*	−*	+	++	(40)
CD58 (LFA-3)	−	+	++	++	−	+	++	++	(44,48,49)
CD62L (L-selectin)	+++	++	++	−	+++	+++	++	−	(38,44,46–49)
CD69	−*	−*	+	+	−*	−*	+	+	(38,55)
<b>CD95 (FAS)</b>	−	++	+++	++	−	++	+++	++	(37,44,46–51)
CD122 (IL-2Rβ)	−	+	+	++	−	+	+	++	(37,44,46–49)
CD127 (IL-7Rα)	++	++	++	+/−	+	++	++	+	(44,47,53,56)
CD130	+++	++	++	+	+++	++	++	+	(37)
CD134 (OX40)	+*	+*	+++	++	+*	+*	++	++	(40)
CD183 (CXCR3)	−	+	+	++	−	++	++	+/−	(37,38,44,48,50,57,58)
CD184 (CXCR4)	+++	++	++	+	+	+++	+++	++	(38,44)
CD195 (CCR5)	−	+	+	+++	−	+	++	+++	(37,50)
<b>CD197 (CCR7)</b>	+++	+++	++	−	+++	+++	++	−	(38,44,46–51)
CD218a (IL-18Rα)	−	+	++	+++	−	++	++	+++	(37)
CD244 (2B4)	−*	−*	−	+	−*	−*	−	+	(40)
CD278 (ICOS)	+*	+*	++	++	+*	+*	++	+	(40)
HLA-DR	−	+	+	++	−	+	+	++	(37,38)
<i>Intracellular</i>									
ACTN1					+++	++	+	−	(44)
EOMES	−	+	+	++	−	+	++	+++	(24,44,46)
FOXO1					+	+	+	−	(59)
FOXP1	+++	++	++	+	+++	++	++	+	(32,44,46)
ID2					+	+	++	+++	(46)
ID3					+++	+++	++	+	(46)
IL6ST					+++	+	+	−	(44)
LASS6					+++	++	+	−	(44)
LEF1	+++	++	+	−	+++	++	+	−	(32,44,46)
KLF7					+++	++	+	−	(46)
PRDM1	−	+	++	+++	−	+	++	+++	(32,44,46)
PRF1	−	−			−	−	+	+++	(44,50)
TAF4B					+++	++	+	−	(44)
TBX21	−	+	+	+++	−	+	+	+++	(32,44,46)
TCF7	+++	+++	++	+	+++	+++	++	+	(32,46)
Zeb2	−	+	++	+++	−	+	++	+++	(32,46)
<i>Intracellular upon stimulation</i>									
IFNγ	−	+	++	+++	−	+	++	+++	(38,44,46,47,50,58)
IL-2	−/+	+++	+++	+	−/+	+++	++	+/−	(38,46,47,58)
TNFα	−	−/+	++	+++	−	+	++	+++	(46,47,50)
GZMA	−	+		++	−	+	+	+++	44,50,60,61
GZMB	−	+/−		++	−	−*	−*	++	(50,60–62)
Perforin				++	−	+*	+*	+++	(60,61)

\*No discrimination between T<sub>N</sub> and T<sub>SCM</sub> cells. Empty boxes are not specifically described for the indicated subset.

cell adhesion molecule and homing receptor L-selectin (CD62L, Fig. 2B), co-stimulatory molecules CD27 and CD28, and cytokine receptors IL-2R $\beta$  (CD122) and IL-7R $\alpha$  (CD127) are included in effector memory differentiation analyses. Upon stimulation, besides expression of co-stimulatory markers, activated T- $\alpha\beta$  cells can also start to express co-inhibitory receptors, including CTLA-4, PD-1, TIGIT, and TIM-3 (39,40). While this prevents sustained T cell activation via a natural feedback loop, it can also hamper

immune responses in patients with chronic viral infections or cancer (41).

The analysis of all the described subsets should be always preceded by a proper gating strategy which includes the exclusion of doublets using a single cell gate (42) (Fig. 3C), dead cells with a life/dead stain (Fig. 3D), and, when necessary, autofluorescent cells using an empty fluorescence channel. This is especially important when small (low frequent) populations including T<sub>SCM</sub> and antigen-specific T cells are analyzed. Antigen-specific CD8<sup>+</sup> T- $\alpha\beta$  cells can be visualized using single or double (combinatorial encoding) tetramer or pentamer stainings (Fig. 3E). Tetramers and pentamers consist of respectively 4 or 5 biotinylated peptide-MHC complexes linked to a fluorescent label. The use of double tetramer staining with two different fluorescent labels eliminates the chance of false positive results, thereby increasing the analysis specificity. Combinatorial staining with 6 different fluorochromes results in 15 unique color combinations, and makes it possible to identify up to 15 different epitopes with a high specificity (43). Additionally, in particular when performing deep multiparametric phenotypes, also Fluorescence Minus One (FMO) controls should be taken into consideration. An FMO control includes cells stained with all fluorochromes of the panel except one, and can be used to verify the cut-off point of background fluorescence and thereby the positive population (Fig. 3F) after optimum concentrations have been established via an antibody titration assay. Finally, single stains on the cells of interest should be performed to set the appropriate compensations followed by a check of compensation settings when using the complete panel. It is essential to synchronize flow cytometer settings over time and also harmonize your analyzers to acquire identical data on different machines. The right settings, together with accurate gating and a complete flow cytometry panel can be used for comprehensive phenotyping of your T-cell subset of interest.



**Figure 3.** Gating for flow cytometry. Peripheral blood mononuclear cells were resuspended in PBS containing 0.5% bovine serum albumin followed by extracellular staining for 30 min at 4°C in the dark. (A) Gating strategy to discriminate CD4<sup>+</sup> and CD8<sup>+</sup> T cell memory populations within single, live, CD56<sup>+</sup>TCR $\gamma\delta$ <sup>+</sup>CD3<sup>+</sup> cells. TN-like cells are CD45RO<sup>+</sup>CCR7<sup>+</sup>, TCM cells are CD45RO<sup>+</sup>CCR7<sup>+</sup> and TEM/EFF cells are CD45RO<sup>+</sup>CCR7<sup>-</sup>. The discrimination between TN and TSCM within TN-like was based on the CD197-differentiation pattern of total CD4<sup>+</sup> or CD8<sup>+</sup> T cells (– →). (B) Gating strategy to discriminate CD4<sup>+</sup> and CD8<sup>+</sup> T cell memory populations with CD45RO, CD62L, and CD27 within single, live, CD56<sup>+</sup>TCR $\alpha\beta$ <sup>+</sup>CD3<sup>+</sup> cells. TN-like cells are CD45RO<sup>+</sup>CD62L<sup>+</sup>CD27<sup>+</sup>, TCM cells are CD45RO<sup>+</sup>CD62L<sup>+</sup>CD27<sup>+</sup> and TEM/EFF cells are CD45RO<sup>+</sup>CD62L<sup>+</sup>CD27<sup>-</sup>. (C) Discrimination of single cells by plotting forward scatter width (FS-W) against forward scatter area (FS-A) as well as plotting side scatter width (SS-W) against side scatter area (SS-A). Events deviating from the linear cell population are called doublets. (D) Live cells can be evaluated by gating on viability dye-negative cells. (E) Antigen-specific CD8<sup>+</sup> T cells are analyzed by using antigen-specific tetramers, labeled with two different fluorescent labels. (F) Fluorescence minus one (FMO) control for CD45RO expression within FL3 contains all antibodies except CD45RO-ECD. This is used to determine the position of CD45RO<sup>+</sup> cells



# LITERATURE CITED

- Schwarz BA, Bhandoola A. Trafficking from the bone marrow to the thymus: A prerequisite for thymopoiesis. *Immunol Rev* 2006;209:47–57.
- Lind EF, Prockop SE, Porritt HE, Petrie HT. Mapping precursor movement through the postnatal thymus reveals specific microenvironments supporting defined stages of early lymphoid development. *J Exp Med* 2001;194(2):127–134.
- Terstappen LW, Huang S, Picker LJ. Flow cytometric assessment of human T-cell differentiation in thymus and bone marrow. *Blood* 1992;79(3):666–677.
- Laribi K, Lemaire P, Sandrini J, Baugier de Materre A. Advances in the understanding and management of T-cell prolymphocytic leukemia. *Oncotarget* 2017;8(61):104664–104686.
- Dudley EC, Petrie HT, Shah LM, Owen MJ, Hayday AC. T cell receptor beta chain gene rearrangement and selection during thymocyte development in adult mice. *Immunity* 1994;1(2):83–93.
- Passoni L, Hoffman ES, Kim S, Crompton T, Pao W, Dong MQ, Owen MJ, Hayday AC. Intrathymic delta selection events in gammadelta cell development. *Immunity* 1997;7(1):83–95.
- Garcillan B, Marin AV, Jimenez-Reinoso A, Briones AC, Munoz-Ruiz M, et al. Gammadelta T lymphocytes in the diagnosis of human T cell receptor Immunodeficiencies. *Front Immunol* 2015;6:20.
- Kalyan S, Kabelitz D. Defining the nature of human gammadelta T cells: A biographical sketch of the highly empathetic. *Cell Mol Immunol* 2013;10(1):21–29.
- Cowan JE, Jenkinson WE, Anderson G. Thymus medulla fosters generation of natural Treg cells, invariant gammadelta T cells, and invariant NKT cells: What we learn from intrathymic migration. *Eur J Immunol* 2015;45(3):652–660.
- Snyder-Cappione JE, Tincati C, Eccles-James IG, Cappione AJ, Ndhlovu LC, Koth LL, Nixon DF. A comprehensive ex vivo functional analysis of human NKT cells reveals production of MIP1-alpha and MIP1-beta, a lack of IL-17, and a Th1-bias in males. *PLoS One* 2010;5(11):e15412.
- Levelt CN, Carsetti R, Eichmann K. Regulation of thymocyte development through CD3. II. Expression of T cell receptor beta CD3 epsilon and maturation to the CD4+8+ stage are highly correlated in individual thymocytes. *J Exp Med* 1993;178(6):1867–1875.
- Ueno T, Saito F, Gray DH, Kuse S, Hieshima K, et al. CCR7 signals are essential for cortex-medulla migration of developing thymocytes. *J Exp Med* 2004;200(4):493–505.
- Kwan J, Killeen N. CCR7 directs the migration of thymocytes into the thymic medulla. *J Immunol* 2004;172(7):3999–4007.
- Yamano T, Nedjic J, Hinterberger M, Steinert M, Koser S, Pinto S, Gerdes N, Lutgens E, Ishimaru N, Busslinger M, et al. Thymic B cells are licensed to present self antigens for central T cell tolerance induction. *Immunity* 2015;42(6):1048–1061.
- Hadeiba H, Lahl K, Edalati A, Oderup C, Habtezion A, Pachynski R, Nguyen L, Ghodsi A, Adler S, Butcher EC. Plasmacytoid dendritic cells transport peripheral antigens to the thymus to promote central tolerance. *Immunity* 2012;36(3):438–450.
- Liu G, Li Z, Wei Y, Lin Y, Yang C, Liu T. Direct detection of FoxP3 expression in thymic double-negative CD4-CD8- cells by flow cytometry. *Sci Rep* 2014;4:5781.
- Fischer K, Voelkl S, Heymann J, Przybylski GK, Mondal K, Laumer M, Kunz-Schughart L, Schmidt CA, Andreesen R, Mackensen A. Isolation and characterization of human antigen-specific TCR alpha beta+ CD4-CD8- double-negative regulatory T cells. *Blood* 2005;105(7):2828–2835.
- Martina MN, Noel S, Saxena A, Rabb H, Hamad AR. Double negative (DN) alphabeta T cells: Misperception and overdue recognition. *Immunol Cell Biol* 2015;93(3):305–310.
- Van Acker HH, Capsomidis A, Smits EL, Van Tendeloo VF. CD56 in the immune system: More than a marker for cytotoxicity? *Front Immunol* 2017;8:892.
- Li H, Rostami A. IL-9: Basic biology, signaling pathways in CD4+ T cells and implications for autoimmunity. *J Neuroimmune Pharmacol* 2010;5(2):198–209.
- Annunziato F, Romagnani S. Heterogeneity of human effector CD4+ T cells. *Arthritis Res Ther* 2009;11(6):257.
- Beriou G, Bradshaw EM, Lozano E, Costantino CM, Hastings WD, Orban T, Elyaman W, Khoury SJ, Kuchroo VK, Baecher-Allan C, et al. TGF-beta induces IL-9 production from human Th17 cells. *J Immunol* 2010;185(1):46–54.
- Cosmi L, Annunziato F, Galli MIG, Maggi RME, Nagata K, Romagnani S. CRTH2 is the most reliable marker for the detection of circulating human type 2 Th and type 2 T cytotoxic cells in health and disease. *Eur J Immunol* 2000;30(10):2972–2979.
- Chemin K, Ramskold D, Diaz-Gallo LM, Herrath J, Houtman M, et al. EOMES-positive CD4(+) T cells are increased in PTPN22 (1858T) risk allele carriers. *Eur J Immunol* 2018;48(4):655–669.
- Eyerich S, Eyerich K, Pennino D, Carbone T, Nasorri F, Pallotta S, Cianfarani F, Odorisio T, Traidl-Hoffmann C, Behrendt H, et al. Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest* 2009;119(12):3573–3585.
- Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. *Science* 2003;299(5609):1057–1061.
- Kondo T, Takata H, Matsuki F, Takiguchi M. Cutting edge: Phenotypic characterization and differentiation of human CD8+ T cells producing IL-17. *J Immunol* 2009;182(4):1794–1798.
- Maggi L, Santarlasci V, Capone M, Peired A, Frosali F, Crome SQ, Querci V, Fambrini M, Liotta F, Levings MK, et al. CD161 is a marker of all human IL-17-producing T-cell subsets and is induced by RORC. *Eur J Immunol* 2010;40(8):2174–2181.
- Mittrucker HW, Visekruna A, Huber M. Heterogeneity in the differentiation and function of CD8(+) T cells. *Arch Immunol Ther Exp (Warsz)* 2014;62(6):449–458.
- Wong MT, Ye JJ, Alonso MN, Landrigan A, Cheung RK, Engleman E, Utz PJ. Regulation of human Th9 differentiation by type I interferons and IL-21. *Immunol Cell Biol* 2010;88(6):624–631.
- Liu W, Putnam AL, Xu-Yu Z, Szot GL, Lee MR, et al. CD127 expression inversely correlates with FoxP3 and suppressive function of human CD4+ T reg cells. *J Exp Med* 2006;203(7):1701–1711.
- Durek P, Nordstrom K, Gasparoni G, Salhab A, Kressler C, et al. Epigenomic profiling of human CD4(+) T cells supports a linear differentiation model and highlights molecular regulators of memory development. *Immunity* 2016;45(5):1148–1161.
- van Aalderen MC, van den Biggelaar M, Remmerswaal EBM, van Alphen FPI, Meijer AB, ten Berge IJM, van Lier RAW. Label-free analysis of CD8(+) T cell subset proteomes supports a progressive differentiation model of human-virus-specific T cells. *Cell Rep* 2017;19(5):1068–1079.
- Verbist KC, Guy CS, Milasta S, Liedmann S, Kaminski MM, et al. Metabolic maintenance of cell asymmetry following division in activated T lymphocytes. *Nature* 2016;532(7599):389–393.
- Herndler-Brandstetter D, Ishigame H, Shinnakasu R, Plajer V, Stecher C, et al. KLRG1(+) effector CD8(+) T cells lose KLRG1, differentiate into all memory T cell lineages, and convey enhanced protective immunity. *Immunity* 2018;48(4):716–729.
- Lugli E, Dominguez MH, Gattinoni L, Chattopadhyay PK, Bolton DL, Song K, Klatt NR, Brenchley JM, Vaccari M, Gostick E, et al. Superior T memory stem cell persistence supports long-lived T cell memory. *J Clin Invest* 2013;123(2):594–599.
- Lugli E, Gattinoni L, Roberto A, Mavilio D, Price DA, Restifo NP, Roederer M. Identification, isolation and in vitro expansion of human and nonhuman primate T stem cell memory cells. *Nat Protoc* 2013;8(1):33–42.
- Sallusto F, Lenig D, Forster R, Lipp M, Lanzavecchia A. Two subsets of memory T lymphocytes with distinct homing potentials and effector functions. *Nature* 1999;401(6754):708–712.
- Blackburn SD, Shin H, Haining WN, Zou T, Workman CJ, Polley A, Betts MR, Freeman GJ, Vignali DAA, Wherry EJ. Coregulation of CD8+ T cell exhaustion by multiple inhibitory receptors during chronic viral infection. *Nat Immunol* 2009;10(1):29–37.
- Hutten TJA, Norde WJ, Woestenenk R, Wang RC, Maas F, Kester M, Falkenburg JHF, Berglund S, Luznik L, Jansen JH, et al. Increased Coexpression of PD-1, TIGIT, and KLRG-1 on tumor-reactive CD8(+) T cells during relapse after allogeneic stem cell transplantation. *Biol Blood Marrow Transplant* 2018;24(4):666–677.
- Norde WJ, Maas F, Hobo W, Korman A, Quigley M, Kester MGD, Hebeda K, Falkenburg JHF, Schaap N, de Witte TM, et al. PD-1/PD-L1 interactions contribute to functional T-cell impairment in patients who relapse with cancer after allogeneic stem cell transplantation. *Cancer Res* 2011;71(15):5111–5122.
- Snow CBK. Usefulness of DNA fluorescence ratioing in reducing DNA measurement artifacts caused by aggregates. *Cytometry A* 1994;7A:50.
- Hombirnk P, Hadrup SR, Bakker A, Kester MG, Falkenburg JH, et al. High-throughput identification of potential minor histocompatibility antigens by MHC tetramer-based screening: Feasibility and limitations. *PLoS One* 2011;6(8):e22523.
- Gattinoni L, Lugli E, Ji Y, Pos Z, Paulos CM, Quigley MF, Almeida JR, Gostick E, Yu Z, Carpenito C, et al. A human memory T cell subset with stem cell-like properties. *Nat Med* 2011;17(10):1290–1297.
- Walling BL, Kim M. LFA-1 in T cell migration and differentiation. *Front Immunol* 2018;9(952). <https://doi.org/10.3389/fimmu.2018.00952>.
- Gattinoni L, Klebanoff CA, Restifo NP. Paths to stemness: Building the ultimate antitumor T cell. *Nat Rev Cancer* 2012;12(10):671–684.
- Rosenblum MD, Way SS, Abbas AK. Regulatory T cell memory. *Nat Rev Immunol* 2016;16(2):90–101.
- Chahroudi A, Silvestri G, Lichterfeld M. T memory stem cells and HIV: A long-term relationship. *Curr HIV/AIDS Rep* 2015;12(1):33–40.
- Hamann D, Baars PA, Rep MH, Hooibrink B, Kerkhof-Garde SR, et al. Phenotypic and functional separation of memory and effector human CD8+ T cells. *J Exp Med* 1997;186(9):1407–1418.
- Mpande CAM, Dintwe OB, Musvosvi M, Mabwe S, Bilek N, et al. Functional, antigen-specific stem cell memory (TSCM) CD4(+) T cells are induced by human *Mycobacterium tuberculosis* infection. *Front Immunol* 2018;9:324.
- Zanon V, Pilipow K, Scamardella E, De Paoli F, De Simone G, et al. Curtailed T-cell activation curbs effector differentiation and generates CD8(+) T cells with a naturally-occurring memory stem cell phenotype. *Eur J Immunol* 2017;47(9):1468–1476.
- Booiman T, Wit FW, Girigorie AF, Maurer I, De Francesco D, et al. Terminal differentiation of T cells is strongly associated with CMV infection and increased in HIV-positive individuals on ART and lifestyle matched controls. *PLoS One* 2017;12(8):e0183357.
- Golubovskaya V, Wu L. Different subsets of T cells, memory, effector functions, and CAR-T immunotherapy. *Cancers (Basel)* 2016;8:3.
- Michie CA, McLean A, Alcock C, Beverley PC. Lifespan of human lymphocyte subsets defined by CD45 isoforms. *Nature* 1992;360(6401):264–265.
- Wang X, He Q, Shen H, Lu XJ, Sun B. Genetic and phenotypic difference in CD8(+) T cell exhaustion between chronic hepatitis B infection and hepatocellular carcinoma. *J Med Genet* 2018;56(1):18–21.
- Ammann S, Lehmerberg K, Zur Stadt U, Janka G, Rensing-Ehl A, et al. Primary and secondary hemophagocytic lymphohistiocytosis have different patterns of T-cell activation, differentiation and repertoire. *Eur J Immunol* 2017;47(2):364–373.
- Groom JR, Richmond J, Murooka TT, Sorensen EW, Sung JH, Bankert K, von Andrian UH, Moon JJ, Mempel TR, Luster AD. CXCR3 chemokine receptor-ligand

- interactions in the lymph node optimize CD4+ T helper 1 cell differentiation. *Immunity* 2012;37(6):1091–1103.
58. Takeshita M, Suzuki K, Kassai Y, Takiguchi M, Nakayama Y, Otomo Y, Morita R, Miyazaki T, Yoshimura A, Takeuchi T. Polarization diversity of human CD4+ stem cell memory T cells. *Clin Immunol* 2015;159(1):107–117.
  59. Delpoux A, Michelini RH, Verma S, Lai CY, Omilusik KD, Utzschneider DT, Redwood AJ, Goldrath AW, Benedict CA, Hedrick SM. Continuous activity of Foxo1 is required to prevent anergy and maintain the memory state of CD8(+) T cells. *J Exp Med* 2018;215(2):575–594.
  60. Takata H, Takiguchi M. Three memory subsets of human CD8+ T cells differently expressing three cytolytic effector molecules. *J Immunol* 2006;177(7):4330–4340.
  61. Bengsch B, Ohtani T, Herati RS, Bovenschen N, Chang KM, Wherry EJ. Deep immune profiling by mass cytometry links human T and NK cell differentiation and cytotoxic molecule expression patterns. *J Immunol Methods* 2018;453:3–10.
  62. Lin L, Couturier J, Yu X, Medina MA, Kozinetz CA, Lewis DE. Granzyme B secretion by human memory CD4 T cells is less strictly regulated compared to memory CD8 T cells. *BMC Immunol* 2014;15:36.