

The force-free magnetosphere of a Kerr black hole: the role of boundary conditions

Zhen Pan^{1,*}

¹*Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada*

(Dated: June 25, 2018)

The role of boundary conditions in the structure of a force-free black hole magnetosphere was rarely discussed, since previous studies have been focused on the the field lines entering the horizon which is causally disconnected and on which the boundary condition imposed usually makes no difference to the magnetosphere structure. However, recent high-resolution general relativistic (GR) force-free simulation shows that there are both field lines entering the horizon and field lines ending up on the equatorial current sheet within the ergosphere for asymptotic uniform field. For the latter, the equatorial boundary condition is well approximated being marginally force-free, i.e., $B^2 - E^2 = 0$, where B and E are the magnetic and electric field strength, respectively. In this paper, we revisit this problem by solving the GR Grad-Shafranov equation which governs the structure of the force-free magnetosphere in steady state and self-consistently imposing the marginally force-free equatorial condition. We also discuss the applicability of this boundary condition and the numerical algorithm proposed in this paper for general magnetic field configurations.

I. INTRODUCTION

The paper is organized as follows. In Section II, we outline the basic governing equations. In Section III, we clarify the radiation condition, boundary conditions and numerical algorithm for the uniform field solution. In Section IV, we discuss the solution uniqueness, near-horizon field line configuration and the application to general field line configurations. Summary is given in Section V.

II. BASIC EQUATIONS

In this paper, we adopt the Kerr-Schild coordinate with the line element

$$ds^2 = - \left(1 - \frac{2r}{\Sigma}\right) dt^2 + \left(\frac{4r}{\Sigma}\right) dr dt + \left(1 + \frac{2r}{\Sigma}\right) dr^2 \\ + \Sigma d\theta^2 - \frac{4ar \sin^2 \theta}{\Sigma} d\phi dt - 2a \left(1 + \frac{2r}{\Sigma}\right) \sin^2 \theta d\phi dr \\ + \frac{\beta}{\Sigma} \sin^2 \theta d\phi^2$$

where $\Sigma = r^2 + a^2 \mu^2$, $\Delta = r^2 - 2r + a^2$, $\beta = \Delta \Sigma + 2r(r^2 + a^2)$, and a is the dimensionless BH spin. In the force-free approximation, electromagnetic energy greatly exceeds that of matter. Consequently, the force-free magnetosphere is governed by energy conservation equation of electromagnetic field, or conventionally called as the GS equation. In the Kerr spacetime, the axisymmetric and

steady GS equation can be written in a compact form [1]

$$\begin{aligned} & \left[A_{\phi,rr} + \frac{\sin^2 \theta}{\Delta} A_{\phi,\mu\mu} \right] \mathcal{K}(r, \theta; \Omega) \\ & + \left[A_{\phi,r} \partial_r^\Omega + \frac{\sin^2 \theta}{\Delta} A_{\phi,\mu} \partial_\mu^\Omega \right] \mathcal{K}(r, \theta; \Omega) \\ & + \frac{1}{2} \left[A_{\phi,r}^2 + \frac{\sin^2 \theta}{\Delta} A_{\phi,\mu}^2 \right] \Omega' \partial_\Omega \mathcal{K}(r, \theta; \Omega) \\ & - \frac{\Sigma}{\Delta} II' = 0, \end{aligned} \quad (1)$$

with $\mathcal{K}(r, \theta; \Omega) = \left[\frac{\beta}{\Sigma} \Omega^2 \sin^2 \theta - \frac{4ra}{\Sigma} \Omega \sin^2 \theta - \left(1 - \frac{2r}{\Sigma}\right) \right]$ being the LS function, $\mu \equiv \cos \theta$, and the primes designate derivatives with respect to A_ϕ . $\partial_i^\Omega (i = r, \mu)$ denotes the partial derivative with respect to coordinate i with Ω fixed, and ∂_Ω is the derivative with respect to Ω .

III. UNIFORM FIELD SOLUTION

A. Solution uniqueness and radiation condition

For common field configurations, including split monopole and parabola, there exist two LSs where the LS function vanishes and the GS equation degrades from second order to first order. As proposed by Contopoulos et.al. [2], one can adjust the two free functions $\Omega(A_\phi)$ and $I(A_\phi)$ enabling field lines smoothly cross the two LSs, then the solution $\{\Omega(A_\phi), I(A_\phi), A_\phi(r, \theta)\}$ is uniquely found. But for the vertical field lines, their exist only one LS, which is insufficient to determine two free functions. Therefore people have been wondering the possibility of many solutions though the many-solutions scenario is not supported by force-free dynamics simulations. To tackle this problem, we proposed that the two functions are not independent; instead, they are related by the radiation condition [1, 3]

$$I = 2\Omega A_\phi, \quad (2)$$

* zpan@perimeterinstitute.ca

which has been confirmed by the recent high-accuracy simulations [4].

B. Boundary conditions

The boundary conditions at infinity (inner infinity $r = r_+$ and outer infinity $r \rightarrow \infty$) and on the polar axis can be simply set as

$$\begin{aligned} A_{\phi,r}|_{r=r_+, \infty} &= 0, \\ A_{\phi}|_{\mu=1} &= 0, \end{aligned} \quad (3)$$

where r_+ is the radius of the event horizon, while the equatorial boundary conditions are more unclear until recent simulations [4] come out which show that there exists an equator current sheet within the ergosphere where the magnetic dominance marginally loses, i.e., $B^2 - E^2 \approx 0$. Motivated by these simulations, we choose the following equatorial boundary condition in our numerical solutions,

$$\begin{aligned} A_{\phi,\mu}(\mu = 0, r > 2) &= 0, \\ B^2 - E^2(\mu = 0, r_+ \leq r \leq 2) &= 0. \end{aligned} \quad (4a) \quad (4b)$$

In fact, Equation (4b) is neither a Dirichlet nor a Neumann boundary condition, since

$$\begin{aligned} B^2 - E^2 \\ = \frac{1}{\Sigma \sin^2 \theta} \left[-\mathcal{K} \left(A_{\phi,r}^2 + \frac{\sin^2 \theta}{\Delta} A_{\phi,\mu}^2 \right) + \frac{\Sigma}{\Delta} I^2 \right], \end{aligned} \quad (5)$$

which involves both derivatives $A_{\phi,\mu}$ and $A_{\phi,r}$ on the boundary. Therefore, it is numerically non-trivial to impose this boundary condition in computation.

In addition, there is a coordinate singularity $1/\Delta$ in the expression of $B^2 - E^2$. To avoid possible numerical difficulty, we use following prescription

$$\int_{A_{\phi}^{\text{HE}}}^{A_{\phi}^{\text{EE}}} \left(\frac{B^2 - E^2}{B^2 + E^2} \right)^2 dA_{\phi} \Big/ (A_{\phi}^{\text{EE}} - A_{\phi}^{\text{HE}}) < 10^{-3}. \quad (6)$$

in our computation, as a proxy of the marginally force-free equatorial boundary condition (4b), where A_{ϕ}^{HE} and A_{ϕ}^{EE} are the magnetic flux entering the horizon and the ergosphere, respectively; “HE” and “EE” are short for Horizon-Equator and Ergosphere-Equator, respectively. For definiteness, we choose $B^2 + E^2$ to be the energy density measured by the zero-angular-momentum-observers. Explicitly, we have

$$\begin{aligned} B^2 + E^2 \\ = \frac{1}{\Sigma \sin^2 \theta} \left[\left(\mathcal{K} + \frac{\Delta \Sigma}{\beta} \right) \left(A_{\phi,r}^2 + \frac{\sin^2 \theta}{\Delta} A_{\phi,\mu}^2 \right) + \frac{\Sigma}{\Delta} I^2 \right] \end{aligned} \quad (7)$$

C. Generic features of the BH magnetosphere structure

Before delving into the details of numerically solving the GS equations, we point out that from the radiation condition (2) and the marginally force-free boundary condition (4b) themselves contains rich information about the BH magnetosphere structure.

Let's first find out where the LS intersect with the equator, $r_{\text{LS}}|_{\mu=0}$. From the expression of the LS function, it is straightforward to see $r_{\text{LS}}|_{\mu=0} \leq 2$ for $\Omega \geq 0$. At this point, the LS function \mathcal{K} vanishes, and I must also vanish for satisfying the boundary condition (4b), which in turns indicates a vanishing angular velocity Ω from the radiation condition (2). Due to the static limit, Ω must be non-zero inside the ergosphere. Therefore the only possibility is $r_{\text{LS}}|_{\mu=0} = 2$, i.e. to satisfy the boundary condition (4b), the LS must interact the equator at $r = 2$, which also justifies our choice of equatorial boundary conditions (4a, 4b).

From above analysis, we expect several generic features in the magnetospheres that are independent of the GS equation: (1) the LS runs from $r = r_+$ to $r = 2$ as θ varies from 0 to $\pi/2$, (2) there is no current sheet within the magnetosphere except the equatorial current sheet extending from r_+ to 2, which gives rise to a cusp ($A_{\phi,\mu} \neq 0$) to the equatorial magnetic field lines, (3) magnetic field lines entering the ergosphere end up either on the horizon or on the equatorial current sheet, both of which carry electric current and therefore Poynting flux. With the guidance of the qualitative features above, we are to numerically solve the GS equation and quantify these features.

D. Numerical method

We aim to find a pair of $\Omega(A_{\phi})$ and $I(A_{\phi})$ satisfying the radiation condition (2) and enabling field lines smoothly crossing the LS, and suitable normal derivative $A_{\phi,\mu}(\mu = 0, r_+ \leq r \leq 2)$ on the equator guaranteeing the boundary condition (4b), which itself cannot be directly used in computation.

In our computation, we define a new radial coordinate $R = r/(1+r)$, confine our computation domain $R \times \mu$ in the region $[R(r_+), 1] \times [0, 1]$, and implement a uniform 512×64 grid. The numerical algorithm for searching desired $\{\Omega(A_{\phi}), I(A_{\phi}), A_{\phi,\mu}(\mu = 0, r_+ \leq r \leq 2)\}$ is detailed in the following steps.

1. We choose an initial guess for the field configuration, functions $\{\Omega(A_{\phi}), I(A_{\phi})\}$ and equatorial boundary

condition as follows

$$A_\phi = \frac{B_0}{2} r^2 \sin^2 \theta, \quad (8a)$$

$$\Omega = 0.5\Omega_H (1 - A_\phi/A_\phi^{\text{HE}}), \quad (8b)$$

$$I = \Omega_H A_\phi (1 - A_\phi/A_\phi^{\text{HE}}), \quad (8c)$$

$$A_{\phi,\mu}(\mu = 0, r_+ \leq r \leq 2) = -(r/r_+)^3, \quad (8d)$$

where $\Omega_H = a/2r_+$ is the angular velocity of the BH.

2. We evolve the GS equation (1) using the well-known relaxation method [5] and adjust $II'(A_\phi)$ until field lines smoothly cross the LS [see e.g. 2, 3, 6, 7, for more details].

3. Usually the current I found in Step 2 neither satisfies the radiation condition (2) nor guarantees the boundary condition (4b). We adjust $A_{\phi,\mu}(\mu = 0, r_+ \leq r \leq 2)$ as follows

$$A_{\phi,\mu}|_{\text{new}} = A_{\phi,\mu}|_{\text{old}} + \zeta_1 \times [2\Omega A_\phi (2\Omega A_\phi)' - II'], \quad (9)$$

where ζ_1 is an empirical step size. For each new $A_{\phi,\mu}$, we repeat Step 2 until $A_{\phi,\mu}(\mu = 0, r_+ \leq r \leq 2)$ converges, i.e. the condition $2\Omega A_\phi (2\Omega A_\phi)' = II'$ is achieved for $A_\phi \in (A_\phi^{\text{HE}}, A_\phi^{\text{EE}})$.

4. The remaining task is to adjust $\Omega(0 < A_\phi < A_\phi^{\text{HE}})$ enabling the radiation condition (2) for $A_\phi \in (0, A_\phi^{\text{HE}})$ and to adjust $\Omega(A_\phi^{\text{HE}} \leq A_\phi \leq A_\phi^{\text{EE}})$ enabling the boundary condition (4b) for $A_\phi \in [A_\phi^{\text{HE}}, A_\phi^{\text{EE}}]$. The first part is straightforward, i.e.,

$$2A_\phi \Omega_{\text{new}} = I|_{0 < A_\phi < A_\phi^{\text{HE}}}, \quad (10)$$

and the second part can be realized by iterative corrections

$$2A_\phi (\Omega_{\text{new}} - \Omega_{\text{old}}) = -\zeta_2 \times \Delta(B^2 - E^2)|_{\mu=0, r_+ \leq r \leq 2}, \quad (11)$$

where ζ_2 is again an empirical step size, and we have multiplied factor Δ in the correction term to avoid numerical difficulty in the vicinity of the event horizon. To eliminate unphysical discontinuity in the angular velocity at A_ϕ^{HE} , we fit $\Omega_{\text{new}}(A_\phi)$ on the whole range $(0, A_\phi^{\text{EE}})$ via a fifth-order polynomial.

5. For the new angular velocity $\Omega_{\text{new}}(A_\phi)$ obtained in Step 4, we repeat Step 2 to Step 4, until the numerical prescription (6) for the boundary condition (4b) is satisfied.

E. Numerical results

In Figure 1, we plot the magnetic field lines enclosing a BH with spin $a = 0.99$ as an example, and in the upper panel of Figure 2, we show the angular velocity functions $\Omega(A_\phi)$ for different BH spins.

With the angular velocity $\Omega(A_\phi)$ available, the energy extraction rate from the BH is known as

$$\dot{E} = 4\pi \int_0^{A_\phi^{\text{EE}}} \Omega \times I dA_\phi. \quad (12)$$




FIG. 1. The configuration of field lines for the magnetosphere of a Kerr BH with spin $a = 0.99$, where the solid/red line is the ergosphere and the dashed/black line is the LS, both of which intersect with the equator at $r = 2M$.




FIG. 2. Upper panel: the angular velocity functions $\Omega(A_\phi)$ for different BH spins. Lower Panel: comparison of our numerical results (solid lines) with the simulation results in [4] (dashed lines).




FIG. 3. Comparison of the energy extraction rate $\dot{E}(\Omega_H)$ obtained from our numerical solutions and from the high-resolution force-free simulations [4], where the former is lower than the latter by $\sim 10\%$ for large spins.

In the limit of slow rotation, analytic solution is known as [4, 8],

$$\Omega = \Omega_H \frac{\sqrt{1-\psi}}{1+\sqrt{1-\psi}}, \quad (13)$$

where $\psi = A_\phi/(2B_0 M^2)$, and the energy extraction rate is

$$\dot{E} = 128\pi \left(\frac{17}{24} - \ln 2 \right) B_0^2 M^4 \Omega_H^2. \quad (14)$$

In the lower panel of Figure 2 and in Figure 3, we compare our numerical results with recent high-resolution simulations done by East and Yang [4]. In Figure 3, the points are taken from either the simulations or our numerical solutions, while the solid lines are corresponding polynomial fitting curves which require to approach

Eq.(14) for small spins and to be flat for extremal spins. We find the angular velocity functions obtained from two different approaches yield a good agreement (Figure 2), while the energy extraction rate \dot{E} is $\sim 10\%$ lower than the corresponding simulation results, which is traced to $\sim 5\%$ smaller magnetic flux A_ϕ^{EE} in our numerical solutions.

IV. DISCUSSIONS

A. General application

In real astrophysics environment, we expect the field configurations are more close to parabolas instead of being strictly vertical. Due to the same topology, it is reasonable to expect the marginally force-free boundary condition (4b) is also a good work approximation. The features we observed the uniform field solution should also apply here. The numerical method for solving the GS equation and self-consistently imposing the boundary condition (4b).

B. Near-horizon field lines

In one of our previous papers [3], we made a claim that “in the steady axisymmetric force-free magnetosphere around a Kerr BH, all magnetic field lines that cross the infinite-redshift surface must intersect the event horizon” based on two. However, simulations shows that there exists an equatorial current sheet within the ergosphere, where the force-free condition breaks down.

V. SUMMARY

In this paper, we revisit the problem of the uniform field solution to the BZ mechanism.

-
- [1] Z. Pan, C. Yu, and L. Huang, *Astrophys. J.* **836**, 193 (2017).
 - [2] I. Contopoulos, D. Kazanas, and D. Papadopoulos, *Astrophys. J.* **765**, 113 (2013), [arXiv:1212.0320 \[astro-ph.HE\]](#).
 - [3] Z. Pan and C. Yu, *Astrophys. J.* **816**, 77 (2016).
 - [4] W. E. East and H. Yang, (2018), [arXiv:1805.05952](#).
 - [5] W. Press, S. Teukolsky, W. Vetterling, B. Flannery, E. Ziegel, W. Press, B. Flannery, S. Teukolsky, and W. Vetterling, *Technometrics* (Cambridge University Press, 1987).
 - [6] A. Nathanail and I. Contopoulos, *Astrophys. J.* **788**, 186 (2014), [arXiv:1404.0549 \[astro-ph.HE\]](#).
 - [7] J. F. Mahlmann, P. Cerdá-Durán, and M. A. Aloy, *Mon. Not. R. Astron. Soc.* **477**, 3927 (2018), [arXiv:1802.00815](#).
 - [8] Z. Pan and C. Yu, (2014), [arXiv:1406.4936](#).