

On the Subcritical Self-Catalytic Branching Brownian Motions (SBBM)

Zhenyao Sun

Beijing institute of Technology (BIT)

Sept. 2025

University POLITEHNICA Bucharest

Based on joint work with Clayton Barnes (AWS) and Leonid Mytnik (Technion),
and joint work with Haojie Hou (BIT)

- **Model: Self-Catalytic Branching Brownian Motions (SBBM)**

A particle system that extends the **Branching Brownian Motion** by incorporating **catalytic branching** through pairwise interactions between particles (which will be defined in the next slide).

- **Objectives:**

- Construct an SBBM that supports **infinitely many particles**.
- Characterize the law of the SBBM.
- Investigate the **coming down from infinity (CDI)** property of the SBBM.

- **Motivation:**

- SBBMs offer insights into complex population dynamics, incorporating biological dispersal and intraspecific competition/cooperation.
- SBBMs serve as **moment duals** to a class of stochastic reaction-diffusion equations with multiplicative noise.

SBBM Dynamics:

- **Initial Configuration:**

The positions of the initial particles are given by $(x_i)_{i=1}^n \subset \mathbb{R}$.

- **Particle Movement:**

Each particle performs independent Brownian motion on \mathbb{R} .


- **Ordinary Branching:**

Particles branch independently at rate β_o , replaced by k (random) offspring according to the law (p_k) .

- **Catalytic Branching:**

Each pair of particles independently branches at rate β_c , based on their intersection local times, replaced by k (random) offspring according to the law (q_k) .

An illustration of SBBM



- A graph illustration of SBBM with $p_3 = 1$ and $q_1 = 1$.

Assumptions

- **Catalytic Branching is Subcritical:**

$$\sum kq_k < 2.$$

- **Catalytic Branching is Not Parity-Preserving:**

There exists an odd k such that $q_k > 0$.

- **A Technical Assumption:**

There exists $R > 1$ such that

$$\sum R^k p_k < \infty \quad \text{and} \quad \sum R^k q_k < \infty.$$

The Explosion Problem for SBBM

A Priori Consideration:

- The SBBM model is well-defined only up to its explosion time τ_∞ .

Definition of the Explosion Time τ_∞ :

- Define a sequence of stopping times $\{\tau_k\}_{k \in \mathbb{Z}_+}$ such that:
 - $\tau_0 = 0$.
 - For each $k \geq 0$,

$$\tau_{k+1} := \inf\{t > \tau_k : \text{a branching occurs at time } t\}.$$

- The explosion time is defined as:

$$\tau_\infty := \lim_{k \rightarrow \infty} \tau_k.$$

The Explosion Problem:

- Does $\tau_\infty = \infty$?

Non-Explosion Property

Athreya-Tribe (2000, Ann. Probab.)

If the ordinary branching is subcritical, i.e., $\sum kp_k < 1$, then:

$$\mathbb{P}(\tau_\infty = \infty) = 1.$$

Hou-S. (2025, arXiv)

Same result holds without assuming $\sum kp_k < 1$.

Implications:

- The SBBM model remains well-defined for all time.
- For $t \geq 0$, define $Z_t^{(n)}(A) := \#\{\text{particles in set } A \text{ at time } t\}$.
- The process $(Z_t^{(n)})_{t \geq 0}$ is a Markov process taking values in:

$$\mathcal{N} := \{\text{locally finite point measures on } \mathbb{R}\}.$$

- We refer to $Z_t^{(n)}$ as an SBBM (with n initial particles).

The Infinitely Initial Particles Problem

- Consider the scenario where the initial configuration of an SBBM consists of infinitely many particles located at $(x_i)_{i=1}^{\infty} \subset \mathbb{R}$.
- In this case, $\tau_{\infty} = 0$ almost surely, and the model is a priori not well-defined.

The Infinitely Initial Particles Problem:

- Let $(Z_t^{(n)})_{t \geq 0}$ be an SBBM with initial value $\sum_{i=1}^n \delta_{x_i}$.
- What is the limit of the processes $(Z_t^{(n)})_{t \geq 0}$ as $n \rightarrow \infty$?

Characterizing the Law:

- How can we characterize the law of $(Z_t^{(n)})_{t \geq 0}$?
- Is the family of processes $\{(Z_t^{(n)})_{t \geq 0} : n \in \mathbb{N}\}$ tight?

The Dual SPDE

- **The dual SPDE of SBBM:**

$$\begin{cases} \partial_t u_t(x) = \frac{\Delta}{2} u_t(x) - \Phi(u_t(x)) + \sqrt{\Psi(u_t(x))} \dot{W}_t(x), & t > 0, x \in \mathbb{R}, \\ u_0(x) = f(x), & x \in \mathbb{R}. \end{cases}$$

- **Space-time White Noise:**

$(W_t)_{t \geq 0} :=$ a cylindrical Wiener process on $L^2(\mathbb{R})$, such that

$$\mathbb{E}[W_t(\phi)W_s(\psi)] = (t \wedge s)\langle \phi, \psi \rangle.$$

- **Ordinary Branching Mechanism:**

$$\Phi(z) := \beta_o \left(\sum_{k=0}^{\infty} p_k (1-z)^k - (1-z) \right).$$

- **Catalytic Branching Mechanism:**

$$\Psi(z) := \beta_c \left(\sum_{k=0}^{\infty} q_k (1-z)^k - (1-z)^2 \right).$$

The Dual SPDE

- Let $z^* := \inf\{z \in [1, 2] : \Psi(z) = 0\}$.
- Under the assumption that the catalytic branching is not parity-preserving, we have $z^* \in [1, 2)$.
- Let $C(\mathbb{R}, [0, z^*]) := \{\text{Continuous functions from } \mathbb{R} \text{ to } [0, z^*]\}$.
- We say a $C(\mathbb{R}, [0, z^*])$ -valued continuous process $(u_t)_{t \geq 0}$ is a weak solution, if \exists a space-time white noise \dot{W} such that $\forall \phi \in C_c^\infty(\mathbb{R})$,

$$\begin{aligned} & \langle u_t, \phi \rangle - \langle f, \phi \rangle \\ &= \int_0^t \langle u_s, \frac{\Delta}{2} \phi \rangle ds - \int_0^t \langle \Phi(u_s), \phi \rangle ds + \int_0^t \langle \sqrt{\Psi(u_s)} \phi, dW_s \rangle, \quad \text{a.s.} \end{aligned}$$

Shiga (1994, Can. J. Math.)

For each initial value $f \in C(\mathbb{R}, [0, z^*])$, there exists a weak solution to the dual SPDE.

The Duality

- For any $[0, z^*]$ -valued function g and point measure μ , define

$$(1 - g)^\mu := \prod_{x \in \mathbb{R}} (1 - g(x))^{\mu(\{x\})}.$$

Athreya-Tribe (2000, Ann. Probab.)

If the ordinary branching is subcritical, i.e., $\sum kp_k < 1$, then:

$$\mathbb{E} \left[(1 - u_t)^{Z_0^{(n)}} \right] = \mathbb{E} \left[(1 - u_0)^{Z_t^{(n)}} \right].$$

Hou-S. (2025, arXiv)

The result above holds without assuming $\sum kp_k < 1$.

- Corollary:** The uniqueness in law holds for the dual SPDE.

SBBM with Infinitely Many Initial Particles

- Let the state space \mathcal{N} be equipped with the vague topology.

Initial Trace (Λ, μ) :

- $\Lambda := \{\text{sub-sequential limits of } (x_i)_{i=1}^\infty\}$.
- $\mu := \sum_{x_i \notin \Lambda} \delta_{x_i}$.

Hou-S. (2025, arXiv)

There exists an \mathcal{N} -valued càdlàg Markov process $(Z_t)_{t \geq 0}$ such that $(Z_t^{(n)})_{t \geq 0}$ converges to $(Z_t)_{t \geq 0}$ as $n \rightarrow \infty$ in finite-dimensional distributions. The law of the process $(Z_t)_{t \geq 0}$ is determined by the two branching mechanisms (Φ, Ψ) and the initial trace (Λ, μ) .

- We call $(Z_t)_{t \geq 0}$ an SBBM with initial trace (Λ, μ) and branching mechanisms (Φ, Ψ) .

Coming Down from Infinity (CDI)

The Local-Time Coalescing Brownian Motions (LCBM):

- If the ordinary branching rate $\beta_o = 0$ and the catalytic branching law satisfies $q_1 = 1$, then the SBBM degenerates into the LCBM.
- In this case, $\Phi = 0$ and $\Psi(z) = z(1 - z)$.

Barnes-Mytnik-S. (2024, Ann. Probab.)

Suppose that $(Z_t)_{t>0}$ is an LCBM. Let U be any open interval. Then, almost surely, for every $t > 0$,

$$Z_t(U) < \infty \iff (\Lambda \cup \text{supp}(\mu)) \cap U \text{ is bounded.}$$

Hou-S. (2025, arXiv)

The same result holds for SBBM.

- **The CDI property:** Almost surely, for every $t > 0$,

$$Z_t(\mathbb{R}) < \infty \iff \sup\{|x_i| : i \in \mathbb{N}\} < \infty.$$

The Mean Field Equation (MFE)

From a physic's point of view:

- The MFE for a system of independent Brownian motions is given by the heat equation $\partial_t h = \frac{\Delta}{2} h$. In the sense that

$$\mathbb{E}[\#\{\text{particles in } (x - \frac{1}{2}, x + \frac{1}{2}) \text{ at time } t\}] \approx h_t(x).$$

- The MFE for LCBM is $\partial_t v = \frac{\Delta}{2} v - \frac{\Psi'(0+)}{2} v^2$.

Le Gall (1996, J. Appl. Math. Stochastic Anal.)

There exists a unique non-negative solution $(v_t(x))_{t>0, x \in \mathbb{R}}$ to the PDE

$$\begin{cases} \partial_t v_t(x) = \frac{\Delta}{2} v_t(x) - \frac{\Psi'(0+)}{2} v_t(x)^2, & t > 0, x \in \mathbb{R}, \\ \left\{ y \in \mathbb{R} : \forall r > 0, \lim_{t \rightarrow 0} \int_{y-r}^{y+r} v_t(x) dx = \infty \right\} = \Lambda, \\ \lim_{t \rightarrow 0} \langle v_t, \phi \rangle = \langle \mu, \phi \rangle, \quad \phi \in C_c(\Lambda^c). \end{cases}$$

Speed of CDI for LCBM

The Speed of CDI Problem

- Assume CDI holds for a process $(N_t)_{t \geq 0}$.
- Can we find a rate function $a(t)$ such that $N_t/a(t) \rightarrow 1$ as $t \downarrow 0$?

Barnes-Mytnik-S. (2024, Ann. Probab.)

Suppose that $(Z_t)_{t > 0}$ is an LCBM with initial trace (Λ, μ) . Let U be an open interval. Suppose that $(\Lambda \cap \text{supp}(\mu)) \cap U$ is bounded and $\Lambda \cap \bar{U} \neq \emptyset$. Then,

$$\left(\int_U v_t(x) dx \right)^{-1} Z_t(U) \xrightarrow[t \downarrow 0]{L^1} 1,$$

where $(v_t(x))_{t > 0, x \in \mathbb{R}}$ is the solution to the corresponding MFE with initial trace (Λ, μ) .

Hou-S. (2025, arXiv)

The same result holds for SBBM.

Criticality of the Branching:

- It is crucial for our result that the catalytic branching is subcritical, i.e., $\sum kq_k < 2$.
- Barnes-Mytnik-S. (2025, Probab. Theory Related Fields) constructed an SBBM with $p_\infty = 1$ and $q_1 = 1$, and showed that the total population in this model is “reflecting from infinity”.
- When the catalytic branching is supercritical, i.e., $\sum kq_k > 2$, we believe that the SBBM will explode in finite time.
- When there is no ordinary branching, i.e., $\beta_o = 0$, and the catalytic branching is critical, i.e., $\sum kq_k = 2$, we believe that the SBBM is non-explosive and rescales to the stochastic heat equation:

$$\partial_t u = \frac{\Delta}{2} u + u \dot{W}.$$

About the Parity:

- It is crucial for our result that the catalytic branching is not parity-preserving, i.e., there exists an odd number k such that $q_k > 0$.
- Consider an SBBM with no ordinary branching, i.e., $\beta_0 = 0$, and $q_0 = 1$. We call this model the local-time annihilating Brownian motion (LABM).
- LABM is non-explosive and can be defined up to all time, provided there are only finitely many initial particles.
- It can be shown that $\{Z_t^{(n)} : n \in \mathbb{N}\}$ is tight.
- However, the subsequential convergence-in-distribution limit of $\{Z_t^{(n)} : n \in \mathbb{N}\}$ is not unique.
- [Hammer-Ortgiese-Völlering \(2021, Stochastic Process. Appl.\)](#): The entrance laws of the (hard) annihilating Brownian motion are characterized.
- Characterization of all entrance laws of LABM is still open.

Examples of Duality:

- We say two Markov processes $(X_t)_{t \geq 0}$ and $(Y_t)_{t \geq 0}$ are dual to each other if there exists a large class of functions $H(x, y)$ such that

$$\mathbb{E}[H(X_t, Y_0)] = \mathbb{E}[H(X_0, Y_t)].$$

- [Bachelier \(1900, Ann. Sci. École Norm. Sup.\)](#):
Brownian motion and the heat equation $\partial_t h = \frac{\Delta}{2} h$.
- [McKean \(1975, Comm. Pure Appl. Math.\)](#): Branching Brownian motion and the FKPP equation $\partial_t v = \frac{\Delta}{2} v + v(1 - v)$.
- [Harris \(1978, Ann. Probab.\)](#):
Coalescing random walk and the voter model.
- [Shiga \(1986, Math. Appl.\)](#): LCBM and the stochastic FKPP equation $\partial_t v = \frac{\Delta}{2} v + \sqrt{v(1 - v)} \dot{W}$.
- [Tóth-Werner \(1998, Probab. Theory Relat. Fields\)](#):
(Hard) Coalescing Brownian motions and itself.
- **Folklore**: Stochastic heat equation $\partial_t u = \frac{\Delta}{2} u + u \dot{W}$ and itself.
- ...


Coming Down from Infinity (CDI):

- Feller (1954, Trans. Amer. Math. Soc.): Some diffusion processes in one dimension.
- Aldous (1999, Bernoulli): Kingman's coalescent.
- Schweinsberg (2000, Electron. Comm. Probab.) and Berestycki-Berestycki-Limic (2010, Ann. Probab.): Λ -coalescent.
- Limic-Sturm (2006, Electron. J. Probab.) and Angel-Berestycki-Limic (2012, Probab. Theory Related Fields): Coalescing random walks on graphs.
- Mourrat-Weber (2017, Comm. Math. Phys.): Dynamical Φ_3^4 model (leading to a new construction of the Euclidean Φ_3^4 Field Theory).
- ...

The Mean Field Equation (MFE)

- The CDI rate of SBBM is characterized by $\partial_t v = \frac{\Delta}{2} v - \frac{\Psi'(0+)}{2} v^2$ despite that the true MFE is $\partial_t \tilde{v} = \frac{\Delta}{2} \tilde{v} + \Phi'(0+) \tilde{v} - \frac{\Psi'(0+)}{2} \tilde{v}^2$.
- This is because $v(s, y) \asymp \tilde{v}(s, y)$ uniformly for $(s, y) \in [0, 1] \times \mathbb{R}$.
- The equation $\partial_t v = \frac{\Delta}{2} v - v|v|^\alpha$ with initial trace (Λ, μ) was studied by [Marcus-Véron \(1999, Comm. Partial Differential Equations\)](#) in the PDE literature.
- [Watanabe \(1968, J. Math. Kyoto Univ.\)](#):
The equation $\partial_t v = \frac{\Delta}{2} v - \frac{\Psi'(0+)}{2} v^2$ is the Laplace dual to the Super-Brownian motion $(X_t)_{t \geq 0}$.
- [Le Gall \(1996, J. Appl. Math. Stochastic Anal.\)](#) used the equation $\partial_t v = \frac{\Delta}{2} v - \frac{\Psi'(0+)}{2} v^2$ to study the Brownian snake, which is related to the super-Brownian motion through a Ray-Knight type theorem.

Theory Roadmap



Thanks!