


Entropic gateway · N = 4 · T_N = 20480 t.u.


n_steps = 819,200,000 · h = 2.5e-05 · B = N n_steps = 3.277e+09 walker-steps · complete seeds: ABF 32/32, ABF+FR 32/32

Walker traces of slots 0-3, seed 8100


(a) ABF · seed 8100 · slots 0-3 · first 40 t.u.




(b) ABF · seed 8100 · slots 0-3 · whole run (T_N = 20480 t.u.)



(c) ABF+FR · seed 8100 · slots 0-3 · first 40 t.u.



(d) ABF+FR · seed 8100 · slots 0-3 · whole run (T_N = 20480 t.u.)



- ABF (every walker its own lineage)
- ABF+FR lineage of ancestor 1 (85 trace points)
- ABF+FR lineage of ancestor 2 (29 trace points)
- ABF+FR lineage of ancestor 3 (9628 trace points)
- ABF+FR lineage of ancestor 0 (38 trace points)
- ABF+FR: all other lineages

Trace seed 8100 = the first seed of the config with every arm complete (reproducible choice). Dots = saved trace points; lines join consecutive points only on the dense trace grid ($\Delta t = 0.1$ t.u.) and never across a lineage change, a rebirth or a jump > 0.6 (walker reborn as a copy). FR colours: the 4 whole-run ancestor labels with the most trace points among the traced slots; the rest grey.

Small N (N = 4, i.e. ≤ 4): each seed contributes only 4 walkers to an instantaneous histogram, so instantaneous marginals are necessarily discrete ($TV_{\text{inst floor } E_N[TV]} = 0.796$).

Shaded band: gate region $|x| \leq 0.5$; dashed: well minima $x = \pm 1$.

eqb_plot_config/2 · config gateway_production.json · 2026-10-10 08:09 UTC