
**Gateway family expA/shift_lam16 · N = 128 · T_N = 640 t.u.**

n_steps = 25,600,000 · h = 2.5e-05 · B = N n_steps = 3.277e+09 walker-steps · complete seeds: ABF 32/32, ABF+FR 32/32

Mean-force snapshots (bin mean force) vs reference (top) and difference (bottom)




ABF (n = 32) · ABF+FR (n = 32) · interquartile range · reference · EM-consistent − analytic reference

Line = median across seeds; band = interquartile range (25th-75th percentile) across seeds. Metrics are the frozen SCIENTIFIC_PLAN section-4 quantities, unsmoothed.
Bin mean force Gamma = M / (C + 1) (step) vs the analytic F′ (curve); bottom: Gamma minus the bin average of F′_ref, whose RMS over the eval window is the floor-free companion e_F′_stat; e_F′² = e_F′_stat² + floor², floor = 0.03227 (the within-bin variation of F′_ref: a hard lower bound of e_F′, not visible in this row). Empty bins read 0 (estimator convention).
Dashed guides: region boundaries x = ±0.5 (left well / gate / right well); shaded |x| > 1.5: outside the scored eval window. Top-row y range = the reference over the scored window ± 35 % of its span (larger excursions are clipped there and show in the difference row).
No smoothing.

eqb_plot_config/2 · config shift_lam16.json · 2026-10-11 09:26 UTC