


Gateway family conditional_relaxation/lam1 · N = 2048 · T_N = 40 t.u.


n_steps = 1,600,000 · h = 2.5e-05 · B = N n_steps = 3.277e+09 walker-steps · complete seeds: ABF 32/32, ABF+FR 32/32

Walker traces of slots 0-31, seed 8100


(a) ABF · seed 8100 · slots 0-31 · first 4 t.u.




(b) ABF · seed 8100 · slots 0-31 · whole run (T_N = 40 t.u.)



(c) ABF+FR · seed 8100 · slots 0-31 · first 4 t.u.



(d) ABF+FR · seed 8100 · slots 0-31 · whole run (T_N = 40 t.u.)



—●— ABF (every walker its own lineage)

—●— ABF+FR lineage of ancestor 361 (1120 trace points)

—●— ABF+FR lineage of ancestor 1850 (531 trace points)

—●— ABF+FR lineage of ancestor 283 (469 trace points)

—●— ABF+FR lineage of ancestor 5 (401 trace points)

—●— ABF+FR: all other lineages

Trace seed 8100 = the first seed of the config with every arm complete (reproducible choice). Dots = saved trace points; lines join consecutive points only on the dense trace grid ($\Delta t = 0.1$ t.u.) and never across a lineage change, a rebirth or a jump > 0.6 (walker reborn as a copy). FR colours: the 4 whole-run ancestor labels with the most trace points among the traced slots; the rest grey.

Shaded band: gate region $|x| \leq 0.5$; dashed: well minima $x = \pm 1$.

eqb_plot_config/2 · config lam1.json · 2026-10-10 14:20 UTC