


Ethane/LTA 300 K · N = 1 · T_N = 61440 t.u.

n_steps = 307,200,000 · h = 0.0002 · B = N n_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16 (N = 1: ABF only, no FR arm)

Seed-averaged instantaneous CV density vs budget fraction u

(a) ABF (mean over 16 seeds of hist_inst / N)



Identical diverging colour scale for every panel (and every N): blue = below the uniform target, grey = at target, red = above; clipped at 4x. Displayed on 36 CV bins (5 production bins each). Columns = the 200 uniform budget saves $u = k/200$.

Each column pools only $N \times$ seeds = 16 walkers over 36 cells (one walker = $2.2 \times$ uniform), so the map is necessarily speckled; read it for where the population sits, not for cell-level values.

Small N (N = 1, i.e. ≤ 4): each seed contributes only 1 walker to an instantaneous histogram, so instantaneous marginals are necessarily discrete (TV_inst floor $E_N[\text{TV}] = 0.944$).

Dashed horizontal guides: $|\phi| = 0.791$ rad (window $|z| < 1.5$ Å) and $|\phi| = 2.109$ rad (cage $|z| > 4$ Å).

eqb_plot_config/2 · config lta_300K_production.json · 2026-10-10 07:12 UTC