


Ethane/LTA 300 K · N = 1 · T_N = 61440 t.u.

n_steps = 307,200,000 · h = 0.0002 · B = N n_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16 (N = 1: ABF only, no FR arm)

Configuration summary vs physical time



Line = median across seeds; band = interquartile range (25th-75th percentile) across seeds. Metrics are the frozen SCIENTIFIC_PLAN section-4 quantities, unsmoothed.

Each column pools only $N \times \text{seeds} = 16$ walkers over 36 cells (one walker = $2.2 \times$ uniform), so the map is necessarily speckled; read it for where the population sits, not for cell-level values.

(1)-(3): markers = persistent crossing of the seed-median curve; 'cens. a | b' = seeds with a censored tau per arm; shaded below the grey line = MC reference noise (RMS of the per-bin SE). (4) top F, bottom F' at $u = 0.1$ (left) and 1 (right), same alignment as e_F . (5) seed-mean instantaneous density / uniform, diverging scale clipped at 4.

Small N (N = 1, i.e. ≤ 4): each seed contributes only 1 walker to an instantaneous histogram, so instantaneous marginals are necessarily discrete (TV_inst floor $E_N[\text{TV}] = 0.944$).

eqb_plot_config/2 · config lta_300K_production.json · 2026-10-10 07:12 UTC