


# Ethane/LTA 300 K · N = 1 · T\_N = 61440 t.u.

n\_steps = 307,200,000 · h = 0.0002 · B = N n\_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16 (N = 1: ABF only, no FR arm)

Configuration summary vs budget fraction u (budget-normalised)



Line = median across seeds; band = interquartile range (25th-75th percentile) across seeds. Metrics are the frozen SCIENTIFIC\_PLAN section-4 quantities, unsmoothed.

Each column pools only  $N \times \text{seeds} = 16$  walkers over 36 cells (one walker =  $2.2 \times$  uniform), so the map is necessarily speckled; read it for where the population sits, not for cell-level values.

(1)-(3): markers = persistent crossing of the seed-median curve; 'cens. a | b' = seeds with a censored tau per arm; shaded below the grey line = MC reference noise (RMS of the per-bin SE). (4) top F, bottom F' at  $u = 0.1$  (left) and 1 (right), same alignment as  $e_F$ . (5) seed-mean instantaneous density / uniform, diverging scale clipped at 4.

Small N (N = 1, i.e.  $\leq 4$ ): each seed contributes only 1 walker to an instantaneous histogram, so instantaneous marginals are necessarily discrete (TV\_inst floor  $E_N[\text{TV}] = 0.944$ ).

eqb\_plot\_config/2 · config lta\_300K\_production.json · 2026-10-10 07:12 UTC