


Ethane/LTA 300 K · N = 16 · T_N = 3840 t.u.


n_steps = 19,200,000 · h = 0.0002 · B = N n_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16, ABF+FR 16/16

Walker traces of slots 0-15, seed 30000


(a) ABF · seed 30000 · slots 0-15 · first 60 t.u.




(b) ABF · seed 30000 · slots 0-15 · whole run (T_N = 3840 t.u.)



(c) ABF+FR · seed 30000 · slots 0-15 · first 60 t.u.



(d) ABF+FR · seed 30000 · slots 0-15 · whole run (T_N = 3840 t.u.)



- ABF (every walker its own lineage)
- ABF+FR lineage of ancestor 14 (3561 trace points)
- ABF+FR lineage of ancestor 0 (1584 trace points)
- ABF+FR lineage of ancestor 2 (10723 trace points)
- ABF+FR lineage of ancestor 15 (3420 trace points)
- ABF+FR: all other lineages

Trace seed 30000 = the first seed of the config with every arm complete (reproducible choice). Dots = saved trace points; lines join consecutive points only on the dense trace grid ($\Delta t = 0.05$ t.u.) and never across a lineage change or a jump > 0.75 rad (periodic wrap or slot overwritten by a copy). FR colours: the 4 whole-run ancestor labels with the most trace points among the traced slots; the rest grey.

Shaded: window $|\phi| < 0.791$ rad (darker) and cages $|\phi| > 2.109$ rad (lighter).

eqb_plot_config/2 · config lta_300K_production.json · 2026-10-10 07:10 UTC