


# Ethane/LTA 300 K · N = 2 · T\_N = 30720 t.u.

n\_steps = 153,600,000 · h = 0.0002 · B = N n\_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16, ABF+FR 16/16

Seed-averaged instantaneous CV density vs physical time



Identical diverging colour scale for every panel (and every N): blue = below the uniform target, grey = at target, red = above; clipped at 4x. Displayed on 36 CV bins (5 production bins each). Columns = every save (log time; budget grid + physical checkpoints).

Each column pools only  $N \times \text{seeds} = 32$  walkers over 36 cells (one walker =  $1.1 \times$  uniform), so the map is necessarily speckled; read it for where the population sits, not for cell-level values.

Small N (N = 2, i.e.  $\leq 4$ ): each seed contributes only 2 walkers to an instantaneous histogram, so instantaneous marginals are necessarily discrete (TV\_inst floor  $E_N[\text{TV}] = 0.892$ ).

Dashed horizontal guides:  $|\phi| = 0.791$  rad (window  $|z| < 1.5$  Å) and  $|\phi| = 2.109$  rad (cage  $|z| > 4$  Å).

eqb\_plot\_config/2 · config lta\_300K\_production.json · 2026-10-10 07:11 UTC