


Entropic gateway · N = 4 · T_N = 20480 t.u.

n_steps = 819,200,000 · h = 2.5e-05 · B = N n_steps = 3.277e+09 walker-steps · complete seeds: ABF 32/32, ABF+FR 32/32

Configuration summary vs physical time



Line = median across seeds; band = interquartile range (25th-75th percentile) across seeds. Metrics are the frozen SCIENTIFIC_PLAN section-4 quantities, unsmoothed.

Each column pools only $N \times \text{seeds} = 128$ walkers over 36 cells (one walker = $0.28 \times$ uniform), so the map is necessarily speckled; read it for where the population sits, not for cell-level values.

(1)-(3): markers = persistent crossing of the seed-median curve; 'cens. a | b' = seeds with a censored tau per arm; shaded below the grey line = EM bias at h (analytic vs EM-consistent reference); dash-dot in (2) = the zero-noise floor 0.0323 of $e_{F'}$ (hard lower bound). (4)

top F, bottom F' at $u = 0.1$ (left) and 1 (right), same alignment as e_F . (5) seed-mean instantaneous density / uniform, diverging scale clipped at 4.

Small N (N = 4, i.e. ≤ 4): each seed contributes only 4 walkers to an instantaneous histogram, so instantaneous marginals are necessarily discrete (TV_inst floor $E_N[\text{TV}] = 0.796$).

eqb_plot_config/2 · config_gateway_production.json · 2026-10-10 09:47 UTC