


# Entropic gateway · N = 8 · T\_N = 10240 t.u.

n\_steps = 409,600,000 · h = 2.5e-05 · B = N n\_steps = 3.277e+09 walker-steps · complete seeds: ABF 32/32, ABF+FR 32/32

Establishment and crossings vs budget fraction u



Line = median across seeds; band = interquartile range (25th-75th percentile) across seeds. Metrics are the frozen SCIENTIFIC\_PLAN section-4 quantities, unsmoothed.

Establishment (frozen): first save after which the instantaneous far-state fraction (right well  $x > 0.5$ ) stays  $\geq \frac{1}{2}$  of its uniform target 0.361; the cumulative-visitation version uses C\_all (far-state bins) against  $\frac{1}{2}$  of their bin share. True events never count a copy (clones inherit labels).

Null calibration at N = 8: an EXACTLY uniform population (far count  $\sim \text{Bin}(N, 0.361)$  per save) fails the instantaneous criterion at a save with  $p = 0.153$  and passes every save of the second half with  $P = 0.000 \rightarrow \text{UNRELIABLE}$  here: the instantaneous establishment time is censored / late by chance even when established, and FR birth-death can push the count below binomial scatter (an FR advantage in it is not evidence of faster convergence); read the cumulative version and the first arrival.

eqb\_plot\_config/2 · config gateway\_production.json · 2026-10-10 09:47 UTC