


Ethane/LTA 150 K · N = 256 · T_N = 240 t.u.


n_steps = 1,200,000 · h = 0.0002 · B = N n_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16, ABF+FR 16/16

Marginal convergence: TV_half and TV_inst with the finite-N floor vs physical time

(a) TV_half: visitation over the trailing half $C_{all}(t) - C_{all}(\approx t/2)$
censored tau: 0/16 | 0/16



(b) TV_inst: instantaneous walker histogram (descriptive)



- ABF (n = 16)
- ABF+FR (n = 16)
- interquartile range
- TV_half threshold 0.1 (marker: persistent crossing of the median curve)
- finite-N floor $E_N[TV] = 0.1029$ (exact, multinomial(N, uniform))

Line = median across seeds; band = interquartile range (25th-75th percentile) across seeds. Metrics are the frozen SCIENTIFIC_PLAN section-4 quantities, unsmoothed.

TV_half uses the saved step nearest $t/2$ (or the run start) as the window's lower edge (eqb_metrics.tv_half); tau for the marginal is defined on TV_half. TV_inst is never used for tau: at small N it is dominated by discreteness (17/18 at $N = 1$).

eqb_plot_config/2 · config lta_150K_production.json · 2026-10-10 09:46 UTC