


# Ethane/LTA 300 K · N = 1024 · T\_N = 60 t.u.

n\_steps = 300,000 · h = 0.0002 · B = N n\_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16, ABF+FR 16/16


Marginal convergence: TV\_half and TV\_inst with the finite-N floor vs physical time

(a) TV\_half: visitation over the trailing half  $C_{all}(t) - C_{all}(\approx t/2)$

censored tau: 0/16 | 0/16



(b) TV\_inst: instantaneous walker histogram (descriptive)



- ABF (n = 16)
- ABF+FR (n = 16)
- interquartile range
- O — TV\_half threshold 0.1 (marker: persistent crossing of the median curve)
- ..... finite-N floor  $E_N[TV] = 0.0514$  (exact, multinomial(N, uniform))

Line = median across seeds; band = interquartile range (25th-75th percentile) across seeds. Metrics are the frozen SCIENTIFIC\_PLAN section-4 quantities, unsmoothed.

TV\_half uses the saved step nearest  $t/2$  (or the run start) as the window's lower edge (eqb\_metrics.tv\_half); tau for the marginal is defined on TV\_half. TV\_inst is never used for tau: at small N it is dominated by discreteness (17/18 at  $N = 1$ ).

eqb\_plot\_config/2 · config lta\_300K\_production.json · 2026-10-10 09:46 UTC