





Ethane/LTA 150 K · N = 4 · T_N = 15360 t.u.

n_steps = 76,800,000 · h = 0.0002 · B = N n_steps = 3.072e+08 walker-steps · complete seeds: ABF 16/16, ABF+FR 16/16

Walker traces of slots 0-3, seed 15000



- ABF (every walker its own lineage)
- ABF+FR lineage of ancestor 1 (2018 trace points)
- ABF+FR lineage of ancestor 0 (415 trace points)
- ABF+FR lineage of ancestor 3 (7283 trace points)
- ABF+FR lineage of ancestor 2 (1208 trace points)
- ABF+FR: all other lineages

Trace seed 15000 = the first seed of the config with every arm complete (reproducible choice). Dots = saved trace points; lines join consecutive points only on the dense trace grid ($\Delta t = 0.05$ t.u.) and never across a lineage change or a jump > 0.75 rad (periodic wrap or slot overwritten by a copy). FR colours: the 4 whole-run ancestor labels with the most trace points among the traced slots; the rest grey.

Small N (N = 4, i.e. ≤ 4): each seed contributes only 4 walkers to an instantaneous histogram, so instantaneous marginals are necessarily discrete ($TV_{\text{inst}} \text{ floor } E_N[TV] = 0.796$).

Shaded: window $|\phi| < 0.791$ rad (darker) and cages $|\phi| > 2.109$ rad (lighter).

eqb_plot_config/2 · config lta_150K_production.json · 2026-10-10 07:19 UTC