

Manifold Fitting by Successive Tangent-Space Projection

Yuqing Xia¹, Bingjie Li², and Zhigang Yao^{3,*}

¹Zhejiang University of Finance and Economics, School of Data Sciences, Hangzhou, 310018, China

²Department of Epidemiology and Biostatistics, Ministry of Education Key Laboratory of Environment and Health, School of Public Health, Tongji Medical College, Huazhong University of Science and Technology, Wuhan, Hubei, China

³Department of Statistics & Data Science, National University of Singapore, 21 Lower Kent Ridge Road, Singapore

Abstract

Manifold fitting seeks to recover the geometric structure underlying noisy ambient observations. We propose Successive Tangent Space Projection (STSP), an iterative manifold-fitting method that reduces normal noise while preserving tangential variation. The former improves fitting accuracy, whereas the latter helps retain geometric coverage. We characterise the nearby population fixed-point set of STSP as a manifold-fitting object. Under uniform sampling from a compact smooth manifold with positive reach and isotropic Gaussian noise, this set lies within $O(\sigma^2)$ of the underlying manifold, and nearby population orbits converge geometrically to it. At the finite-sample level, with high probability, fixed points in the local tube lie within $O(\sigma^2)$ of the underlying manifold up to sampling error, and empirical orbits initialised within that tube and generated from a fixed reference sample enter and remain in the same neighbourhood. We further develop a multi-scale extension, MS-STSP, designed to reduce curvature bias while preserving the $O(\sigma^2)$ geometric localization order of STSP at both the population and finite-sample levels. Numerical experiments show that STSP compares favourably with competing methods in balancing fitting accuracy and geometric coverage, and that MS-STSP reduces curvature-induced shrinkage, particularly under high noise.

Keywords: Manifold fitting; Tangent-space projection; Geometric coverage; Multi-scale extrapolation; Ambient-space denoising.

1 Introduction

Manifold fitting aims to recover a smooth low-dimensional structure from noisy observations in a high-dimensional ambient space. It builds on the manifold hypothesis, which views the apparent high dimensionality of data as arising from variation around a much simpler underlying structure. Fefferman et al. (2016) connected this hypothesis to the statistical problem of determining whether a distribution can be well approximated by a smooth manifold, while Whiteley et al. (2026) explained how such structure can arise from latent variables, correlation, and stationarity. The purpose of manifold fitting is to estimate the geometric structure of possible signals in their original space, including their local variation and connectivity (Yao and Su, 2026).

The literature approaches manifold fitting from several directions. Minimax theory establishes fundamental limits for estimating a manifold under Hausdorff loss (Genovese et al., 2012a,b). Implicit reconstruction methods define the fitted manifold as a ridge or a zero set constructed from the data (Fefferman et al., 2018;

*Corresponding author: zhigang.yao@nus.edu.sg.

Genovese et al., 2014; Yao and Xia, 2025); the definition is global, but the manifold must still be recovered numerically. In practice, numerical manifold fitting commonly relies on neighbourhood information, either to recover an implicitly defined manifold or to construct fitted points directly. Two representative local methods are local PCA and mean shift. Local PCA estimates tangent geometry from a neighbourhood covariance matrix and projects points onto the resulting affine approximation (Cheng and Wu, 2013). Structure-adaptive manifold estimation also uses local geometric information, iteratively refining local weights to reconstruct a smooth manifold and denoise the observations (Puchkin and Spokoiny, 2022). Mean shift is a mode-seeking procedure based on a kernel density estimate (Comaniciu and Meer, 2002), while subspace-constrained mean shift (SCMS) adapts the mean-shift update to estimate density ridges (Genovese et al., 2014; Ozertem and Erdogmus, 2011).

The differing behaviour of local PCA and mean shift highlights two aspects of manifold fitting. Fitting accuracy measures whether the fitted points lie close to the underlying manifold, whereas geometric coverage measures whether they represent all parts of it. Figure 1 illustrates how the two methods affect these criteria. By retaining tangential positions, local PCA preserves broad geometric coverage but leaves visible variation normal to the sphere. Mean shift updates points along the full mean-shift vector, producing a more concentrated fit but also uneven density and local gaps.




Figure 1: (a) The underlying sphere; (b) the local PCA output; and (c) the mean-shift output. The same near-boundary region is enlarged in each panel.

These observations motivate Successive Tangent Space Projection (STSP), an iterative manifold-fitting algorithm whose nearby population fixed-point set forms a fitted manifold. Each update removes the estimated normal displacement from a point to its local mean while retaining its tangential position. The normal correction improves fitting accuracy, whereas the preservation of tangential position limits drift along the manifold and helps retain geometric coverage. Repeated application of the update gives an explicit construction of the fixed-point set. The multi-scale extension, MS-STSP, combines results obtained at several neighbourhood scales through squared-scale extrapolation to reduce curvature bias. Our contributions are as follows.

- *Local geometric refinement.* We introduce STSP, which combines local averaging and tangent-space projection to correct normal displacement while retaining tangential variation.
- *Fixed-point geometry and population convergence.* We show that the complete nearby population fixed-point set is a d -dimensional manifold whose nearest-point projection onto $\mathcal{M}$ is a diffeomorphism. When h is proportional to σ , this fixed-point manifold lies within $O(\sigma^2)$ of $\mathcal{M}$, and every sufficiently close population orbit converges geometrically to one of its points.
- *Finite-sample orbit localization.* For a fixed reference sample, we derive uniform local-moment and tangent-projector bounds and use them to control empirical orbits initialised within the local tube.

After a geometric transient, each orbit remains within $O(\sigma^2 + hr_N)$ of the underlying manifold, where r_N depends on the intrinsic local sample size Nh^d .

- *Multi-scale correction.* We develop MS-STSP by combining corresponding fixed-reference branches through squared-scale extrapolation. The theory shows that stable extrapolation preserves the $O(\sigma^2)$ localization order, while the experiments investigate its ability to reduce curvature-induced shrinkage.

Section 2 states the model and assumptions and defines the STSP update. Section 3 develops the local-moment expansions, population fixed-point manifold, and finite-sample localization theory. Section 4 analyses the curvature bias of local averaging and develops the multi-scale procedure. Section 5 reports numerical studies. Section 6 concludes. Full proofs are given in the Supplementary Material.

2 Manifold Fitting via Successive Tangent-Space Projection

2.1 Model, assumptions, and notation

Let $\mathcal{M} \subset \mathbb{R}^D$ be a compact d -dimensional manifold, with $1 \leq d < D$. We observe

$$\mathbf{y}_i = \mathbf{x}_i + \sigma \boldsymbol{\xi}_i, \quad i = 1, \dots, N,$$

where $\mathbf{x}_i \in \mathcal{M}$ is the signal and $\boldsymbol{\xi}_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(\mathbf{0}, I_D)$ is standardised ambient noise, independent of $\{\mathbf{x}_i\}_{i=1}^N$.

Assumption 1. The manifold and sampling satisfy the following.

- (A1) The signals $\{\mathbf{x}_i\}_{i=1}^N$ are drawn independently from the uniform distribution on $\mathcal{M}$.
- (A2) $\mathcal{M}$ is a C^5 embedded submanifold without boundary. In geodesic normal charts of the radius used below, the embedding and its derivatives through order five are bounded by a fixed constant K_{geo} .
- (A3) $\text{reach}(\mathcal{M}) \geq \tau > 0$, where

$$\text{reach}(\mathcal{M}) = \sup\{r > 0 : \text{every } \mathbf{y} \text{ with } \text{dist}(\mathbf{y}, \mathcal{M}) < r \text{ has a unique nearest point in } \mathcal{M}\}.$$

- (A4) Let $\iota_{\mathcal{M}} = \inf_{x \in \mathcal{M}} \text{inj}_x(\mathcal{M}) > 0$ be the global injectivity radius of the induced Riemannian metric, and set $r_{\text{geo}} = \min\{\tau, \iota_{\mathcal{M}}\}$. The dimensions d, D are fixed, $\sigma < 1$, and σ/r_{geo} is sufficiently small.

Conditions (A2) and (A3) control the geometric complexity of $\mathcal{M}$: within a neighbourhood of radius $h \ll \tau$, the manifold deviates from its tangent space by at most order h^2 , and the positive reach excludes sharp features and near self-intersections that would make recovery ill-posed. The injectivity radius in (A4) is the largest intrinsic scale on which geodesic normal coordinates are single-valued uniformly over $\mathcal{M}$. Condition (A4) ensures the noise is small relative to both the reach and injectivity scales, so that normal fluctuations can be distinguished from genuine structure.

We use the following notation throughout. For a nonempty set $\mathcal{S} \subset \mathbb{R}^D$, define $\text{dist}(\mathbf{y}, \mathcal{S}) = \inf_{\mathbf{s} \in \mathcal{S}} \|\mathbf{y} - \mathbf{s}\|$. For nonempty compact sets $\mathcal{S}_1$ and $\mathcal{S}_2$, their Hausdorff distance is $d_H(\mathcal{S}_1, \mathcal{S}_2) = \max\{\sup_{u \in \mathcal{S}_1} \text{dist}(u, \mathcal{S}_2), \sup_{v \in \mathcal{S}_2} \text{dist}(v, \mathcal{S}_1)\}$. Whenever the nearest point is unique, $\Pi_{\mathcal{S}}(\mathbf{y})$ denotes that point. In particular, if $\text{dist}(\mathbf{y}, \mathcal{M}) < \tau$, then (A3) gives the unique projection $\Pi_{\mathcal{M}}(\mathbf{y})$. For $x \in \mathcal{M}$, $T_x \mathcal{M}$ and $N_x \mathcal{M}$ denote the tangent and normal spaces, $\mathbf{P}_x$ is the orthogonal projector onto $T_x \mathcal{M}$, and $\mathbf{Q}_x := I_D - \mathbf{P}_x$ is the orthogonal projector onto $N_x \mathcal{M}$. Thus $\Pi_{\mathcal{S}}$ returns a projected point, whereas bold capital $\mathbf{P}$ and $\mathbf{Q}$ denote linear projector operators. The symbol Π_x denotes the second fundamental form, and I_m denotes the $m \times m$ identity matrix. For $\mathbf{b} \in \mathbb{R}^D$ and $r > 0$, $B_E(\mathbf{b}, r) = \{\mathbf{y} \in \mathbb{R}^D : \|\mathbf{y} - \mathbf{b}\| \leq r\}$ is the ambient Euclidean ball. For $x \in \mathcal{M}$, $B_g(x, r) = \{z \in \mathcal{M} : d_g(x, z) \leq r\}$ is the geodesic ball, and d_g is geodesic

distance. The symbol dA denotes the induced d -dimensional Riemannian volume measure on $\mathcal{M}$. If $0 < r < \text{inj}_x(\mathcal{M})$, the exponential map $\exp_x$ is one-to-one on $\{u \in T_x \mathcal{M} : \|u\| < r\}$; writing $z = \exp_x(u)$ gives geodesic normal coordinates centred at x , and then $d_g(x, z) = \|u\|$. Bold lowercase symbols denote observed, reference, or query vectors, whereas plain lowercase symbols denote deterministic manifold points or local-coordinate vectors. Hats denote empirical quantities, and parenthesized superscripts denote iterates or repeated composition. Constants may depend on $d, D, \tau, K_{\text{geo}}, \iota_{\mathcal{M}}$ and $\text{Area}(\mathcal{M})$, but not on N, σ , or h , and their values may change between occurrences.

2.2 From an implicit fitting equation to an explicit construction

Yao and Xia (2025) define a fitted manifold through the zero set of a smoothed local normal residual, denoted here by $\mathbf{G}_h$:

$$\widehat{\mathcal{M}}_{h,YX} = \{\mathbf{y} : \mathbf{G}_h(\mathbf{y}) = \mathbf{0}\}.$$

The fitted locations have zero estimated normal fitting bias, and their analysis places this zero-level set close to the underlying manifold. This construction provides a geometric starting point for manifold fitting: characterizing fitted locations through local normal residuals. However, an implicit equation characterizes the fitted set but still requires a numerical procedure to produce points on it. Their implementation minimizes $\|\mathbf{G}_h(\mathbf{y})\|_2^2$ by gradient descent with line search. Membership in the fitted set requires a zero objective, whereas a stationary numerical solution may have a positive objective. This distinction motivates an explicit construction whose iterates can be analysed directly relative to the underlying manifold.

For an affine-space correction, let $\mathbf{m}(\mathbf{y})$ be a weighted local centre, let $T_{\text{loc}}(\mathbf{y})$ be a local tangent space, and let $\mathbf{P}_{\text{loc}}(\mathbf{y})$ be its orthogonal projector. The associated local affine space and normal residual are

$$\mathcal{A}(\mathbf{y}) = \mathbf{m}(\mathbf{y}) + T_{\text{loc}}(\mathbf{y}), \quad \mathbf{R}(\mathbf{y}) = \mathbf{y} - \Pi_{\mathcal{A}(\mathbf{y})}(\mathbf{y}) = \{I_D - \mathbf{P}_{\text{loc}}(\mathbf{y})\}\{\mathbf{y} - \mathbf{m}(\mathbf{y})\}.$$

The zero-residual set of this affine-space construction is

$$\widehat{\mathcal{M}} = \{\mathbf{y} : \mathbf{R}(\mathbf{y}) = \mathbf{0}\}.$$

Its geometric condition is that the displacement from the local centre has zero projection onto the estimated normal space. Accordingly, for any affine-space residual of the form above, define the corresponding explicit correction by

$$\mathcal{T}(\mathbf{y}) = \mathbf{m}(\mathbf{y}) + \mathbf{P}_{\text{loc}}(\mathbf{y})\{\mathbf{y} - \mathbf{m}(\mathbf{y})\} = \mathbf{y} - \mathbf{R}(\mathbf{y}).$$

Then $\widehat{\mathcal{M}} = \text{Fix}(\mathcal{T})$: the implicit zero-residual equation is equivalently the fixed-point equation of a local tangent-space correction.

Motivated by the normal-residual characterization, STSP constructs an explicit iteration from hard-radius conditional moments. Section 3 studies its fixed-point set and local trajectories. Let $\mathbf{Y}$ denote a generic observation from the model and set $p_h(\mathbf{y}) = \Pr\{\mathbf{Y} \in B_E(\mathbf{y}, h)\}$. Wherever $p_h(\mathbf{y}) > 0$, let $\boldsymbol{\mu}_h(\mathbf{y})$ and $\boldsymbol{\Sigma}_h(\mathbf{y})$ be the conditional mean and covariance of $\mathbf{Y}$ given $\mathbf{Y} \in B_E(\mathbf{y}, h)$. Let $\mathcal{D}_h$ collect points with positive local probability and a strict eigengap. For $\mathbf{y} \in \mathcal{D}_h$, let $T_h(\mathbf{y})$ be the uniquely defined leading- d eigenspace of $\boldsymbol{\Sigma}_h(\mathbf{y})$, and let $\mathbf{P}_h(\mathbf{y})$ be its orthogonal projector. The population domain, local affine space, residual, fitting object, and explicit correction are

$$\mathcal{D}_h = \{\mathbf{y} : p_h(\mathbf{y}) > 0, \lambda_d\{\boldsymbol{\Sigma}_h(\mathbf{y})\} > \lambda_{d+1}\{\boldsymbol{\Sigma}_h(\mathbf{y})\}\},$$

$$\mathcal{A}_h(\mathbf{y}) = \boldsymbol{\mu}_h(\mathbf{y}) + T_h(\mathbf{y}),$$

$$\mathbf{F}_h(\mathbf{y}) = \{I_D - \mathbf{P}_h(\mathbf{y})\}\{\mathbf{y} - \boldsymbol{\mu}_h(\mathbf{y})\}, \quad \mathcal{S}_{h,\text{pop}} = \{\mathbf{y} \in \mathcal{D}_h : \mathbf{F}_h(\mathbf{y}) = \mathbf{0}\},$$

$$\mathcal{T}_h(\mathbf{y}) = \Pi_{\mathcal{A}_h(\mathbf{y})}(\mathbf{y}) = \boldsymbol{\mu}_h(\mathbf{y}) + \mathbf{P}_h(\mathbf{y})\{\mathbf{y} - \boldsymbol{\mu}_h(\mathbf{y})\} = \mathbf{y} - \mathbf{F}_h(\mathbf{y}).$$

Consequently, $\mathcal{S}_{h,\text{pop}} = \text{Fix}(\mathcal{T}_h)$, and manifold fitting is implemented by the successive corrections $\mathbf{y}^{(t+1)} = \mathcal{T}_h(\mathbf{y}^{(t)})$. Figure 2 illustrates normal-residual fitting and successive tangent-space corrections.




Figure 2: Implicit and explicit manifold fitting. (a) Blue intensity represents the length $\|\mathbf{R}(\mathbf{y})\|$ of the local affine-space residual, and the white locus is the generic fitted set $\widehat{\mathcal{M}} = \{\mathbf{y} : \mathbf{R}(\mathbf{y}) = \mathbf{0}\}$. (b) A schematic sequence of projections onto the query-local affine spaces $\mathcal{A}(\mathbf{y})$ approaching the fitted set. For STSP, the residual and affine space are $\mathbf{R} = \mathbf{F}_h$ and $\mathcal{A} = \mathcal{A}_h$, respectively.

2.3 The STSP update

Set $\mathbf{z}_j = \mathbf{y}_j$ and keep $\mathcal{Z} = \{\mathbf{z}_j\}_{j=1}^N$ fixed throughout the algorithm. At iteration t only the query locations $\widehat{\mathbf{y}}_i^{(t)}$ change. For each query, first form the radius- h neighbourhood in the fixed reference cloud,

$$\mathcal{I}_{\mathcal{Z},h}(\mathbf{y}) = \{j : \|\mathbf{z}_j - \mathbf{y}\| \leq h\}.$$

Let $n_{\mathcal{Z},h}(\mathbf{y}) = |\mathcal{I}_{\mathcal{Z},h}(\mathbf{y})|$ denote the corresponding count. The radius h must be large enough for stable estimation and small enough that the manifold is nearly linear within it. Whenever $n_{\mathcal{Z},h}(\mathbf{y}) > 0$, compute the local mean $\widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y}) = n_{\mathcal{Z},h}(\mathbf{y})^{-1} \sum_{j \in \mathcal{I}_{\mathcal{Z},h}(\mathbf{y})} \mathbf{z}_j$ and covariance

$$\widehat{\boldsymbol{\Sigma}}_{\mathcal{Z},h}(\mathbf{y}) = \frac{1}{n_{\mathcal{Z},h}(\mathbf{y})} \sum_{j \in \mathcal{I}_{\mathcal{Z},h}(\mathbf{y})} \{\mathbf{z}_j - \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y})\} \{\mathbf{z}_j - \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y})\}^\top.$$

If, in addition, the d th covariance eigenvalue is strictly greater than the $(d+1)$ st, let $\widehat{T}_{\mathcal{Z},h}(\mathbf{y})$ be its uniquely defined leading- d eigenspace, and let $\widehat{\mathbf{P}}_{\mathcal{Z},h}(\mathbf{y})$ be the orthogonal projector onto $\widehat{T}_{\mathcal{Z},h}(\mathbf{y})$. On this empirical domain, define

$$\widehat{\mathcal{T}}_{\mathcal{Z},h}(\mathbf{y}) = \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y}) + \widehat{\mathbf{P}}_{\mathcal{Z},h}(\mathbf{y})\{\mathbf{y} - \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y})\}.$$

All iterations apply this same deterministic map through $\widehat{\mathbf{y}}_i^{(t+1)} = \widehat{\mathcal{T}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}_i^{(t)})$. Conditional on $\mathcal{Z}$, the N query orbits are uncoupled. The update retains the estimated tangential component relative to the local mean and discards the estimated normal component. The hard-radius operator is well defined along a trajectory when every visited neighbourhood is nonempty and its d th and $(d+1)$ st covariance eigenvalues are separated. The finite-sample theory below supplies uniform conditions for both properties.

2.4 Relation to local PCA, mean shift, and SCMS

Write $\mathbf{Q}_h(\mathbf{y}) = \mathbf{I}_D - \mathbf{P}_h(\mathbf{y})$. The population STSP update has the projected-shift form

$$\mathcal{T}_h(\mathbf{y}) = \mathbf{y} + \mathbf{Q}_h(\mathbf{y})\{\boldsymbol{\mu}_h(\mathbf{y}) - \mathbf{y}\}. \quad (1)$$

Algorithm 1 Successive Tangent-Space Projection (STSP)

Require: Reference points $\mathcal{Z} = \{\mathbf{z}_j\}_{j=1}^N \subset \mathbb{R}^D$, initial queries $\widehat{\mathbf{y}}_i^{(0)} = \mathbf{z}_i$, dimension d , radius h , and iteration horizon L .

Ensure: Fitted queries $\{\widehat{\mathbf{y}}_i^{(L)}\}_{i=1}^N$.

```

1: for  $t = 0$  to  $L - 1$  do
2:   for  $i = 1$  to  $N$  do
3:      $\mathcal{I}_i^{(t)} := \mathcal{I}_{\mathcal{Z},h}(\widehat{\mathbf{y}}_i^{(t)})$ .
4:     Set  $\widehat{\boldsymbol{\mu}}_i^{(t)} := \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}_i^{(t)})$  and  $\widehat{\boldsymbol{\Sigma}}_i^{(t)} := \widehat{\boldsymbol{\Sigma}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}_i^{(t)})$ , computed from  $\{\mathbf{z}_j : j \in \mathcal{I}_i^{(t)}\}$ .
5:     Set  $\widehat{\mathbf{P}}_i^{(t)} := \widehat{\mathbf{P}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}_i^{(t)})$ , the leading- $d$  projector of  $\widehat{\boldsymbol{\Sigma}}_i^{(t)}$ .
6:      $\widehat{\mathbf{y}}_i^{(t+1)} = \widehat{\boldsymbol{\mu}}_i^{(t)} + \widehat{\mathbf{P}}_i^{(t)}(\widehat{\mathbf{y}}_i^{(t)} - \widehat{\boldsymbol{\mu}}_i^{(t)})$ .
7:   end for
8: end for
9: return  $\{\widehat{\mathbf{y}}_i^{(L)}\}_{i=1}^N$ .

```

It therefore applies the estimated normal component of the displacement from the query to its hard-radius conditional mean while preserving the query’s tangential component.

The local-PCA comparator used in Section 5 applies one projection onto an estimated affine tangent space. STSP recomputes the query-local conditional moments as the query moves and repeatedly applies the normal correction in (1), while the distribution or sample that supplies those moments remains fixed. Mean shift instead takes the full displacement to the local mean.

This algebraic form is closely related to SCMS. A standard SCMS update can be written schematically as

$$\mathcal{T}_{\text{SCMS},h}(\mathbf{y}) = \mathbf{y} + \widetilde{\mathbf{Q}}_h(\mathbf{y})\{\widetilde{\boldsymbol{\mu}}_h(\mathbf{y}) - \mathbf{y}\},$$

where $\widetilde{\boldsymbol{\mu}}_h(\mathbf{y}) - \mathbf{y}$ is a kernel mean-shift vector and $\widetilde{\mathbf{Q}}_h(\mathbf{y})$ projects onto the estimated normal eigenspace of a kernel-density Hessian. For Gaussian weights, write $\boldsymbol{\mu}_{G,h}$ and $\boldsymbol{\Sigma}_{G,h}$ for the weighted mean and covariance, and $f_{G,h}$ for the corresponding Gaussian kernel density estimate. Then $\nabla \log f_{G,h} = (\boldsymbol{\mu}_{G,h} - \mathbf{y})/h^2$ and $\nabla^2 \log f_{G,h} = \boldsymbol{\Sigma}_{G,h}/h^4 - I_D/h^2$. Thus a projected update using Gaussian moments and the leading covariance eigenspace has the same local form as SCMS-log-KDE. The STSP studied here instead uses hard-radius conditional moments. Our contribution concerns the geometry of its nearby fixed-point set and finite-sample control of its iterates, rather than the projected mean-shift form alone. Section 3 establishes these local results. Table 1 summarizes the local quantities, updates, and targets of these methods.

Table 1: Schematic projection-based comparison of the local updates considered here.

Method	Local quantities	Update	Target or role
Mean shift	Local mean	Full displacement toward the local mean	Density modes
SCMS	Kernel mean-shift vector and normal eigenspace of the density Hessian	Normal projection of the kernel mean-shift vector	Density ridge
Local PCA	Estimated affine tangent space	Projection onto the affine tangent space	Local affine approximation
STSP	Local mean and covariance within a fixed radius	Repeated normal correction using fixed reference data	Fixed-point manifold near the underlying manifold

3 Theoretical Analysis

Fix $c > 0$ and set $h = c\sigma$, with $c\sigma \leq r_{\text{geo}}/2$. The population results are stated as $\sigma \rightarrow 0$. In the finite-sample results, $N \rightarrow \infty$ jointly with $\sigma \rightarrow 0$. With d, D, c, A fixed, the condition $N\sigma^{d+2}/\{D \log(1/\sigma) + \log(1/\delta)\} \geq C_0$ for a sufficiently large constant C_0 gives uniform empirical consistency and preserves the population $O(\sigma^2)$ localization order.

3.1 Flat reference model and uniform local expansions

Proofs of the results in this subsection are given in Supplementary Sections S2.1–S2.2. Let $\Phi_\rho = \{\mathbf{y} : \text{dist}(\mathbf{y}, \mathcal{M}) \leq \rho\}$. For $\mathbf{y} \in \Phi_\rho$, write $x = \Pi_{\mathcal{M}}(\mathbf{y})$ and $\mathbf{v}(\mathbf{y}) = \mathbf{y} - x \in N_x \mathcal{M}$. We analyse the population operator on $\Phi_{A\sigma}$ for a fixed $A < \infty$.

Put $q = D - d$. For $a \in \mathbb{R}^q$, let $W = (W_T, W_N) \in \mathbb{R}^d \times \mathbb{R}^q$ have density

$$p_{c,a}(w) = \frac{\mathbf{1}\{\|w\| \leq c\} \exp\{-\|a + w_N\|^2/2\}}{\int_{\|v\| \leq c} \exp\{-\|a + v_N\|^2/2\} dv}. \quad (2)$$

Let $\mathbb{E}_{c,a}$ and $\text{cov}_{c,a}$ denote expectation and covariance under $p_{c,a}$, respectively. Define

$$m_c(a) = a + \mathbb{E}_{c,a} W_N, \quad S_c(a) = \text{cov}_{c,a}(W).$$

The law in (2) is the exact scaled conditional law for an affine d -plane. We call (c, A) *spectrally admissible* when there is $g_0 > 0$ such that, for every $\|a\| \leq A$,

$$\lambda_{\min}\{\text{cov}_{c,a}(W_T)\} - \lambda_{\max}\{\text{cov}_{c,a}(W_N)\} \geq g_0. \quad (3)$$

The two ingredients have distinct roles. Spectral admissibility identifies the leading- d covariance eigenspace with the tangent space, whereas the score identity

$$Dm_c(a) = I_q - \text{cov}_{c,a}(W_N) \quad (4)$$

and the Brascamp–Lieb inequality imply $\sup_{\|a\| \leq A} \|Dm_c(a)\|_{\text{op}} \leq \kappa_0 < 1$.

Remark 1 (Scale-free spectral condition). The normalised gap in (3) depends only on (d, q, c, A) , where $q = D - d$, and not on σ , the reach, or other geometric features of $\mathcal{M}$. It can therefore be checked once for each dimension pair, radius ratio, and normalised tube width. Supplementary Section S2.1 reports the minimum gaps for the values used in the numerical studies.

Lemma 1 (Uniform local-moment expansion). *Under Assumption 1 and spectral admissibility, let $\mathbf{y} \in \Phi_{A\sigma}$, $x = \Pi_{\mathcal{M}}(\mathbf{y})$, and write $\mathbf{v}(\mathbf{y}) = \sigma a$ in orthonormal tangent–normal frames at x . Uniformly over $\Phi_{A\sigma}$,*

$$\mu_h(\mathbf{y}) = x + \sigma(\mathbf{0}_d, m_c(a)) + O(\sigma^2), \quad \Sigma_h(\mathbf{y}) = \sigma^2 S_c(a) + O(\sigma^3),$$

and

$$\|\mathbf{P}_h(\mathbf{y}) - \mathbf{P}_x\|_{\text{op}} \leq C\sigma.$$

Consequently,

$$\mathbf{Q}_x\{\mathcal{T}_h(\mathbf{y}) - x\} = \sigma m_c(a) + O(\sigma^2), \quad \|\mathbf{P}_x\{\mathcal{T}_h(\mathbf{y}) - x\}\| \leq C\sigma^2.$$

3.2 Population fixed-point manifold and attraction

Proofs of the population recursion and fixed-point manifold theorem are given in Supplementary Section S2.2. Throughout this subsection, (c, A) is spectrally admissible and $c\sigma \leq r_{\text{geo}}/2$.

Theorem 1 (Local normal-distance recursion). *Under Assumption 1 and the spectral admissibility condition above, there are $\sigma_{\text{con}} > 0$, $C < \infty$, and $\kappa \in (\kappa_0, 1)$ such that, for $\sigma \leq \sigma_{\text{con}}$ and $\mathbf{y} \in \Phi_{A\sigma}$,*

$$\text{dist}(\mathcal{T}_h(\mathbf{y}), \mathcal{M}) \leq \kappa \text{dist}(\mathbf{y}, \mathcal{M}) + C\sigma^2.$$

For sufficiently small σ , $\Phi_{A\sigma}$ is forward invariant, and any population orbit starting there obeys

$$\text{dist}(\mathcal{T}_h^{(t)}(\mathbf{y}), \mathcal{M}) \leq \kappa^t \text{dist}(\mathbf{y}, \mathcal{M}) + \frac{C\sigma^2}{1 - \kappa}.$$

To lift normal localization to ambient convergence, we identify the complete nearby zero set of the update residual and control the map transverse to that set. Recall $\mathbf{F}_h$ and $\mathcal{S}_{h,\text{pop}}$ from Section 2.2.

Theorem 2 (Population fixed-point manifold and local orbit convergence). *Fix $0 < A_0 < A$. Under the conditions of Theorem 1, there are $C, C_{\text{step}}, \rho_0 > 0$ and $\bar{\kappa} \in (0, 1)$ such that, for all sufficiently small σ , the fixed-point set in $\Phi_{A_0\sigma}$ is the compact C^1 normal graph*

$$\mathcal{M}_h = \{x + b_h(x) : x \in \mathcal{M}, b_h(x) \in N_x \mathcal{M}\}, \quad \|b_h\|_{C^1} \leq C\sigma^2.$$

Its nearest-point projection onto $\mathcal{M}$ is a C^1 diffeomorphism and

$$d_H(\mathcal{M}_h, \mathcal{M}) \leq C\sigma^2.$$

There is a neighbourhood $\mathcal{U}_h = \{\mathbf{y} : \text{dist}(\mathbf{y}, \mathcal{M}_h) < \rho_0\sigma\} \subset \Phi_{A\sigma}$ such that every $\mathbf{y}^{(0)} \in \mathcal{U}_h$ generates an orbit $\mathbf{y}^{(t+1)} = \mathcal{T}_h(\mathbf{y}^{(t)})$ that converges to a fixed point $\mathbf{y}^{(\infty)} \in \mathcal{M}_h$. More precisely,

$$\text{dist}(\mathbf{y}^{(t)}, \mathcal{M}_h) \leq \bar{\kappa}^t \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_h)$$

and

$$\|\mathbf{y}^{(t)} - \mathbf{y}^{(\infty)}\| \leq \frac{C_{\text{step}} \bar{\kappa}^t}{1 - \bar{\kappa}} \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_h).$$

In particular, $\text{dist}(\mathbf{y}^{(\infty)}, \mathcal{M}) \leq C\sigma^2$.

3.3 Finite-sample localization

Proofs of the uniform empirical approximation, fixed-point localization, and orbit localization results are given in Supplementary Section S3.1. We analyse the empirical operator $\widehat{\mathcal{T}}_{\mathcal{Z},h}$ constructed from the fixed i.i.d. reference sample $\mathcal{Z}$.

Let $0 < \delta < 1$ and define

$$r_N = \sqrt{\frac{D \log(1/h) + \log(1/\delta)}{Nh^d}} + \frac{D \log(1/h) + \log(1/\delta)}{Nh^d}.$$

Lemma 2 (Uniform empirical approximation). *Under the conditions of Lemma 1, there are constants $C, r_0, c_{\text{gap}} > 0$ such that, if $r_N \leq r_0$, then with probability at least $1 - \delta$,*

$$\begin{aligned} \sup_{\Phi_{A\sigma}} \|\widehat{\boldsymbol{\mu}}_{\mathcal{Z},h} - \boldsymbol{\mu}_h\| &\leq Chr_N, & \sup_{\Phi_{A\sigma}} \|\widehat{\boldsymbol{\Sigma}}_{\mathcal{Z},h} - \boldsymbol{\Sigma}_h\|_{\text{op}} &\leq Ch^2 r_N, \\ \sup_{\Phi_{A\sigma}} \|\widehat{\mathbf{P}}_{\mathcal{Z},h} - \mathbf{P}_h\|_{\text{op}} &\leq Cr_N, & \inf_{\Phi_{A\sigma}} \{\widehat{\lambda}_d - \widehat{\lambda}_{d+1}\} &\geq c_{\text{gap}} h^2. \end{aligned}$$

Corollary 1 (Localization of empirical fixed points). *Under the conditions of Theorem 1, there are constants $C, r_{\text{fp}} > 0$ such that, if $r_N \leq r_{\text{fp}}$, then with probability at least $1 - \delta$, every fixed point $\widehat{\mathbf{y}}_N \in \Phi_{A\sigma}$ of $\widehat{\mathcal{T}}_{\mathcal{Z},h}$, whenever one exists, satisfies*

$$\text{dist}(\widehat{\mathbf{y}}_N, \mathcal{M}) \leq C\{\sigma^2 + hr_N\}.$$

Consequently, for fixed $d, D, c, A, h = c\sigma$ and

$$N \geq C\sigma^{-(d+2)}\{D \log(1/\sigma) + \log(1/\delta)\}$$

suffice for the right-hand side to be $O(\sigma^2)$.

Theorem 3 (Finite-sample orbit localization). *For the same fixed $0 < A_0 < A$, under the conditions of Theorem 1 and Lemma 2, there are $\sigma_{\text{orb}}, r_{\text{orb}}, C > 0$ such that, if $\sigma \leq \sigma_{\text{orb}}$ and $r_N \leq r_{\text{orb}}$, then with probability at least $1 - \delta$, every empirical orbit*

$$\widehat{\mathbf{y}}^{(t+1)} = \widehat{\mathcal{T}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}^{(t)}), \quad \widehat{\mathbf{y}}^{(0)} \in \Phi_{A_0\sigma},$$

remains in $\Phi_{A\sigma}$ and satisfies, for every $t \geq 0$,

$$\text{dist}(\widehat{\mathbf{y}}^{(t)}, \mathcal{M}) \leq \kappa^t \text{dist}(\widehat{\mathbf{y}}^{(0)}, \mathcal{M}) + \frac{C(\sigma^2 + hr_N)}{1 - \kappa}.$$

In particular, after the geometric transient the entire empirical orbit lies in an $O(\sigma^2 + hr_N)$ neighbourhood of $\mathcal{M}$; if $r_N = O(\sigma)$ and $h = c\sigma$, this neighbourhood has width $O(\sigma^2)$.

Theorem 3 is uniform over all initial queries in $\Phi_{A_0\sigma}$, including data-dependent observations that fall in this tube. Supplementary Section S3.2 quantifies the corresponding fraction of observed initializations. It also gives the theoretical iteration complexity. In particular, if

$$L \geq \frac{\lceil \log \{ (1 - \kappa) d_0 / [C(\sigma^2 + hr_N)] \} \rceil_+}{|\log \kappa|}, \quad d_0 = \text{dist}(\widehat{\mathbf{y}}^{(0)}, \mathcal{M}),$$

the transient term is no larger than the statistical error radius. Thus $L = O\{\log(1/\sigma)\}$ when $d_0 = O(\sigma)$ and $\sigma^2 + hr_N = O(\sigma^2)$.

4 Multi-scale STSP

The radius h controls a bias–variance trade-off: large neighbourhoods stabilize local moments but increase curvature bias, whereas small neighbourhoods can give unstable tangent estimates. MS-STSP combines a fixed finite collection of radii. The numerical studies use three radii; the theory below allows any fixed finite number of radii.

4.1 Motivation, algorithm, and localization results

Proofs of the quadratic bias expansion and the two MS-STSP localization theorems are given in Supplementary Sections S4.1, S4.4, and S4.5; Supplementary Sections S4.2–S4.3 provide the supporting weight and aggregation results. For a sufficiently small fixed geometric constant $c_0 \in (0, 1)$ and $0 < h \leq c_0 r_{\text{geo}}$, define the clean local-average map

$$f_h(x) = \frac{1}{\text{Area}\{B_E(x, h) \cap \mathcal{M}\}} \int_{B_E(x, h) \cap \mathcal{M}} y \, dA(y), \quad x \in \mathcal{M}.$$

Lemma 3 (Quadratic structure of local averaging bias). *Under conditions (A2)–(A4), uniformly in $x \in \mathcal{M}$,*

$$f_h(x) = x + \frac{1}{2}h^2\mathbf{B}(x) + O(h^4).$$

Here, for any orthonormal basis $\{e_1, \dots, e_d\}$ of $T_x\mathcal{M}$,

$$\mathbf{B}(x) = \frac{1}{d+2} \sum_{k=1}^d \Pi_x(e_k, e_k) \in N_x\mathcal{M}, \quad (5)$$

and the remainder is uniform in x .

For the same x , the vector $\mathbf{B}(x)$ is independent of h . Consequently, the ordinary least-squares intercept weights for a regression on $s_j = h_j^2$ satisfy

$$\sum_{j=1}^M w_j = 1, \quad \sum_{j=1}^M w_j h_j^2 = 0,$$

and cancel the displayed h^2 term in this ideal clean local-average model. The clean local-average expansion motivates extrapolation in the squared neighbourhood radius. For noisy STSP branches, the results below establish that stable aggregation preserves the single-scale localization order.

Algorithm 2 Multi-Scale STSP (MS-STSP)

Require: Original cloud $\mathcal{Z} = \{\mathbf{z}_i\}_{i=1}^N$, dimension d , radii $\{h_j\}_{j=1}^M$ with $M \geq 2$ and at least two distinct values, and iteration horizon $L \geq 1$.

Ensure: Multi-scale estimates $\{\hat{\mathbf{y}}_{i,\text{MS}}\}_{i=1}^N$.

- 1: **for** $j = 1, \dots, M$ **do**
 - 2: Apply Algorithm 1 with reference $\mathcal{Z}$, initial queries $\hat{\mathbf{y}}_{i,j}^{(0)} = \mathbf{z}_i$, dimension d , radius h_j , and horizon L , obtaining $\{\hat{\mathbf{y}}_{i,j}^{(L)}\}_{i=1}^N$.
 - 3: **end for**
 - 4: Compute the OLS intercept weights $w_1, \dots, w_M$ for the regressor h_j^2 .
 - 5: **for** $i = 1, \dots, N$ **do**
 - 6: $\hat{\mathbf{y}}_{i,\text{MS}} \leftarrow \sum_{j=1}^M w_j \hat{\mathbf{y}}_{i,j}^{(L)}$.
 - 7: **end for**
 - 8: **return** $\{\hat{\mathbf{y}}_{i,\text{MS}}\}_{i=1}^N$.
-

The following theorems establish population and finite-sample localization for the fixed-reference multi-scale branches in Algorithm 2. The finite-sample result permits scale-specific horizons L_j ; Algorithm 2 is the common case $L_j = L$.

Theorem 4 (Population MS-STSP localization). *Let M be fixed and $h_j = c_j\sigma$. Suppose every (c_j, A) is spectrally admissible, let the common initial point satisfy $\mathbf{y}^{(0)} \in \cap_{j=1}^M \mathcal{U}_{h_j}$, and suppose the weights satisfy*

$$\sum_j w_j = 1, \quad \sum_j |w_j| \leq C_w,$$

where C_w is a fixed design bound. Let $\mathbf{y}_j^{(\infty)} = \lim_{t \rightarrow \infty} \mathcal{T}_{h_j}^{(t)}(\mathbf{y}^{(0)})$ and $\mathbf{y}_{\text{MS}}^{(\infty)} = \sum_j w_j \mathbf{y}_j^{(\infty)}$. Then all projected points $\Pi_{\mathcal{M}}(\mathbf{y}_j^{(\infty)})$ lie in one $O(\sigma)$ geodesic patch and

$$\text{dist}(\mathbf{y}_{\text{MS}}^{(\infty)}, \mathcal{M}) \leq CC_w \sigma^2.$$

Thus population MS-STSP retains the $O(\sigma^2)$ localization order of the single-scale branches.

Theorem 5 (Finite-sample MS-STSP localization). *Let $0 < \delta < 1$. For $j = 1, \dots, M$, define*

$$r_{N,j} = \sqrt{\frac{D \log(1/h_j) + \log(M/\delta)}{Nh_j^d}} + \frac{D \log(1/h_j) + \log(M/\delta)}{Nh_j^d}.$$

Assume the conditions of Theorem 4 and the initialization, sampling-error, and finite-horizon conditions of Supplementary Theorem S5 in Section S4.5. These conditions keep each branch in its attraction tube, reduce its transient error to the localization scale, and bound accumulated tangential drift up to its prescribed horizon L_j . For $t = 0, \dots, L_j$, define the empirical branch orbit by $\widehat{\mathbf{y}}_j^{(t)} = \widehat{\mathcal{T}}_{\mathcal{Z},h_j}^{(t)}(\mathbf{y}^{(0)})$, and set $\widehat{\mathbf{y}}_{\text{MS}} = \sum_j w_j \widehat{\mathbf{y}}_j^{(L_j)}$. Then, with probability at least $1 - \delta$,

$$\text{dist}(\widehat{\mathbf{y}}_{\text{MS}}, \mathcal{M}) \leq CC_w \left\{ \sigma^2 + \max_j h_j r_{N,j} \right\}.$$

In particular, if $h_j = c_j \sigma$ and

$$N \geq C \sigma^{-(d+2)} \{D \log(1/\sigma) + \log(M/\delta)\},$$

the right-hand side is $O(\sigma^2)$.

5 Numerical Studies

5.1 Experimental setup

We evaluate STSP and MS-STSP on synthetic manifolds and mouse dentate-gyrus single-cell data. The synthetic study uses four curves (circle, tennis curve, trefoil knot, and sixfolds) and two surfaces (sphere and torus), shown in Supplementary Figure S1, with ten independently generated noisy clouds per setting. Let $\mathcal{Y} = \{\mathbf{y}_i\}_{i=1}^N$ denote the noisy input and $\widehat{\mathcal{Y}} = \{\widehat{\mathbf{y}}_i\}_{i=1}^N$ a fitted output; data-generation and implementation details are given in Supplementary Section S5. Let $\mathcal{X}_{\text{eval}} = \{\mathbf{x}_{\text{eval},\ell}\}_{\ell=1}^{N_{\text{eval}}} \subset \mathcal{M}$ be an independent dense evaluation cloud, fixed across repetitions within a setting. Within each repetition, all methods use the same $\mathcal{Y}$ and $\mathcal{X}_{\text{eval}}$.

We use the following two metrics to evaluate $\widehat{\mathcal{Y}}$.

$$e_{\text{fit}} = \frac{1}{N} \sum_{i=1}^N \text{dist}(\widehat{\mathbf{y}}_i, \mathcal{X}_{\text{eval}}), \quad e_{\text{cov}} = \frac{1}{N_{\text{eval}}} \sum_{\ell=1}^{N_{\text{eval}}} \text{dist}(\mathbf{x}_{\text{eval},\ell}, \widehat{\mathcal{Y}}).$$

Both metrics are means over all points in their respective clouds. The first approximates the distance from the fitted cloud to the underlying manifold using the dense evaluation cloud, whereas the second measures how well the fitted cloud covers that evaluation cloud. Smaller values are better.

We compare MS-STSP, STSP, YY-adapted, MF25, local PCA (Cheng and Wu, 2013), and mean shift (Comaniciu and Meer, 2002), and two SCMS variants (Ozertem and Erdogmus, 2011; Zhang and Chen, 2023). SCMS-KDE uses the Hessian of a Gaussian kernel density estimate to define its projection subspace; SCMS-log-KDE uses the Hessian of the log density. Both project the Gaussian mean-shift vector onto the selected normal subspace. We refer to the estimator of Yao et al. (2023) as the Yao–Yau (YY) estimator, following the terminology of Yao and Su (2026). YY-adapted denotes our dimension-aware adaptation of its local mean-displacement construction, using a d -dimensional tangent estimate to construct its averaging neighbourhood. MF25 numerically approximates the residual zero set of the method described in Section 2.2. The curve experiments use $N = 5000$ and $\sigma = 0.08$ for all four curves, with an additional circle experiment at $\sigma \in \{0.1, 0.2, 0.3\}$. The surface experiments use $N = 20000$ and $\sigma = 0.08$ for both surfaces. Noise is isotropic Gaussian in $\mathbb{R}^3$. For each method with tunable parameters, the synthetic experiments select the candidate minimizing $J_{\text{mean}} = \max(e_{\text{fit}}, e_{\text{cov}})$. The implementation code is available at <https://github.com/zhigang-yao/STSP>.

5.2 Fitting and coverage errors on synthetic manifolds

We first examine how the competing methods balance fitting accuracy and geometric coverage across the six synthetic manifolds; Table 2 summarises the corresponding errors, while Figure 3 illustrates the fitted clouds on the torus.

Table 2: Fitting and coverage errors over ten independent realisations on six synthetic manifolds (mean $\pm$ standard deviation, in units of 10^{-3}). For each manifold and error measure, the smallest mean is bold and the second smallest is underlined; rankings use unrounded means.

Method	Error	Circle	Tennis curve	Trefoil knot	Sixfolds	Sphere	Torus
Local PCA	e_{fit}	18.90 ± 0.60	22.56 ± 0.87	15.39 ± 0.63	33.11 ± 0.84	24.67 ± 0.30	32.55 ± 0.37
	e_{cov}	<u>8.07 ± 0.45</u>	<u>11.19 ± 0.57</u>	<u>12.76 ± 0.44</u>	<u>14.31 ± 0.28</u>	20.41 ± 0.19	<u>21.56 ± 0.20</u>
Mean shift	e_{fit}	15.05 ± 3.99	23.07 ± 1.40	24.51 ± 6.75	25.61 ± 3.44	21.72 ± 0.57	<u>23.31 ± 0.76</u>
	e_{cov}	24.66 ± 2.24	29.54 ± 1.74	39.42 ± 2.22	31.26 ± 2.27	42.13 ± 1.07	43.19 ± 0.67
SCMS-KDE	e_{fit}	25.56 ± 1.00	30.94 ± 1.01	15.91 ± 0.96	47.04 ± 1.94	29.32 ± 0.56	40.03 ± 0.75
	e_{cov}	17.27 ± 0.78	22.81 ± 1.21	14.87 ± 0.71	24.19 ± 1.44	22.06 ± 0.36	24.86 ± 0.38
SCMS-log-KDE	e_{fit}	10.68 ± 0.71	14.11 ± 1.21	12.86 ± 0.78	22.24 ± 1.43	18.33 ± 0.46	27.66 ± 0.55
	e_{cov}	10.66 ± 0.75	14.19 ± 1.22	13.58 ± 0.70	20.44 ± 1.54	19.71 ± 0.35	28.13 ± 0.53
YY-adapted	e_{fit}	<u>8.91 ± 0.83</u>	<u>11.44 ± 0.80</u>	15.32 ± 0.58	<u>18.41 ± 0.86</u>	<u>17.23 ± 0.33</u>	25.86 ± 0.47
	e_{cov}	8.95 ± 0.71	11.75 ± 0.84	17.33 ± 0.58	18.56 ± 0.85	23.51 ± 0.23	29.85 ± 0.46
MF25	e_{fit}	10.15 ± 0.81	13.37 ± 1.01	<u>12.54 ± 0.77</u>	21.50 ± 1.45	17.62 ± 0.44	30.90 ± 0.98
	e_{cov}	10.12 ± 0.78	13.33 ± 1.07	13.26 ± 0.69	20.76 ± 1.48	19.22 ± 0.36	23.97 ± 0.50
STSP	e_{fit}	10.27 ± 0.80	13.31 ± 1.00	12.81 ± 0.69	20.48 ± 1.13	18.57 ± 0.38	24.01 ± 0.42
	e_{cov}	9.85 ± 0.81	12.77 ± 1.00	13.27 ± 0.61	17.53 ± 1.53	<u>19.16 ± 0.25</u>	23.77 ± 0.37
MS-STSP	e_{fit}	5.98 ± 0.69	8.92 ± 0.83	11.56 ± 0.51	15.04 ± 0.40	11.49 ± 0.20	16.67 ± 0.20
	e_{cov}	5.83 ± 0.67	8.53 ± 0.81	12.09 ± 0.51	12.49 ± 0.43	13.65 ± 0.19	17.33 ± 0.16

The results reveal a consistent trade-off between fitting and coverage errors. Local PCA provides particularly small coverage errors, ranking second on five of the six manifolds, but has substantially larger fitting errors. Mean shift sometimes fits more closely, including the second-smallest torus fitting error, but has the largest coverage error on every manifold. SCMS-log-KDE improves markedly on SCMS-KDE and gives a more balanced fit, while YY-adapted has the second-smallest fitting error on four manifolds at the cost of larger coverage errors, especially on the trefoil knot, sphere, and torus. MF25 is also competitive, attaining the second-smallest trefoil fitting error and coverage errors on the sphere and torus close to those of STSP. This broadly similar performance is consistent with MF25 and STSP targeting related manifold-fitting objects, both defined as zero sets of local normal residuals, although their constructions and numerical procedures differ. Across all six manifolds, STSP reduces fitting error by 17–46% relative to local PCA and coverage error by 44–66% relative to mean shift, thereby performing well on both criteria. MS-STSP has the smallest fitting and coverage errors in every setting, reducing the corresponding STSP errors by 9–42%. For example, on the torus its fitting and coverage errors are 0.01667 and 0.01733, compared with 0.02401 and 0.02377 for STSP.

Figure 3 uses the torus as a representative example. The local-PCA fit in panel (a) covers the surface but has a visibly rough, spiked boundary. The mean-shift and YY-adapted fits in panels (b) and (e) contain clear gaps, consistent with their larger coverage errors. SCMS-KDE and MF25 in panels (c) and (f) recover smoother boundaries but retain scattered points away from the surface. SCMS-log-KDE and STSP in panels (d) and (g) give coherent, well-covered surfaces with only a small number of such points. MS-STSP in panel (h) gives the cleanest reconstruction, with a smooth boundary, few visible outliers, and even coverage. These comparisons are consistent with the intended effects of successive normal correction and its multi-scale extension. Corresponding fitted-cloud comparisons for the other manifolds are provided in Supplementary Figures S4–S9.




Figure 3: Torus fits for the first of ten independent realisations. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.

5.3 Multi-scale fitting under high noise

To assess robustness beyond the baseline noise level, Table 3 tracks the two errors on the circle as σ increases. Both errors increase for every method, but at markedly different rates. Local PCA continues to favour coverage, attaining the smallest coverage error at $\sigma = 0.3$ while its fitting error becomes large; mean shift shows the converse pattern, with the second-smallest fitting error at $\sigma = 0.2$ and 0.3 but substantially larger coverage errors. SCMS-log-KDE remains more stable than SCMS-KDE, while YY-adapted preserves coverage better than fitting and MF25 deteriorates sharply at the highest noise level. STSP remains competitive on fitting but loses coverage as noise increases. MS-STSP provides the most consistent balance: it has the smallest fitting error at all three noise levels and the smallest or second-smallest coverage error. At $\sigma = 0.3$, its fitting and coverage errors are 0.03323 and 0.02706, reductions of approximately 57% and 60% from the corresponding STSP errors of 0.07739 and 0.06815.

Table 3: Fitting and coverage errors over ten independent realisations of the high-noise circle experiment (mean $\pm$ standard deviation, in units of 10^{-3}). Within each noise level and for each error measure separately, the smallest mean is marked in bold and the second smallest is underlined. Rankings use unrounded means.

Method	Error	$\sigma = 0.1$	$\sigma = 0.2$	$\sigma = 0.3$
Local PCA	e_{fit}	26.27 ± 0.76	78.86 ± 3.16	150.48 ± 5.16
	e_{cov}	<u>10.27 ± 0.46</u>	<u>18.74 ± 0.82</u>	26.51 ± 1.28
Mean shift	e_{fit}	16.94 ± 6.26	<u>37.22 ± 5.81</u>	<u>72.96 ± 9.49</u>
	e_{cov}	26.92 ± 2.38	46.24 ± 2.75	77.62 ± 13.68
SCMS-KDE	e_{fit}	38.13 ± 1.08	125.19 ± 5.35	230.17 ± 10.26
	e_{cov}	26.13 ± 1.10	63.66 ± 7.25	66.47 ± 12.18
SCMS-log-KDE	e_{fit}	14.22 ± 1.36	43.25 ± 3.32	91.09 ± 12.33
	e_{cov}	13.86 ± 1.39	39.34 ± 3.90	83.21 ± 11.31
YY-adapted	e_{fit}	<u>11.05 ± 1.39</u>	37.52 ± 4.44	96.51 ± 6.03
	e_{cov}	10.84 ± 1.20	26.51 ± 2.36	30.92 ± 1.39
MF25	e_{fit}	13.51 ± 1.31	43.50 ± 3.80	154.62 ± 13.85
	e_{cov}	13.27 ± 1.34	38.69 ± 3.65	78.32 ± 11.98
STSP	e_{fit}	13.34 ± 1.00	38.04 ± 3.18	77.39 ± 12.10
	e_{cov}	12.55 ± 1.22	33.05 ± 3.65	68.15 ± 10.72
MS-STSP	e_{fit}	11.03 ± 1.27	17.20 ± 2.35	33.23 ± 3.33
	e_{cov}	9.36 ± 1.15	14.58 ± 2.19	<u>27.06 ± 3.83</u>




Figure 4: Circle fitting for the first of ten independent realisations at $\sigma = 0.3$. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively. Blue points are fitted observations and the thin red curve is the underlying circle.

Figure 4 shows the corresponding fits at $\sigma = 0.3$; fits at the lower noise levels are provided in Supplementary Section S5. Local PCA and YY-adapted in panels (a) and (e) form rough, diffuse bands; the mean-shift fit in panel (b) is sparse and uneven along the circle; and SCMS-KDE and MF25 in panels (c) and (f) retain many points away from the underlying curve. More importantly, some apparently smooth fits lie systematically inside the red reference circle, most visibly for SCMS-log-KDE and MF25 in panels (d) and (f), illustrating curvature bias under high noise. STSP in panel (g) removes most outlying points but retains some irregularity and inward displacement. MS-STSP in panel (h) follows the reference circle more closely

while maintaining nearly complete coverage, consistent with multi-scale extrapolation reducing curvature bias without reintroducing substantial fitting variability.

5.4 Empirical convergence and radius sensitivity

We next examine agreement between the empirical behaviour of STSP and the theory in three respects: the scaling of fitting error with noise, geometric convergence of the iterates to an error floor, and sensitivity to the neighbourhood radius.

Figure 5 gives a hard-radius finite-sample diagnostic for the circle and sphere over a wide noise range. The dashed guide has the theoretical $O(\sigma^2)$ slope; its vertical position is chosen for visual comparison with the observed errors, rather than calculated from a theoretical constant.




Figure 5: STSP fitting error versus noise on log-log axes for (a) the circle and (b) the sphere. Points show means over ten independent repetitions; error bars show one standard deviation on either side of the mean. Dashed lines indicate $O(\sigma^2)$ scaling.

Figure 6 shows the first few iterations for the circle and sphere. The finite-sample bound in Theorem 3 has the form $\text{dist}(\hat{\mathbf{y}}^{(t)}, \mathcal{M}) \leq \kappa^t \text{dist}(\hat{\mathbf{y}}^{(0)}, \mathcal{M}) + C(\sigma^2 + hr_N)/(1 - \kappa)$: a geometrically decreasing term followed by a finite-sample error floor. The observed fitting errors fall rapidly over the first few iterations and then level off, qualitatively consistent with this form.

Figure 7 examines sensitivity to the neighbourhood radius and exhibits the expected bias-variance pattern. Too-small radii give unstable local moments, whereas too-large radii introduce curvature bias. The minimizing ratio changes with the noise level.

5.5 Dentate gyrus single-cell data

We analyse single-cell RNA-sequencing data from the mouse dentate gyrus (Hochgerner et al., 2018). The data are available at https://raw.githubusercontent.com/theislabs/scvelo_notebooks/master/data/DentateGyrus/10X43_1.h5ad, which contain 2,930 cells assigned to 14 populations. Five annotations describe granule-cell neurogenesis: radial glia-like cells, neuronal intermediate progenitor cells (nIPC), neuroblasts, immature granule cells, and mature granule cells. The remaining annotations distinguish




Figure 6: STSP fitting error versus iteration for (a) the circle and (b) the sphere. Solid curves show means over ten independent repetitions; shaded bands show one standard deviation on either side of the mean. Dashed curves schematically illustrate the geometric-decay-plus-floor form of Theorem 3.




Figure 7: STSP fitting error versus neighbourhood radius on a circle at two noise levels. Curves show means over ten independent repetitions; error bars show one standard deviation on either side of the mean. Stars mark minima of the mean errors.

astrocytes, oligodendrocyte precursor cells (OPC), oligodendrocytes (OL), microglia, endothelial cells, and the Cajal–Retzius, Cck–Tox, GABAergic, and mossy neuronal populations.

Counts are library-normalised, log-transformed, and restricted to the 2,000 most variable genes before coordinatewise standardisation and PCA. All methods use the same unwhitened representation at a given dimension. The scores are divided by their root-mean-square cloud radius. For each method, candidate parameters are selected by the average Euclidean silhouette score (SIL) of the 14 supplied annotations. The principal analysis uses $p = 30$ and $d = 1$; candidate sets and iteration settings are given in Supplementary Section S5.

Table 4: Results for the dentate gyrus data using 30 principal components. SIL uses the 14 supplied annotations. Accuracy uses optimal one-to-one matching of cluster and annotation labels. Clustering uses 14 clusters; entries are means over 10 shared k -means seeds, with five k -means replicates per seed. Larger values are better.

Method	Average SIL	ARI	Accuracy	NMI
Input	0.1615	0.4439	0.5747	0.6848
Mean shift	0.2504	0.4575	0.6514	0.7009
SCMS-KDE	0.3086	0.4701	0.6282	0.6332
SCMS-log-KDE	0.3841	0.4621	0.6174	0.7100
Local PCA	0.3643	0.4628	0.6230	0.7058
YY-adapted	0.3742	0.5153	0.6804	0.7227
MF25	0.3466	0.5061	0.6788	0.7302
STSP	0.4224	0.5048	0.6659	0.7275
MS-STSP	0.4300	0.5474	0.6905	0.7338

Table 4 shows that STSP raises average SIL from 0.1615 in the input and 0.2504 after mean shift to 0.4224. Its ARI, accuracy, and NMI also exceed those of the input, mean shift, local PCA, and both SCMS variants. Three-scale MS-STSP achieves the largest values on all four measures: SIL 0.4300, ARI 0.5474, accuracy 0.6905, and NMI 0.7338.

Figure 8 shows t-SNE plots computed separately from the STSP and MS-STSP fitted representations, coloured by the supplied cell annotations. In each plot, neuroblasts, immature granule cells, and mature granule cells occupy neighbouring regions, while the smaller radial glia-like and nIPC populations remain visible. Several glial, vascular, and other neuronal populations form separate labelled regions; OPC and OL lie near one another.

Figure 9 compares average SIL across $p = 20, 30, 40, 50$. At each dimension, parameters are selected separately for each method using the same full-cell annotation-based SIL criterion; the plot therefore shows selected performance at each p , not the behaviour of one fixed configuration across dimensions. The input SIL decreases as more components are retained. All fitted representations improve upon the input at each displayed dimension. MS-STSP has the highest SIL throughout. STSP remains above 0.40 at $p = 20, 30, 40$ and is second only to MS-STSP at all four dimensions; local PCA and SCMS-log-KDE also retain substantially higher SIL than the input as p increases.




Figure 8: Dentate gyrus cell populations in t-SNE plots of (a) the STSP-fitted and (b) the three-scale MS-STSP-fitted representations using 30 principal components. Colours indicate the 14 supplied annotations.




Figure 9: Average annotation-based silhouette versus the number of retained principal components for the dentate gyrus data.

6 Conclusion

STSP connects successive local tangent-space corrections to a fitted manifold through the geometry of its nearby population fixed points. Under the stated small-noise conditions, this fixed-point set is a C^1 normal graph over the underlying manifold, diffeomorphic to it and within $O(\sigma^2)$ in Hausdorff distance.

Population orbits initialised sufficiently close to this set converge geometrically to individual fixed points. With high probability, fixed-reference empirical orbits initialised within the local tube enter and remain in an $O(\sigma^2 + hr_N)$ neighbourhood of the underlying manifold after a geometric transient.

The experiments assess fitting accuracy and geometric coverage jointly; at higher noise levels, multi-scale aggregation reduces observed curvature-induced shrinkage. The theory establishes that stable aggregation preserves the single-scale localisation order. Further work could establish empirical-orbit convergence and joint guarantees for coverage and data-driven scale selection.

Conflict of interest

The authors declare no competing interests.

Acknowledgments

Zhigang Yao has received support from the Singapore Ministry of Education through Tier 1 grants A-8004146-00-00 and A-8002931-00-00 at the National University of Singapore. Yuqing Xia was supported by the National Natural Science Foundation of China (grant No. 12401500). The authors are grateful for the resources and facilities provided by the Shanghai Institute for Mathematics and Interdisciplinary Sciences (SIMIS), which were essential for the completion of this work.

Supplementary material

The Supplementary Material contains all proofs, supporting theoretical results, manifold definitions, and detailed parameter settings for the numerical experiments.

Supplementary Material: Proofs and Numerical Details

S1 Notation and geometric preliminaries

The supplement is organized by the three levels of theory in the main paper. Section S1 records notation and geometric estimates. Section S2 establishes the population fixed-point and orbit results for STSP, Section S3 gives the finite-sample results, and Section S4 develops the localization theory for MS-STSP. Numerical definitions and implementation details are collected in Section S5.

Notation follows the main paper. In particular, Π_x denotes the second fundamental form at x , and K_{geo} is the geometric derivative bound in Assumption 1(A2).

Fix $c > 0$ and set $h = c\sigma$, with $c\sigma \leq r_{\text{geo}}/2$. Population asymptotics are taken as $\sigma \rightarrow 0$; in Section S3, $N \rightarrow \infty$ jointly with $\sigma \rightarrow 0$. The normalised curvature remainder in the flat-law expansion is $O(c^2\sigma/r_{\text{geo}})$, which tends to zero for fixed c .

We first record two standard consequences of bounded curvature and positive reach, used repeatedly below.

Lemma S1 (Tangent approximation). *For $x \in \mathcal{M}$ and $z \in \mathcal{M}$ with $\|z - x\| \leq h \leq \tau/2$, write $z - x = \mathbf{P}_x(z - x) + \mathbf{n}(z)$ with $\mathbf{n}(z) \in N_x\mathcal{M}$. Then*

$$\|\mathbf{n}(z)\| \leq \frac{\|z - x\|^2}{2\tau}, \quad 0 \leq \|z - x\| - \|\mathbf{P}_x(z - x)\| \leq \frac{\|z - x\|^3}{4\tau^2}.$$

Proof. By Proposition 6.3 of Niyogi et al. (2008) and the assumption $\text{reach}(\mathcal{M}) \geq \tau$, $\text{dist}(z, x + T_x\mathcal{M}) \leq \|z - x\|^2/(2\tau)$ for $z \in \mathcal{M}$. Since $\text{dist}(z, x + T_x\mathcal{M}) = \|(I_D - \mathbf{P}_x)(z - x)\| = \|\mathbf{n}(z)\|$, this gives the first inequality. For the second inequality, set $r = \|z - x\|$, $r_T = \|\mathbf{P}_x(z - x)\|$, and $r_\perp = \|\mathbf{n}(z)\|$. Because the tangent and normal components are orthogonal, $r^2 = r_T^2 + r_\perp^2$, so $r_T \leq r$. If $r = 0$ the claim is trivial. Otherwise,

$$0 \leq r - r_T = \frac{r^2 - r_T^2}{r + r_T} = \frac{r_\perp^2}{r + r_T} \leq \frac{r_\perp^2}{r} \leq \frac{r^3}{4\tau^2},$$

where the last step uses $r_\perp \leq r^2/(2\tau)$ from the first inequality. This proves the stated bound. $\square$

Lemma S2 (Volume comparison). *There exist constants $c_0 \in (0, 1)$ and $C > 0$ such that, for $x \in \mathcal{M}$ and $h \leq c_0 r_{\text{geo}}$,*

$$B_g(x, h(1 - Ch^2)) \subset B_E(x, h) \cap \mathcal{M} \subset B_g(x, h(1 + Ch^2)).$$

Moreover,

$$\text{Area}\{B_E(x, h) \cap \mathcal{M}\} = \omega_d h^d (1 + O(h^2)),$$

where ω_d is the volume of the unit d -ball and the $O(h^2)$ term is uniform over $\mathcal{M}$ and controlled by the curvature bound K_{geo} in (A2).

Proof. Because $h \leq c_0 r_{\text{geo}} \leq \iota_{\mathcal{M}}$, the exponential map at x gives geodesic normal coordinates on the intrinsic balls considered below, while positive reach excludes distant intrinsic sheets from the corresponding Euclidean ball. A vector $u \in T_x\mathcal{M}$ in the injectivity ball represents $\exp_x(u)$ and $d_g(x, \exp_x(u)) = \|u\|$. The standard volume expansion in normal coordinates, valid under a uniform curvature bound (do Carmo, 1992, Chapter 3), says that the area element has density $1 + O(\|u\|^2)$ with respect to Lebesgue measure du on $T_x\mathcal{M}$. Hence

$$\text{Area}\{B_g(x, r)\} = \int_{\|u\| \leq r} \{1 + O(\|u\|^2)\} du = \omega_d r^d \{1 + O(r^2)\}.$$

We now compare the geodesic ball with the ambient Euclidean ball $B_E(x, h) \cap \mathcal{M}$. In the same exponential coordinates, the embedded geodesic has the Taylor expansion $\exp_x(u) - x = u + \frac{1}{2}\Pi_x(u, u) + O(\|u\|^3)$, where

Π_x is the second fundamental form and the quadratic term is normal to $T_x\mathcal{M}$. Hence $\|\exp_x(u) - x\|^2 = \|u\|^2 + O(\|u\|^4)$. Choose a constant $C_0 > 0$, independent of x and h , such that

$$\left| \frac{\|\exp_x(u) - x\|}{\|u\|} - 1 \right| \leq C_0 \|u\|^2 \quad (0 < \|u\| \leq r_{\text{geo}}/2).$$

Equivalently, whenever $\|u\| \leq r_{\text{geo}}/2$,

$$\|u\|(1 - C_0\|u\|^2) \leq \|\exp_x(u) - x\| \leq \|u\|(1 + C_0\|u\|^2).$$

Choose

$$c_0 \leq \min \left\{ \frac{1}{4}, \frac{1}{4r_{\text{geo}}\sqrt{C_0}} \right\}.$$

Set $C = 8C_0$. Then $h \leq c_0 r_{\text{geo}}$ implies $h \leq r_{\text{geo}}/4$, $Ch^2 \leq 1/2$, and $C_0(2h)^2 \leq 1/4 < 1/2$. To prove the first inclusion, take $z \in B_g(x, h(1 - Ch^2))$ and write $z = \exp_x(u)$. Since $d_g(x, z) = \|u\|$, we have $\|u\| \leq h(1 - Ch^2) \leq h \leq r_{\text{geo}}/4$, so the distance comparison above is applicable. Moreover,

$$\|z - x\| = \|\exp_x(u) - x\| \leq \|u\|(1 + C_0\|u\|^2) \leq h(1 - Ch^2)(1 + C_0h^2) \leq h,$$

where the last inequality uses $C \geq C_0$. Hence $z \in B_E(x, h) \cap \mathcal{M}$. Conversely, take $z \in B_E(x, h) \cap \mathcal{M}$ and put $r = \|z - x\|$. Proposition 6.3 of Niyogi et al. (2008), applied with $\text{reach}(\mathcal{M}) \geq \tau$ and $r \leq h \leq \tau/4$, gives the global comparison

$$d_g(x, z) \leq \tau\{1 - \sqrt{1 - 2r/\tau}\} \leq 2r \leq 2h.$$

Thus the reach condition excludes every intrinsically distant sheet from the Euclidean ball. Since $2h \leq r_{\text{geo}}/2 \leq \iota_{\mathcal{M}}$, there is now a unique u in the injectivity ball with $z = \exp_x(u)$ and $\|u\| = d_g(x, z) \leq 2h$. The local lower bound is therefore applicable and gives

$$\|u\| \leq \frac{h}{1 - C_0(2h)^2} \leq h\{1 + 2C_0(2h)^2\} \leq h(1 + Ch^2).$$

Since $d_g(x, \exp_x(u)) = \|u\|$, these two implications yield

$$B_g(x, h(1 - Ch^2)) \subset B_E(x, h) \cap \mathcal{M} \subset B_g(x, h(1 + Ch^2)).$$

Applying the preceding geodesic-ball volume expansion to the two bounding balls yields $\text{Area}\{B_E(x, h) \cap \mathcal{M}\} = \omega_d h^d \{1 + O(h^2)\}$, with constants uniform over the compact manifold. $\square$

S2 Population theory for STSP

S2.1 Flat reference model and uniform local expansions

Recall from Section 2.2 of the main paper that $\mathbf{Y}$ is a generic observation, $\boldsymbol{\mu}_h(\mathbf{y})$ and $\boldsymbol{\Sigma}_h(\mathbf{y})$ are its conditional local moments, $T_h(\mathbf{y})$ is the corresponding leading- d population local tangent space, and $\mathbf{P}_h(\mathbf{y})$ is its orthogonal projector. The population update is

$$\mathcal{T}_h(\mathbf{y}) = \boldsymbol{\mu}_h(\mathbf{y}) + \mathbf{P}_h(\mathbf{y})[\mathbf{y} - \boldsymbol{\mu}_h(\mathbf{y})]. \quad (\text{S1})$$

The fixed-point identity used in the main paper follows immediately from (S1): wherever the leading- d eigenspace is unique,

$$\mathcal{T}_h(\mathbf{y}) = \mathbf{y} \iff (I_D - \mathbf{P}_h(\mathbf{y}))\{\mathbf{y} - \boldsymbol{\mu}_h(\mathbf{y})\} = \mathbf{0}.$$

Use the tube Φ_ρ , projection point $x = \Pi_{\mathcal{M}}(\mathbf{y})$, and normal offset $\mathbf{v}(\mathbf{y}) = \mathbf{y} - x \in N_x \mathcal{M}$ defined in the main paper. When $h = c\sigma$, the required uniform conditional-moment expansion holds on the shrinking tube $\Phi_{A\sigma}$, where $A < \infty$ is fixed.

Use q , $W = (W_T, W_N)$, $p_{c,a}$, $\mathbb{E}_{c,a}$, $\text{cov}_{c,a}$, m_c , and S_c as defined in the main paper. For reference, the density is

$$p_{c,a}(w) = \frac{\mathbf{1}\{\|w\| \leq c\} \exp\{-\|a + w_N\|^2/2\}}{\int_{\|v\| \leq c} \exp\{-\|a + v_N\|^2/2\} dv}. \quad (\text{S2})$$

The corresponding normal map and full covariance are

$$m_c(a) = a + \mathbb{E}_{c,a} W_N, \quad S_c(a) = \text{cov}_{c,a}(W).$$

The law in (S2) is the exact scaled conditional law for an affine d -plane. We call (c, A) *spectrally admissible* when there is $g_0 > 0$ such that, for every $\|a\| \leq A$,

$$\lambda_{\min}\{\text{cov}_{c,a}(W_T)\} - \lambda_{\max}\{\text{cov}_{c,a}(W_N)\} \geq g_0. \quad (\text{S3})$$

This finite-dimensional integral condition is sufficient because the required normal contraction follows from the next lemma.

Lemma S3 (Automatic flat-model normal contraction). *For every fixed $c > 0$ and $A < \infty$, the flat-model normal mean satisfies*

$$\text{D}m_c(a) = I_q - S_N(c, a), \quad S_N(c, a) = \text{cov}_{c,a}(W_N).$$

Moreover, there is $\kappa_0 = \kappa_0(c, A, d, q) < 1$ such that

$$\sup_{\|a\| \leq A} \|\text{D}m_c(a)\|_{\text{op}} \leq \kappa_0.$$

Thus only the spectral separation in (S3) needs to be assumed or verified.

Proof. Differentiating the normalised density (S2) under the integral sign gives the score identity

$$\frac{\partial}{\partial a_j} \log p_{c,a}(w) = -\{w_{N,j} - \mathbb{E}_{c,a} W_{N,j}\}.$$

Consequently,

$$\frac{\partial}{\partial a_j} \mathbb{E}_{c,a} W_{N,i} = -\text{cov}_{c,a}(W_{N,i}, W_{N,j}),$$

which proves $\text{D}m_c(a) = I_q - S_N(c, a)$.

Integrating out the tangent coordinates shows that W_N has density, on the open ball $\|n\| < c$,

$$p_N(n) \propto (c^2 - \|n\|^2)^{d/2} \exp\{-\|a + n\|^2/2\} = \exp\{-V_a(n)\},$$

where

$$V_a(n) = \frac{1}{2}\|a + n\|^2 - \frac{d}{2} \log(c^2 - \|n\|^2).$$

Its Hessian is

$$\nabla^2 V_a(n) = I_q + \frac{d}{c^2 - \|n\|^2} I_q + \frac{2d}{(c^2 - \|n\|^2)^2} n n^\top \succeq \left(1 + \frac{d}{c^2}\right) I_q.$$

The Brascamp–Lieb covariance inequality therefore yields

$$0 \prec S_N(c, a) \preceq \mathbb{E}\{\nabla^2 V_a(W_N)^{-1}\} \preceq \frac{c^2}{c^2 + d} I_q \prec I_q.$$

The lower inequality is strict because p_N is positive on the full q -dimensional open ball. The matrix $S_N(c, a)$ depends continuously on a ; compactness of $\{\|a\| \leq A\}$ therefore gives

$$\underline{s}_{c,A} = \inf_{\|a\| \leq A} \lambda_{\min}\{S_N(c, a)\} > 0.$$

All eigenvalues of $Dm_c(a) = I_q - S_N(c, a)$ lie in $(0, 1)$, and hence

$$\|Dm_c(a)\|_{\text{op}} = 1 - \lambda_{\min}\{S_N(c, a)\} \leq 1 - \underline{s}_{c,A} =: \kappa_0 < 1.$$

This proves the uniform contraction. The covariance inequality on the open ball follows by the standard smooth convex-domain approximation in the Brascamp–Lieb theorem (Brascamp and Lieb, 1976). $\square$

Throughout Theorems S1–S3, $h = c\sigma$, (c, A) is spectrally admissible, and $c\sigma \leq r_{\text{geo}}/2$.

Lemma S4 (Scaled population moments near $\mathcal{M}$). *For $\mathbf{y} \in \Phi_{A\sigma}$, choose orthonormal tangent and normal frames at $x = \Pi_{\mathcal{M}}(\mathbf{y})$ and write $\mathbf{v}(\mathbf{y}) = \sigma a$ in the normal frame. Uniformly over this tube,*

$$\boldsymbol{\mu}_h(\mathbf{y}) = x + \sigma(\mathbf{0}_d, m_c(a)) + O(\sigma^2), \quad \boldsymbol{\Sigma}_h(\mathbf{y}) = \sigma^2 S_c(a) + O(\sigma^3),$$

where the displayed vectors and matrices are transported to the ambient frames. Under the eigengap condition in (S3),

$$\|\mathbf{P}_h(\mathbf{y}) - \mathbf{P}_x\|_{\text{op}} \leq C\sigma.$$

Consequently,

$$\mathbf{Q}_x\{\mathcal{T}_h(\mathbf{y}) - x\} = \sigma m_c(a) + O(\sigma^2), \quad \|\mathbf{P}_x\{\mathcal{T}_h(\mathbf{y}) - x\}\| \leq C\sigma^2.$$

Moreover, after writing $\mathbf{y} = x + \sigma a$ in any smooth local orthonormal frame of the normal bundle, the displayed expansions and the projector bound hold uniformly in the C^1 norm with respect to (x, a) on compact sub-tubes $\|a\| \leq A_{\text{sub}} < A$. The $O(D-d)$ -equivariance of m_c and S_c makes these local expansions consistent on overlapping normal-bundle charts.

Proof. Let p_σ be the ambient density of one observation. For $z = x + \sigma b$, the Gaussian-convolution formula and uniform sampling on $\mathcal{M}$ give

$$p_\sigma(z) = \frac{(2\pi\sigma^2)^{-D/2}}{\text{Area}(\mathcal{M})} \int_{\mathcal{M}} \exp\{-\|z - \xi\|^2/(2\sigma^2)\} dA(\xi).$$

Use a smooth normal-bundle chart at x and put $\xi = \exp_x(\sigma s)$. Uniformly for bounded b ,

$$\frac{\|x + \sigma b - \exp_x(\sigma s)\|^2}{\sigma^2} = \|b_T - s\|^2 + \|b_N\|^2 - \sigma \langle b_N, \Pi_x(s, s) \rangle + O\{\sigma^2(1 + \|s\|^6)\},$$

on a fixed chart, and $dA\{\exp_x(\sigma s)\} = \sigma^d\{1 + O(\sigma^2\|s\|^2)\} ds$. After restricting first to $\|s\| \leq C\sqrt{\log(1/\sigma)}$, Taylor expansion under the integral is dominated by a polynomial times a Gaussian. The remaining part of that chart, and the part of $\mathcal{M}$ outside it, are respectively Gaussian-tail and $O\{\exp(-c_{\text{tail}}/\sigma^2)\}$ terms for some $c_{\text{tail}} > 0$. The same domination holds after one derivative in b or in the chart centre and frame. Consequently, uniformly on every bounded bundle chart,

$$\sigma^{D-d} p_\sigma(x + \sigma b) = C_{\text{dens}} \exp(-\|b_N\|^2/2) + O_{C^1}(\sigma).$$

Here the leading integral over $s \in \mathbb{R}^d$ removes b_T , and $C_{\text{dens}} = (2\pi)^{-(D-d)/2}/\text{Area}(\mathcal{M})$.

Now write $\mathbf{y} = x + \sigma a$, $a \in N_x \mathcal{M}$, and $w = (\mathbf{Y} - \mathbf{y})/\sigma$. This query-centred change of variables turns every hard neighbourhood into the fixed ball $\{\|w\| \leq c\}$. Thus differentiation with respect to the query

changes the integrand but introduces no moving-boundary term. On this fixed ball, the preceding expansion is applicable with $b = a + w$ and shows that the unnormalized conditional density is

$$C_{\text{dens}} \exp\{-\|a + w_N\|^2/2\} + O_{C^1}(\sigma).$$

Its integral is uniformly bounded away from zero for $\|a\| \leq A$. Division by this integral therefore gives

$$p_{c,a}(w) + O_{C^1}(\sigma) \quad \text{on } \{\|w\| \leq c\}.$$

The leading law is exactly (S2); tangent reflection gives $\mathbb{E}_{c,a} W_T = \mathbf{0}_d$. Since $\mathbf{Y} = \mathbf{y} + \sigma w$, integration of w and $ww^\top$ now yields

$$\boldsymbol{\mu}_h(\mathbf{y}) = x + \sigma\{\mathbf{0}_d, a + \mathbb{E}_{c,a} W_N\} + O_{C^1}(\sigma^2), \quad \boldsymbol{\Sigma}_h(\mathbf{y}) = \sigma^2 S_c(a) + O_{C^1}(\sigma^3).$$

This proves both moment expansions, including their asserted uniform C^1 form on compact sub-tubes. By (S3), the normalised leading covariance $S_c(a)$ has eigengap at least g_0 . For small σ , the normalised population covariance has gap at least $g_0/2$. The spectral-projector map is C^1 with uniformly bounded derivative on matrices with this gap, so its standard resolvent formula (equivalently, the C^1 Davis–Kahan perturbation bound) gives $\|\mathbf{P}_h(\mathbf{y}) - \mathbf{P}_x\|_{C^1} \leq C\sigma$.

Finally, the leading term of $\mathbf{y} - \boldsymbol{\mu}_h(\mathbf{y})$ is normal and has size $O(\sigma)$. Substitution in (S1) shows that the $O(\sigma)$ projector error contributes only $O(\sigma^2)$. Projection onto $N_x \mathcal{M}$ and $T_x \mathcal{M}$ gives the two stated consequences. $\square$

Lemma S5 (Flat-model eigengap criterion). *For the law (S2), define*

$$\alpha_T(c, \|a\|) = d^{-1} \text{tr}\{\text{cov}_{c,a}(W_T)\}.$$

Then

$$S_c(a) = \begin{pmatrix} \alpha_T(c, \|a\|) I_d & 0 \\ 0 & S_N(c, a) \end{pmatrix}.$$

Hence the leading d eigenspace is tangent uniformly on $\|a\| \leq A$ if and only if

$$\inf_{\|a\| \leq A} \{\alpha_T(c, \|a\|) - \lambda_{\max}(S_N(c, a))\} > 0.$$

This is exactly the spectral admissibility condition used above; normal contraction is automatic by Lemma S3. Both covariance terms are continuous and are given by bounded integrals over the fixed ball $\{\|w\| \leq c\}$.

Proof. Reflection in each tangent coordinate makes every tangent–normal cross-moment zero, and rotation invariance in the tangent coordinates makes the tangent block a scalar multiple of I_d . The spectral statement follows. Continuity follows by differentiation under the integral sign on the bounded domain. $\square$

Remark S1 (Eigenspace separation). Conditioning on the neighbourhood event couples signal and noise, so the conditional eigengap is determined by the limiting integrals in Lemma S5, evaluated for the chosen d, D, c and A . As $c \downarrow 0$ the conditioned law approaches the uniform law on a small ball, the two covariance blocks become isotropic, and their gap vanishes at $O(c^4)$. The localization results assume spectral admissibility at every constituent scale.

For each row, the reported quantity is

$$\inf_{\|a\| \leq 3} [\lambda_{\min}\{\text{cov}_{c,a}(W_T)\} - \lambda_{\max}\{\text{cov}_{c,a}(W_N)\}].$$

The positive values provide a numerical check of spectral separation for these dimension pairs and radius ratios. The theoretical results assume spectral admissibility on $\Phi_{3\sigma}$.

Remark S2 (Effective expansion parameter). The normalised remainder has effective size $c^2\sigma/r_{\text{geo}}$, as is visible from the curvature term in the preceding exponent. Thus a large calibrated c requires $c^2\sigma \ll r_{\text{geo}}$. Under the uniform sampling condition, the leading local law has no additional tangential density-score term.

Table S1: Numerically estimated minimum normalised spectral gaps for the flat reference law at normalised tube width $A = 3$.

$c = h/\sigma$	$d = 1, q = 2$	$d = 2, q = 1$
1.0	0.0207	0.0267
1.25	0.0448	0.0628
1.5	0.0824	0.1243
2.0	0.2106	0.3496
2.5	0.4331	0.6030
3.0	0.7855	0.9267
3.5	1.3115	1.3578
4.0	2.0575	1.9269
5.0	4.3429	3.5847
6.0	7.6804	5.9915
8.0	16.8168	12.7253

S2.2 Population fixed-point manifold and attraction

Theorem S1 (Local normal-distance recursion). *Under Assumption 1 of the main paper and the spectral admissibility condition above, there are $\sigma_{\text{con}} > 0$, $C < \infty$, and $\kappa \in (\kappa_0, 1)$ such that, for $\sigma \leq \sigma_{\text{con}}$ and $\mathbf{y} \in \Phi_{A\sigma}$,*

$$\text{dist}(\mathcal{T}_h(\mathbf{y}), \mathcal{M}) \leq \kappa \text{dist}(\mathbf{y}, \mathcal{M}) + C\sigma^2.$$

For sufficiently small σ , $\Phi_{A\sigma}$ is forward invariant, and any population orbit starting there obeys

$$\text{dist}(\mathcal{T}_h^{(t)}(\mathbf{y}), \mathcal{M}) \leq \kappa^t \text{dist}(\mathbf{y}, \mathcal{M}) + \frac{C\sigma^2}{1 - \kappa}.$$

This result controls normal localization; complete ambient convergence and fixed-point existence are established below.

Proof. With the notation of Lemma S4, $\mathbf{y} - x = \sigma(\mathbf{0}_d, a)$ and

$$\mathcal{T}_h(\mathbf{y}) - x = \sigma(\mathbf{0}_d, m_c(a)) + O(\sigma^2).$$

Indeed, the leading displacement $\mathbf{y} - \mu_h(\mathbf{y})$ is normal, the leading projection is tangent, and multiplying its $O(\sigma)$ projection error by an $O(\sigma)$ displacement contributes only $O(\sigma^2)$. Put $\mathbf{v}_0 = x + \sigma(\mathbf{0}_d, m_c(a))$. It lies on the normal fibre inside the reach tube, so $\text{dist}(\mathbf{v}_0, \mathcal{M}) = \sigma\|m_c(a)\|$. Since distance to a closed set is one-Lipschitz and $\|\mathcal{T}_h(\mathbf{y}) - \mathbf{v}_0\| = O(\sigma^2)$, we have

$$\text{dist}(\mathcal{T}_h(\mathbf{y}), \mathcal{M}) \leq \sigma\|m_c(a)\| + C\sigma^2.$$

Tangent and normal reflection symmetry gives $m_c(\mathbf{0}_q) = \mathbf{0}_q$; the mean-value theorem and Lemma S3 give $\|m_c(a)\| \leq \kappa_0\|a\|$. The preceding displays give $\text{dist}(\mathcal{T}_h(\mathbf{y}), \mathcal{M}) \leq \kappa_0\text{dist}(\mathbf{y}, \mathcal{M}) + C\sigma^2$, which immediately implies the stated inequality for every $\kappa > \kappa_0$. Iteration yields the affine recursion. Forward invariance follows from $\kappa A\sigma + C\sigma^2 \leq A\sigma$ after decreasing σ_{con} . $\square$

Ambient convergence additionally requires control of tangential motion. We obtain it by identifying the complete nearby zero set of the update residual and controlling the map transverse to that set. Recall the population residual $\mathbf{F}_h$ and its zero set $\mathcal{S}_{h,\text{pop}}$ from Section 2.2 of the main paper.

Lemma S6 (Fixed-point manifold and transverse contraction). *Fix $0 < A_0 < A$. Under the conditions of Theorem S1, there are $\sigma_{\text{fix}}, \rho_0, C, C_{\text{step}} > 0$ and $\bar{\kappa} \in (0, 1)$ such that, for $\sigma \leq \sigma_{\text{fix}}$:*

(i) $\mathcal{M}_h := \mathcal{S}_{h, \text{pop}} \cap \Phi_{A_0 \sigma}$ is a compact C^1 d -dimensional normal graph

$$\mathcal{M}_h = \{x + b_h(x) : x \in \mathcal{M}, b_h(x) \in N_x \mathcal{M}\}, \quad \|b_h\|_{C^1} \leq C\sigma^2,$$

and $\Pi_{\mathcal{M}} : \mathcal{M}_h \rightarrow \mathcal{M}$ is a C^1 diffeomorphism;

(ii)

$$d_H(\mathcal{M}_h, \mathcal{M}) \leq C\sigma^2;$$

(iii) on $\mathcal{U}_h = \{\mathbf{y} : \text{dist}(\mathbf{y}, \mathcal{M}_h) < \rho_0 \sigma\}$ contained in $\Phi_{A\sigma}$,

$$\text{dist}(\mathcal{T}_h(\mathbf{y}), \mathcal{M}_h) \leq \bar{\kappa} \text{dist}(\mathbf{y}, \mathcal{M}_h), \quad \|\mathcal{T}_h(\mathbf{y}) - \mathbf{y}\| \leq C_{\text{step}} \text{dist}(\mathbf{y}, \mathcal{M}_h).$$

Proof. Work first in one smooth normal-bundle chart and write $\mathbf{y} = x + \sigma a$, with $x \in \mathcal{M}$ and $\|a\| \leq A_0$. The residual $\mathbf{F}_h(\mathbf{y})$ lies in the estimated normal space $T_h(\mathbf{y})^\perp$. By Lemma S4, this estimated normal space makes angle $O(\sigma)$ with $N_x \mathcal{M}$; hence orthogonal projection from it onto $N_x \mathcal{M}$ is invertible for small σ . Use this projection to identify the residual with a vector in $N_x \mathcal{M}$ and define its scaled coordinate representation $\mathbf{H}_{\sigma, x}(a)$. More explicitly, using $\mathbf{Q}_x = I_D - \mathbf{P}_x$, set

$$\mathbf{H}_{\sigma, x}(a) = \sigma^{-1} \mathbf{Q}_x \mathbf{F}_h(x + \sigma a).$$

The restriction of $\mathbf{Q}_x$ to $T_h(x + \sigma a)^\perp$ is one-to-one, so $\mathbf{H}_{\sigma, x}(a) = \mathbf{0}$ is equivalent to $\mathbf{F}_h(x + \sigma a) = \mathbf{0}$. The uniform C^1 expansion in Lemma S4 gives

$$\mathbf{H}_{\sigma, x}(a) = a - m_c(a) + \mathbf{r}_\sigma(x, a), \quad \|\mathbf{r}_\sigma\|_{C^1} \leq C\sigma.$$

In particular,

$$\mathbf{D}_a \mathbf{H}_{\sigma, x}(a) = I_q - \mathbf{D}m_c(a) + O(\sigma) = \text{cov}_{c, a}(W_N) + O(\sigma).$$

The smallest eigenvalue of $\text{cov}_{c, a}(W_N)$ is bounded away from zero uniformly on $\|a\| \leq A_0$; hence the normal Jacobian is uniformly invertible for sufficiently small σ . Thus $\mathbf{F}_h(x + \sigma a) = \mathbf{0}$ is equivalent to

$$a = m_c(a) - \mathbf{r}_\sigma(x, a).$$

The right-hand side maps the closed A_0 -ball into itself and is a contraction in a , uniformly in x , because $\|\mathbf{D}m_c(a)\|_{\text{op}} \leq \kappa_0 < 1$ and $\|\mathbf{D}_a \mathbf{r}_\sigma\| \leq C\sigma$; on the boundary its norm is at most $\kappa_0 A_0 + C\sigma < A_0$. It has a unique solution $a_h(x)$ for every x , and $a_h(x) = O(\sigma)$ because $m_c(\mathbf{0}_q) = \mathbf{0}_q$ and $\mathbf{r}_\sigma(x, \mathbf{0}_q) = O(\sigma)$. The parameter-dependent contraction theorem, equivalently the implicit-function theorem applied to $\mathbf{H}_{\sigma, x}$, makes a_h a C^1 section of the normal bundle, with $\|a_h\|_{C^1} \leq C\sigma$. On overlaps uniqueness makes these local sections agree. Set $b_h(x) = \sigma a_h(x)$. Consequently

$$\mathcal{M}_h = \{x + b_h(x) : x \in \mathcal{M}\}, \quad \|b_h\|_{C^1} \leq C\sigma^2,$$

is a compact C^1 manifold of fixed points, its nearest-point projection onto $\mathcal{M}$ is a diffeomorphism, and its graph representation gives $d_H(\mathcal{M}_h, \mathcal{M}) \leq C\sigma^2$. The same contraction argument excludes any other residual zero in $\Phi_{A_0 \sigma}$. Choose $\rho_0 < (A - A_0)/2$. After decreasing σ_{fix} so that the graph displacement is smaller than $(A - A_0)\sigma/2$, the resulting attraction neighbourhood $\mathcal{U}_h$ is contained in $\Phi_{A\sigma}$.

It remains to control points off $\mathcal{M}_h$. In bundle coordinates centred at the graph $a_h(x)$, write

$$\mathcal{T}_h(x + \sigma a) = X_\sigma(x, a) + \sigma A_\sigma(x, a), \quad A_\sigma(x, a) \in N_{X_\sigma(x, a)} \mathcal{M}.$$

The C^1 moment and projector expansions, together with the fact that the leading displacement is normal, give

$$X_\sigma(x, a) = x + O_{C^1}(\sigma^2), \quad A_\sigma(x, a) = m_c(a) + O_{C^1}(\sigma).$$

Because the graph consists of pointwise fixed points,

$$X_\sigma\{x, a_h(x)\} = x, \quad A_\sigma\{x, a_h(x)\} = a_h(x).$$

For the transverse coordinate $\Delta = a - a_h(x)$, the mean-value theorem therefore gives

$$A_\sigma(x, a) - a_h\{X_\sigma(x, a)\} = [Dm_c\{a_h(x)\} + O(\sigma) + O(\|\Delta\|)] \Delta.$$

The term from differentiating $a_h\{X_\sigma(x, a)\}$ is included in $O(\sigma)$ because $\|a_h\|_{C^1} = O(\sigma)$ and $D_a X_\sigma = O(\sigma^2)$. Choose $\kappa_0 < \kappa_1 < \bar{\kappa} < 1$. If C_* bounds the remainder derivatives and coordinate distortion, choose $C_*(\sigma_{\text{fix}} + \rho_0) \leq \frac{1}{2} \min\{\kappa_1 - \kappa_0, \bar{\kappa} - \kappa_1\}$. Then the bracket has norm at most κ_1 ; in particular, the attraction radius deteriorates as $\kappa_0 \uparrow 1$. Uniformly in this tube,

$$\text{dist}(x + \sigma a, \mathcal{M}_h) = \sigma \|\Delta\| \{1 + O(\sigma) + O(\|\Delta\|)\}.$$

Shrinking σ_{fix} and ρ_0 once more absorbs this coordinate distortion and gives the first inequality in (iii). Finally, $\mathbf{F}_h$ is C^1 , vanishes on $\mathcal{M}_h$, and has uniformly bounded derivative there. Another application of the mean-value theorem gives

$$\|\mathcal{T}_h(\mathbf{y}) - \mathbf{y}\| = \|\mathbf{F}_h(\mathbf{y})\| \leq C_{\text{step}} \text{dist}(\mathbf{y}, \mathcal{M}_h),$$

which completes the proof. $\square$

Theorem S2 (Population fixed-point manifold and local orbit convergence). *Under the conditions of Lemma S6, the fixed-point set in $\Phi_{A_0\sigma}$ is the normal graph $\mathcal{M}_h$ described there, with $\|b_h\|_{C^1} \leq C\sigma^2$ and $d_H(\mathcal{M}_h, \mathcal{M}) \leq C\sigma^2$. Every $\mathbf{y}^{(0)} \in \mathcal{U}_h$ generates an orbit $\mathbf{y}^{(t+1)} = \mathcal{T}_h(\mathbf{y}^{(t)})$ that converges to a fixed point $\mathbf{y}^{(\infty)} \in \mathcal{M}_h$. More precisely,*

$$\text{dist}(\mathbf{y}^{(t)}, \mathcal{M}_h) \leq \bar{\kappa}^t \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_h)$$

and

$$\|\mathbf{y}^{(t)} - \mathbf{y}^{(\infty)}\| \leq \frac{C_{\text{step}} \bar{\kappa}^t}{1 - \bar{\kappa}} \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_h).$$

In particular, $\text{dist}(\mathbf{y}^{(\infty)}, \mathcal{M}) \leq C\sigma^2$.

Proof. Lemma S6(iii) makes $\mathcal{U}_h$ forward invariant and gives the first display by induction. It also gives

$$\|\mathbf{y}^{(t+1)} - \mathbf{y}^{(t)}\| \leq C_{\text{step}} \bar{\kappa}^t \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_h).$$

The right-hand side is summable. Hence the orbit is Cauchy and converges in $\mathbb{R}^D$ to a point $\mathbf{y}^{(\infty)}$; summing the tail from t onward gives the stated error bound. Since the distance to the closed set $\mathcal{M}_h$ tends to zero, $\mathbf{y}^{(\infty)} \in \mathcal{M}_h$, and every point of $\mathcal{M}_h$ is fixed. Part (ii) of the lemma yields the final distance bound. $\square$

S3 Finite-sample theory for STSP

S3.1 Empirical fixed points and orbit localization

We analyse the empirical operator $\widehat{\mathcal{T}}_{\mathcal{Z},h}$ constructed from the fixed i.i.d. reference sample $\mathcal{Z}$. For reference, the empirical neighbourhood and local moments are

$$\mathcal{I}_{\mathcal{Z},h}(\mathbf{y}) = \{j : \|\mathbf{z}_j - \mathbf{y}\| \leq h\}, \quad n_{\mathcal{Z},h}(\mathbf{y}) = |\mathcal{I}_{\mathcal{Z},h}(\mathbf{y})|,$$

and, whenever $n_{\mathcal{Z},h}(\mathbf{y}) > 0$,

$$\begin{aligned} \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y}) &= \frac{1}{n_{\mathcal{Z},h}(\mathbf{y})} \sum_{j \in \mathcal{I}_{\mathcal{Z},h}(\mathbf{y})} \mathbf{z}_j, \\ \widehat{\boldsymbol{\Sigma}}_{\mathcal{Z},h}(\mathbf{y}) &= \frac{1}{n_{\mathcal{Z},h}(\mathbf{y})} \sum_{j \in \mathcal{I}_{\mathcal{Z},h}(\mathbf{y})} \{\mathbf{z}_j - \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y})\} \{\mathbf{z}_j - \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y})\}^\top. \end{aligned}$$

Let $\widehat{\mathbf{P}}_{\mathcal{Z},h}(\mathbf{y})$ be its leading- d spectral projector whenever the d th and $(d+1)$ st eigenvalues are strictly separated, and define

$$\widehat{\mathcal{T}}_{\mathcal{Z},h}(\mathbf{y}) = \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y}) + \widehat{\mathbf{P}}_{\mathcal{Z},h}(\mathbf{y})\{\mathbf{y} - \widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y})\}.$$

Lemma S7 (Uniform empirical approximation). *Under the conditions of Theorem S1, let $0 < \delta < 1$ and*

$$r_N = \sqrt{\frac{D \log(1/h) + \log(1/\delta)}{Nh^d}} + \frac{D \log(1/h) + \log(1/\delta)}{Nh^d}.$$

There are constants $C, r_0, c_{\text{gap}} > 0$ such that, if $r_N \leq r_0$, then with probability at least $1 - \delta$, every empirical neighbourhood in $\Phi_{A\sigma}$ is nonempty and

$$\begin{aligned} \sup_{\mathbf{y} \in \Phi_{A\sigma}} \|\widehat{\boldsymbol{\mu}}_{\mathcal{Z},h}(\mathbf{y}) - \boldsymbol{\mu}_h(\mathbf{y})\| &\leq Chr_N, \\ \sup_{\mathbf{y} \in \Phi_{A\sigma}} \|\widehat{\boldsymbol{\Sigma}}_{\mathcal{Z},h}(\mathbf{y}) - \boldsymbol{\Sigma}_h(\mathbf{y})\|_{\text{op}} &\leq Ch^2 r_N, \\ \sup_{\mathbf{y} \in \Phi_{A\sigma}} \|\widehat{\mathbf{P}}_{\mathcal{Z},h}(\mathbf{y}) - \mathbf{P}_h(\mathbf{y})\|_{\text{op}} &\leq Cr_N, \quad \inf_{\mathbf{y} \in \Phi_{A\sigma}} \{\widehat{\lambda}_d(\mathbf{y}) - \widehat{\lambda}_{d+1}(\mathbf{y})\} \geq c_{\text{gap}} h^2. \end{aligned}$$

On the same event,

$$\sup_{\mathbf{y} \in \Phi_{A\sigma}} \|\widehat{\mathcal{T}}_{\mathcal{Z},h}(\mathbf{y}) - \mathcal{T}_h(\mathbf{y})\| \leq Chr_N.$$

Proof. The ambient-density expansion in Lemma S4 is uniform for scaled normal and tangential coordinates in bounded sets. Hence, for $\mathbf{y} \in \Phi_{A\sigma}$ and $\mathbf{z} \in B_E(\mathbf{y}, h)$, it gives $p_\sigma(\mathbf{z}) \geq c_1 \sigma^{-(D-d)}$ for all sufficiently small σ . Therefore

$$\inf_{\mathbf{y} \in \Phi_{A\sigma}} \Pr\{\mathbf{Y} \in B_E(\mathbf{y}, h)\} \geq c_1 \sigma^{-(D-d)} \omega_D h^D = c_{\text{mass}} h^d,$$

where $h = c\sigma$ and $c_{\text{mass}} > 0$ depends only on the fixed parameters. The same local density expansion gives the upper bound $\sup_{\mathbf{y} \in \Phi_{A\sigma}} p_h(\mathbf{y}) \leq C_{\text{mass}} h^d$ for the neighbourhood mass p_h defined in the main paper.

Write $\mathbf{v}_y(\mathbf{z}) = (\mathbf{z} - \mathbf{y})/h$ and $I_y(\mathbf{z}) = \mathbf{1}_{\{\|\mathbf{z} - \mathbf{y}\| \leq h\}}$. For $k, \ell = 1, \dots, D$, consider the classes consisting of I_y , $v_{y,k} I_y$, and $v_{y,k} v_{y,\ell} I_y$ as $\mathbf{y}$ ranges over $\Phi_{A\sigma}$. Each has envelope one, and every member has second moment at most $C_{\text{mass}} h^d$. The ball class is VC, and the coordinate-weighted classes are bounded semialgebraic VC-subgraph classes with complexity depending only on D . In particular, each class has a uniform $L_2(Q)$ covering bound of the form $N(\varepsilon, \mathcal{F}, L_2(Q)) \leq (A_D/\varepsilon)^{V_D}$ for every probability measure Q , with finite constants A_D, V_D depending only on the fixed ambient dimension. Applying the VC maximal inequality of

van der Vaart and Wellner (1996) to these bounded classes, followed by the bounded empirical-process tail inequality of Bousquet (2002), uses their variance bound $C_{\text{mass}}h^d$ and yields, simultaneously for the finitely many component classes, that

$$\sup_{\mathbf{y} \in \Phi_{A\sigma}} \left| \widehat{Q}_j(\mathbf{y}) - Q_j(\mathbf{y}) \right| \leq C \left\{ \sqrt{\frac{h^d u}{N}} + \frac{u}{N} \right\}, \quad u = D \log(1/h) + \log(1/\delta),$$

where Q_j denotes any scalar component of the three unnormalized moments and $\widehat{Q}_j$ its empirical counterpart. At the $h^{d/2}$ variance scale, the covering bound contributes a constant multiple of $\log(1/h)$; because D is fixed, its dimension-dependent factor and the finite number of components are absorbed into C in the displayed bound with $u = D \log(1/h) + \log(1/\delta)$. In particular, $Q_0 = p_h \geq c_{\text{mass}}h^d$, and, when r_N is small enough, $\widehat{Q}_0 \geq c_{\text{mass}}h^d/2$. Dividing the first- and second-moment bounds by these denominators and using $\|Q_1\|, \|Q_2\| \leq Q_0$ gives

$$\sup_{\Phi_{A\sigma}} \left\| \frac{\widehat{Q}_1}{\widehat{Q}_0} - \frac{Q_1}{Q_0} \right\| \leq Cr_N, \quad \sup_{\Phi_{A\sigma}} \left\| \frac{\widehat{Q}_2}{\widehat{Q}_0} - \frac{Q_2}{Q_0} \right\|_{\text{op}} \leq Cr_N.$$

Here Q_1 and Q_2 denote the vector and matrix unnormalized moments, respectively. Rescaling them by h and h^2 gives the conditional query-centred moment errors. To make the covariance conversion explicit, define the empirical query-centred conditional moments

$$\widehat{m}_1(\mathbf{y}) = n_{\mathcal{Z},h}(\mathbf{y})^{-1} \sum_{j \in \mathcal{I}_{\mathcal{Z},h}(\mathbf{y})} (\mathbf{z}_j - \mathbf{y}), \quad \widehat{M}_2(\mathbf{y}) = n_{\mathcal{Z},h}(\mathbf{y})^{-1} \sum_{j \in \mathcal{I}_{\mathcal{Z},h}(\mathbf{y})} (\mathbf{z}_j - \mathbf{y})(\mathbf{z}_j - \mathbf{y})^\top.$$

Then, exactly,

$$\widehat{\mu}_{\mathcal{Z},h}(\mathbf{y}) = \mathbf{y} + \widehat{m}_1(\mathbf{y}), \quad \widehat{\Sigma}_{\mathcal{Z},h}(\mathbf{y}) = \widehat{M}_2(\mathbf{y}) - \widehat{m}_1(\mathbf{y})\widehat{m}_1(\mathbf{y})^\top,$$

with the same identities for the population conditional moments. Since $\|\widehat{m}_1\|$ and its population counterpart are at most h , the uniform query-centred first- and second-moment bounds give the displayed centred covariance bound. The factors h and h^2 arise because every observation in a ball is within h of its query. The count bound also makes every empirical count at least a fixed multiple of Nh^d .

The population gap is at least $g_0\sigma^2/2 = g_0h^2/(2c^2)$. Decrease r_0 so that the covariance perturbation is at most $g_0h^2/(8c^2)$. Weyl's inequality, applied to both eigenvalues defining the gap, gives the empirical gap with $c_{\text{gap}} = g_0/(4c^2)$, and Davis–Kahan gives $\sup \|\widehat{\mathbf{P}}_{\mathcal{Z},h} - \mathbf{P}_h\|_{\text{op}} \leq Cr_N$. Since $\|\mathbf{y} - \mu_h(\mathbf{y})\| = O(h)$ on the shrinking tube,

$$\sup_{\mathbf{y} \in \Phi_{A\sigma}} \|\widehat{\mathcal{T}}_{\mathcal{Z},h}(\mathbf{y}) - \mathcal{T}_h(\mathbf{y})\| \leq Chr_N.$$

This proves the lemma. □

Corollary S1 (Localization of empirical fixed points). *Under the conditions of Lemma S7, there are constants $C, r_{\text{fp}} > 0$ such that, if $r_N \leq r_{\text{fp}}$, then with probability at least $1 - \delta$, every fixed point $\widehat{\mathbf{y}}_N \in \Phi_{A\sigma}$ of $\widehat{\mathcal{T}}_{\mathcal{Z},h}$, whenever one exists, satisfies*

$$\text{dist}(\widehat{\mathbf{y}}_N, \mathcal{M}) \leq C\{\sigma^2 + hr_N\}.$$

Consequently, for fixed d, D, c, A , $h = c\sigma$ and

$$N \geq C\sigma^{-(d+2)}\{D \log(1/\sigma) + \log(1/\delta)\}$$

suffice for the right-hand side to be $O(\sigma^2)$.

Proof. On the event in Lemma S7, if $\widehat{\mathbf{y}}_N = \widehat{\mathcal{T}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}_N)$, then

$$\begin{aligned} \text{dist}(\widehat{\mathbf{y}}_N, \mathcal{M}) &\leq \text{dist}(\mathcal{T}_h(\widehat{\mathbf{y}}_N), \mathcal{M}) + \|\widehat{\mathcal{T}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}_N) - \mathcal{T}_h(\widehat{\mathbf{y}}_N)\| \\ &\leq \kappa \text{dist}(\widehat{\mathbf{y}}_N, \mathcal{M}) + C\sigma^2 + Chr_N. \end{aligned}$$

Division by $1 - \kappa$ proves the assertion. The sample-size statement follows by requiring $r_N = O(\sigma)$. $\square$

Theorem S3 (Finite-sample orbit localization). *For the same fixed $0 < A_0 < A$, under the conditions of Corollary S1, there are $\sigma_{\text{orb}}, r_{\text{orb}}, C > 0$ such that, if $\sigma \leq \sigma_{\text{orb}}$ and $r_N \leq r_{\text{orb}}$, then with probability at least $1 - \delta$, every empirical orbit*

$$\widehat{\mathbf{y}}^{(t+1)} = \widehat{\mathcal{T}}_{\mathcal{Z},h}(\widehat{\mathbf{y}}^{(t)}), \quad \widehat{\mathbf{y}}^{(0)} \in \Phi_{A_0\sigma},$$

remains in $\Phi_{A\sigma}$ and satisfies, for every $t \geq 0$,

$$\text{dist}(\widehat{\mathbf{y}}^{(t)}, \mathcal{M}) \leq \kappa^t \text{dist}(\widehat{\mathbf{y}}^{(0)}, \mathcal{M}) + \frac{C(\sigma^2 + hr_N)}{1 - \kappa}.$$

In particular, after the geometric transient the entire empirical orbit lies in an $O(\sigma^2 + hr_N)$ neighbourhood of $\mathcal{M}$; if $r_N = O(\sigma)$ and $h = c\sigma$, this neighbourhood has width $O(\sigma^2)$.

Proof. On the uniform event in Lemma S7, for every $\mathbf{y} \in \Phi_{A\sigma}$,

$$\begin{aligned} \text{dist}(\widehat{\mathcal{T}}_{\mathcal{Z},h}(\mathbf{y}), \mathcal{M}) &\leq \text{dist}(\mathcal{T}_h(\mathbf{y}), \mathcal{M}) + \|\widehat{\mathcal{T}}_{\mathcal{Z},h}(\mathbf{y}) - \mathcal{T}_h(\mathbf{y})\| \\ &\leq \kappa \text{dist}(\mathbf{y}, \mathcal{M}) + C\sigma^2 + Chr_N. \end{aligned}$$

Let $\beta_N = C(\sigma^2 + hr_N)$. Iterating this affine recursion, as long as the orbit remains in $\Phi_{A\sigma}$, gives

$$\text{dist}(\widehat{\mathbf{y}}^{(t)}, \mathcal{M}) \leq \kappa^t \text{dist}(\widehat{\mathbf{y}}^{(0)}, \mathcal{M}) + \beta_N \frac{1 - \kappa^t}{1 - \kappa}.$$

Because $A_0 < A$ and $h = c\sigma$, decrease σ_{orb} and r_{orb} so that $\beta_N/(1 - \kappa) \leq (A - A_0)\sigma$. The last display then proves by induction that every iterate remains in $\Phi_{A\sigma}$, so the recursion is valid for all t . Dropping the factor $1 - \kappa^t$ proves the stated bound. If $r_N = O(\sigma)$, then $\sigma^2 + hr_N = O(\sigma^2)$. The event in Lemma S7 is uniform over the whole tube. Thus the induction uses the same event for every $t \geq 0$ and requires no derivative or continuity of the hard-radius empirical operator. $\square$

S3.2 Observed initialization and iteration complexity

Proposition S1 (Coverage of observed initializations). *Fix $0 < \eta < 1$ and choose constants $A_{\text{init}} < A_0 < A$ such that $\Pr\{\|\boldsymbol{\xi}\| \leq A_{\text{init}}\sigma\} \geq 1 - \eta$ for $\boldsymbol{\xi} \sim \mathcal{N}(\mathbf{0}, I_D)$. Then a generic observation satisfies*

$$\Pr\{\text{dist}(\mathbf{Y}, \mathcal{M}) \leq A_{\text{init}}\sigma\} \geq 1 - \eta.$$

Moreover, with probability at least $1 - \delta_{\text{init}}$, the fraction of the N observed initializations in $\Phi_{A_{\text{init}}\sigma}$ is at least

$$1 - \eta - \sqrt{\frac{\log(1/\delta_{\text{init}})}{2N}}.$$

Every such initialization falls within the fixed tube covered uniformly by Theorem S3.

Proof. Write $\mathbf{y}_i = \mathbf{x}_i + \sigma\boldsymbol{\xi}_i$ with $\boldsymbol{\xi}_i \sim \mathcal{N}(\mathbf{0}, I_D)$. Then $\text{dist}(\mathbf{y}_i, \mathcal{M}) \leq \sigma\|\boldsymbol{\xi}_i\|$, which proves the single-observation bound. The indicators of membership in $\Phi_{A_{\text{init}}\sigma}$ are independent Bernoulli variables with mean at least $1 - \eta$. Hoeffding's inequality gives the displayed sample fraction. Since $A_{\text{init}} < A_0$, every counted observation meets the initialization condition of the finite-sample orbit theorem. $\square$

Theorem S3 also gives a finite-iteration interpretation. Put $d_0 = \text{dist}(\widehat{\mathbf{y}}^{(0)}, \mathcal{M})$. If

$$L \geq \frac{\lceil \log \{ (1 - \kappa) d_0 / [C(\sigma^2 + hr_N)] \} \rceil_+}{|\log \kappa|},$$

then the L th output is within twice the statistical error radius of $\mathcal{M}$. If $d_0 = O(\sigma)$ and $\sigma^2 + hr_N = O(\sigma^2)$, this requires $L = O\{\log(1/\sigma)\}$.

S4 Theory for MS-STSP

We first derive the quadratic expansion of the clean local average that motivates squared-scale extrapolation. We then use stable geometric aggregation to establish localization for the noisy fixed-reference MS-STSP output.

S4.1 Quadratic structure of the clean local average

For $0 < h \leq c_0 r_{\text{geo}}$, recall

$$f_h(x) = \frac{1}{\text{Area}\{B_E(x, h) \cap \mathcal{M}\}} \int_{B_E(x, h) \cap \mathcal{M}} y \, dA(y).$$

Lemma S8 (Quadratic structure of local averaging bias). *Under conditions (A2)–(A4) of the main paper, uniformly in $x \in \mathcal{M}$,*

$$f_h(x) = x + \frac{1}{2} h^2 \mathbf{B}(x) + O(h^4),$$

where, for any orthonormal basis $\{e_1, \dots, e_d\}$ of $T_x \mathcal{M}$,

$$\mathbf{B}(x) = \frac{1}{d+2} \sum_{k=1}^d \Pi_x(e_k, e_k) \in N_x \mathcal{M}. \quad (\text{S4})$$

The remainder is uniform in x .

Proof. Fix $x \in \mathcal{M}$ and use geodesic normal coordinates $u \in T_x \mathcal{M}$. Let $\mathcal{J}_x(u)$ be the area density and $D_h(x) = \{u : \exp_x(u) \in B_E(x, h) \cap \mathcal{M}\}$. Then

$$f_h(x) - x = \frac{\int_{D_h(x)} \{\exp_x(u) - x\} \mathcal{J}_x(u) \, du}{\int_{D_h(x)} \mathcal{J}_x(u) \, du}.$$

Uniform Taylor expansions give

$$\begin{aligned} \exp_x(u) - x &= u + \frac{1}{2} \Pi_x(u, u) + F_{3,x}(u) + F_{4,x}(u) + O(\|u\|^5), \\ \mathcal{J}_x(u) &= 1 + \mathcal{J}_{2,x}(u) + \mathcal{J}_{3,x}(u) + O(\|u\|^4), \end{aligned}$$

where the indexed terms are homogeneous of the indicated degree.

It remains to account for the fact that $D_h(x)$ is not exactly a symmetric tangent ball. Write $u = r\theta$. Orthogonality of u and $\Pi_x(u, u)$ gives

$$\|\exp_x(r\theta) - x\|^2 = r^2 + r^4 H_{4,x}(\theta) + r^5 H_{5,x}(\theta) + O(r^6),$$

where $H_{4,x}$ is even in θ . Solving the boundary equation $\|\exp_x\{\rho_x(\theta, h)\theta\} - x\| = h$ yields

$$\rho_x(\theta, h) = h + h^3 \alpha_x(\theta) + h^4 \beta_x(\theta) + O(h^5), \quad \alpha_x(-\theta) = \alpha_x(\theta).$$

Thus the order- h^3 boundary perturbation has zero pairing with the odd linear term u . Its first possible nonzero boundary contribution to that term is $O(h^{d+4})$. All degree-one and degree-three interior terms also integrate to zero on the leading symmetric ball; the remaining numerator corrections are $O(h^{d+4})$. Moreover,

$$\int_{D_h(x)} \mathcal{J}_x(u) du = \omega_d h^d \{1 + O(h^2)\}.$$

Consequently,

$$f_h(x) - x = \frac{1}{\omega_d h^d} \int_{\|u\| \leq h} \frac{1}{2} \Pi_x(u, u) du + O(h^4).$$

Since

$$\int_{\|u\| \leq h} u_a u_b du = \mathbf{1}\{a = b\} \frac{\omega_d}{d+2} h^{d+2},$$

the leading term is exactly $\frac{1}{2} h^2 \mathbf{B}(x)$. The trace $\sum_a \Pi_x(e_a, e_a)$ is invariant under an orthogonal change of tangent basis, so (S4) is intrinsic. Compactness and the uniform bounds in (A2)–(A4) make the stated $O(h^4)$ remainder uniform over chart centres. $\square$

S4.2 Squared-scale intercept weights

Lemma S9 (OLS intercept identities). *Let $s_j = h_j^2$, $\bar{s} = M^{-1} \sum_j s_j$, $\bar{\mathbf{v}} = M^{-1} \sum_j \mathbf{v}_j$, and $S_{ss} = \sum_j (s_j - \bar{s})^2 > 0$. The intercept in the ordinary least-squares regression of vectors $\mathbf{v}_j \in \mathbb{R}^D$ on $(1, s_j)$ is $\sum_j w_j \mathbf{v}_j$, where*

$$w_j = \frac{1}{M} - \frac{\bar{s}(s_j - \bar{s})}{S_{ss}}.$$

These weights satisfy

$$\sum_j w_j = 1, \quad \sum_j w_j s_j = 0.$$

If $h_j = c_j \sigma$ with fixed positive c_j that are not all equal, then the weights do not depend on σ and $\Lambda_w := \sum_j |w_j| < \infty$.

Proof. The OLS slope is $\hat{\mathbf{b}} = S_{ss}^{-1} \sum_j (s_j - \bar{s}) \mathbf{v}_j$ and the intercept is $\bar{\mathbf{v}} - \bar{s} \hat{\mathbf{b}}$, which gives the displayed formula. Since $\sum_j (s_j - \bar{s}) = 0$,

$$\sum_j w_j = 1.$$

Furthermore,

$$\sum_j (s_j - \bar{s}) s_j = \sum_j (s_j - \bar{s})^2 = S_{ss},$$

so $\sum_j w_j s_j = \bar{s} - \bar{s} = 0$. If $s_j = c_j^2 \sigma^2$, the common factor σ^2 cancels from $\bar{s}(s_j - \bar{s})/S_{ss}$. Thus every w_j is a fixed finite number and so is its finite ℓ_1 norm. $\square$

For the clean local-average map in Lemma S8, the two weight identities give, uniformly in $x \in \mathcal{M}$,

$$\sum_{j=1}^M w_j f_{h_j}(x) = x + O\left(\Lambda_w h_{\max}^4\right), \quad h_{\max} = \max_j h_j.$$

This follows by cancelling the common $h_j^2 \mathbf{B}(x)$ term and bounding the weighted remainders by $C \sum_j |w_j| h_j^4 \leq C \Lambda_w h_{\max}^4$. The noisy-output result below instead uses geometric aggregation.

S4.3 A geometric aggregation lemma

Lemma S10 (Stable affine aggregation in one manifold patch). *Fix $a_{\text{patch}} < \infty$ and an a priori weight cap $C_w < \infty$. There are $\sigma_{\text{agg}}, C < \infty$, depending only on a_{patch}, C_w and the uniform geometric constants, with the following property. Let $0 < \sigma \leq \sigma_{\text{agg}}$, $x_0 \in \mathcal{M}$, and $\mathbf{v}_1, \dots, \mathbf{v}_M$ lie in the reach tube. Put $x_j = \Pi_{\mathcal{M}}(\mathbf{v}_j)$ and suppose*

$$d_g(x_j, x_0) \leq a_{\text{patch}}\sigma, \quad \text{dist}(\mathbf{v}_j, \mathcal{M}) \leq \varepsilon_{\text{agg}}.$$

If $\sum_j w_j = 1$ and $\sum_j |w_j| \leq C_w$, then

$$\text{dist}\left(\sum_j w_j \mathbf{v}_j, \mathcal{M}\right) \leq CC_w(\varepsilon_{\text{agg}} + \sigma^2).$$

The constant C may depend on the fixed cap C_w , but not on $\sigma, \varepsilon_{\text{agg}}, N$, or the particular points.

Proof. Use an orthogonal chart centred at x_0 in which the manifold is the graph

$$x_0 + u + \psi(u), \quad u \in T_{x_0}\mathcal{M},$$

with $\psi(\mathbf{0}) = \mathbf{0}$, $D\psi(\mathbf{0}) = 0$, and $\|\psi(u)\| \leq C\|u\|^2$ on a fixed ball. Positive reach and the derivative bounds give this chart uniformly in x_0 . For sufficiently small σ , geodesic–Euclidean comparability puts every x_j in this chart, so write $x_j = x_0 + u_j + \psi(u_j)$ with $\|u_j\| \leq Ca_{\text{patch}}\sigma$. Also write $\mathbf{v}_j = x_j + n_j$, where $\|n_j\| \leq \varepsilon_{\text{agg}}$.

Set $u_w = \sum_j w_j u_j$. Then $\|u_w\| \leq CC_w a_{\text{patch}}\sigma$, so after reducing σ_{agg} the point $x_w = x_0 + u_w + \psi(u_w)$ belongs to $\mathcal{M}$ in the same chart. Because $\sum_j w_j = 1$,

$$\sum_j w_j \mathbf{v}_j - x_w = \sum_j w_j \psi(u_j) - \psi(u_w) + \sum_j w_j n_j.$$

The normal-error term has norm at most $C_w \varepsilon_{\text{agg}}$. The graph bounds give

$$\left\| \sum_j w_j \psi(u_j) - \psi(u_w) \right\| \leq C \left\{ C_w a_{\text{patch}}^2 + C_w^2 a_{\text{patch}}^2 \right\} \sigma^2.$$

Since the weight cap is fixed and $C_w \geq |\sum_j w_j| = 1$, this is bounded by $CC_w \sigma^2$ after absorbing the fixed cap into C . Taking distance to the particular manifold point x_w proves the result. $\square$

S4.4 Population MS-STSP

Theorem S4 (Population MS-STSP localization). *Let M be fixed, let $h_j = c_j \sigma$, and suppose every (c_j, A) is spectrally admissible. For each scale, let $\mathcal{M}_{h_j}$ and $\mathcal{U}_{h_j}$ be supplied by Lemma S6, and write $\rho_{0,j}$, $\bar{\kappa}_j$, and $C_{\text{step},j}$ for its scale- h_j constants. Assume*

$$\mathbf{y}^{(0)} \in \bigcap_{j=1}^M \mathcal{U}_{h_j}.$$

Let

$$\mathbf{y}_j^{(\infty)} = \lim_{t \rightarrow \infty} \mathcal{T}_{h_j}^{(t)}(\mathbf{y}^{(0)}), \quad \mathbf{y}_{\text{MS}}^{(\infty)} = \sum_{j=1}^M w_j \mathbf{y}_j^{(\infty)},$$

where $\sum_j w_j = 1$ and $\sum_j |w_j| \leq C_w$ for a fixed cap. Then, for all sufficiently small σ ,

$$\text{dist}(\mathbf{y}_j^{(\infty)}, \mathcal{M}) \leq C\sigma^2, \quad d_g\{\Pi_{\mathcal{M}}(\mathbf{y}_j^{(\infty)}), \Pi_{\mathcal{M}}(\mathbf{y}^{(0)})\} \leq C\sigma,$$

uniformly over j , and

$$\text{dist}(\mathbf{y}_{\text{MS}}^{(\infty)}, \mathcal{M}) \leq CC_w \sigma^2.$$

Here and below C may depend on the fixed finite scale design and its weight cap, but not on σ .

Proof. Apply Theorem S2 separately at every scale. Because M is fixed, the maxima of the finitely many constants are still finite. Thus every limit exists, belongs to $\mathcal{M}_{h_j}$, and $\text{dist}(\mathbf{y}_j^{(\infty)}, \mathcal{M}) \leq C\sigma^2$.

The tail estimate in that theorem at $t = 0$ gives

$$\|\mathbf{y}_j^{(\infty)} - \mathbf{y}^{(0)}\| \leq \frac{C_{\text{step},j}}{1 - \bar{\kappa}_j} \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_{h_j}) \leq C\sigma,$$

because $\mathbf{y}^{(0)} \in \mathcal{U}_{h_j}$. Let $x_0 = \Pi_{\mathcal{M}}(\mathbf{y}^{(0)})$ and $x_j = \Pi_{\mathcal{M}}(\mathbf{y}_j^{(\infty)})$. Nearest-point projection is uniformly Lipschitz on a smaller reach tube, so $\|x_j - x_0\| \leq C\sigma$. Local geodesic–Euclidean comparability then gives $d_g(x_j, x_0) \leq C\sigma$. Lemma S10, with $\varepsilon_{\text{agg}} = C\sigma^2$, proves the last display. $\square$

S4.5 Finite-sample MS-STSP

Theorem S5 (Finite-sample MS-STSP localization). *Retain the finite scale design of Theorem S4. Let $0 < \delta < 1$ and assume the common initialization has a uniform attraction-tube buffer,*

$$\text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_{h_j}) \leq \frac{1}{2}\rho_{0,j}\sigma, \quad j = 1, \dots, M.$$

Define

$$r_{N,j} = \sqrt{\frac{D \log(1/h_j) + \log(M/\delta)}{Nh_j^d}} + \frac{D \log(1/h_j) + \log(M/\delta)}{Nh_j^d}.$$

Suppose $\max_j r_{N,j}$ is small enough that the joint empirical event below holds and

$$\frac{Ch_j r_{N,j}}{1 - \bar{\kappa}_j} \leq \frac{1}{2}\rho_{0,j}\sigma \quad \text{for every } j.$$

For predetermined finite iteration horizons L_j and a fixed stopping-horizon bound $C_{\text{hor}} < \infty$, assume

$$\bar{\kappa}_j^{L_j} \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_{h_j}) \leq C\{\sigma^2 + h_j r_{N,j}\}, \quad L_j\{\sigma^2 + h_j r_{N,j}\} \leq C_{\text{hor}}\sigma.$$

For $t = 0, \dots, L_j$, define the empirical branch orbit and its aggregate by

$$\widehat{\mathbf{y}}_j^{(t)} = \widehat{\mathcal{T}}_{\mathcal{Z}, h_j}^{(t)}(\mathbf{y}^{(0)}), \quad \widehat{\mathbf{y}}_{\text{MS}} = \sum_{j=1}^M w_j \widehat{\mathbf{y}}_j^{(L_j)},$$

with $\sum_j w_j = 1$ and $\sum_j |w_j| \leq C_w$. Then, with probability at least $1 - \delta$,

$$\text{dist}(\widehat{\mathbf{y}}_{\text{MS}}, \mathcal{M}) \leq CC_w \left\{ \sigma^2 + \max_j h_j r_{N,j} \right\}.$$

If $h_j = c_j \sigma$ and

$$N \geq C\sigma^{-(d+2)} \{D \log(1/\sigma) + \log(M/\delta)\},$$

then $\max_j r_{N,j} = O(\sigma)$ and the right-hand side is $O(\sigma^2)$.

Proof. Apply the uniform empirical approximation in Lemma S7 at scale j with failure probability δ/M . A union bound gives, with probability at least $1 - \delta$, simultaneously for every j ,

$$\sup_{\mathbf{y} \in \Phi_{A\sigma}} \|\widehat{\mathcal{T}}_{\mathcal{Z}, h_j}(\mathbf{y}) - \mathcal{T}_{h_j}(\mathbf{y})\| \leq Ch_j r_{N,j},$$

and all empirical projectors are uniquely defined by their eigengaps.

On the attraction tube, Lemma S6(iii) and the last display imply

$$\text{dist}(\widehat{\mathcal{T}}_{\mathcal{Z}, h_j}(\mathbf{y}), \mathcal{M}_{h_j}) \leq \bar{\kappa}_j \text{dist}(\mathbf{y}, \mathcal{M}_{h_j}) + Ch_j r_{N,j}.$$

The initialization buffer and the stated smallness condition prove by induction that the empirical branch remains in $\mathcal{U}_{h_j}$. Hence

$$\text{dist}(\widehat{\mathbf{y}}_j^{(L_j)}, \mathcal{M}_{h_j}) \leq \bar{\kappa}_j^{L_j} \text{dist}(\mathbf{y}^{(0)}, \mathcal{M}_{h_j}) + \frac{Ch_j r_{N,j}}{1 - \bar{\kappa}_j}.$$

Since $\sup_{\mathbf{z} \in \mathcal{M}_{h_j}} \text{dist}(\mathbf{z}, \mathcal{M}) \leq C\sigma^2$, the transient condition yields

$$\text{dist}(\widehat{\mathbf{y}}_j^{(L_j)}, \mathcal{M}) \leq C\{\sigma^2 + h_j r_{N,j}\}.$$

We next establish cross-scale locality. For the empirical iterate $\widehat{\mathbf{y}}_j^{(t)}$, the population step bound in Lemma S6(iii) and the uniform perturbation bound give

$$\|\widehat{\mathbf{y}}_j^{(t+1)} - \widehat{\mathbf{y}}_j^{(t)}\| \leq C_{\text{step}, j} \text{dist}(\widehat{\mathbf{y}}_j^{(t)}, \mathcal{M}_{h_j}) + Ch_j r_{N,j}.$$

Summing the preceding affine recursion for $t = 0, \dots, L_j - 1$ gives

$$\|\widehat{\mathbf{y}}_j^{(L_j)} - \mathbf{y}^{(0)}\| \leq C\sigma + CL_j h_j r_{N,j} \leq C\sigma,$$

where the last inequality uses the stopping-horizon condition. Projection Lipschitzness and local geodesic–Euclidean comparability therefore show that all $\Pi_{\mathcal{M}}(\widehat{\mathbf{y}}_j^{(L_j)})$ lie within $C\sigma$ of $\Pi_{\mathcal{M}}(\mathbf{y}^{(0)})$ in geodesic distance.

Apply Lemma S10 with $\varepsilon_{\text{agg}} = C\{\sigma^2 + \max_j h_j r_{N,j}\}$ to obtain the asserted multi-scale bound. Finally, for fixed M and fixed positive c_j , the displayed sample-size lower bound makes every $r_{N,j} = O(\sigma)$, which proves the last statement. $\square$

S5 Numerical and reproducibility details

S5.1 Simulation manifolds

Curves.

1. Tilted Circle ($\vartheta \in [0, 2\pi]$):

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos(\pi/6) & -\sin(\pi/6) \\ 0 & \sin(\pi/6) & \cos(\pi/6) \end{pmatrix} \begin{pmatrix} \cos(\vartheta) \\ \sin(\vartheta) \\ 0 \end{pmatrix}$$

2. Tennis Curve ($\vartheta \in [-\pi, \pi]$):

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \cos^3(\vartheta) \\ \sin^3(\vartheta) \\ \frac{\sqrt{3}}{2} \sin(2\vartheta) \end{pmatrix}$$

3. Trefoil Knot ($\vartheta \in [0, 2\pi]$):

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \sin(\vartheta) + 2 \sin(2\vartheta) \\ \cos(\vartheta) - 2 \cos(2\vartheta) \\ -\sin(3\vartheta) \end{pmatrix}$$

4. Sixfolds ($\vartheta \in [0, 12\pi]$):

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \cos(\vartheta)(1 + 0.3 \cos(6\vartheta)) \\ \sin(\vartheta)(1 + 0.3 \cos(6\vartheta)) \\ 0.3 \sin(6\vartheta) \end{pmatrix}$$

Surfaces.


1. Sphere ($\theta \in [0, 2\pi], \phi \in [0, \pi]$):

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \sin(\phi) \cos(\theta) \\ \sin(\phi) \sin(\theta) \\ \cos(\phi) \end{pmatrix}$$


2. Torus ($\theta, \phi \in [0, 2\pi]$):

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} (1 + 0.3 \cos(\phi)) \cos(\theta) \\ (1 + 0.3 \cos(\phi)) \sin(\theta) \\ 0.3 \sin(\phi) \end{pmatrix}$$


(a)




(b)



(c)



(d)



(e)



(f)



Figure S1: Underlying manifolds used in the synthetic study: (a) circle, (b) tennis curve, (c) trefoil knot, (d) sixfolds, (e) sphere, and (f) torus.

For each independent synthetic repetition, the curve input is sampled by inverting a numerical cumulative arc-length table with 100,001 parameter grid points. The Sixfolds interval traverses the same curve six times; equal-length traversals induce the same arc-length distribution as one fundamental period. Sphere inputs are sampled using a uniform height coordinate and a uniform azimuth. Torus inputs are uniform in surface area: θ is uniform on $[0, 2\pi]$, while ϕ has density $(1 + 0.3 \cos \phi)/(2\pi)$, generated by numerically inverting $F(\phi) = (\phi + 0.3 \sin \phi)/(2\pi)$. Independent Gaussian noise is then added coordinatewise. The evaluation clouds contain 20,000 points for curves and 30,000 points for surfaces and are held fixed across repetitions of a given geometry. Distances from fitted points to the manifold are approximated by nearest-neighbour distances to the corresponding independent evaluation cloud.

The synthetic studies use a common selection rule. Within each synthetic repetition, every method receives the same underlying-manifold realisation, noisy realisation, independent evaluation cloud, sample size, and noise level. Recall the fitted cloud $\hat{\mathcal{Y}}$, independent evaluation cloud $\mathcal{X}_{\text{eval}}$, and the two mean errors defined in Section 5.1 of the main paper:

$$e_{\text{fit}} = \frac{1}{N} \sum_{i=1}^N \text{dist}(\hat{\mathbf{y}}_i, \mathcal{X}_{\text{eval}}), \quad e_{\text{cov}} = \frac{1}{N_{\text{eval}}} \sum_{\ell=1}^{N_{\text{eval}}} \text{dist}(\mathbf{x}_{\text{eval}, \ell}, \hat{\mathcal{Y}}).$$

Tunable parameters are selected separately for each method on the full data set by minimizing

$$J_{\text{mean}} = \max(e_{\text{fit}}, e_{\text{cov}}),$$

with ties resolved first by the analogous joint 95th-percentile error and then by the maximum fitting error. The independent evaluation cloud provides a finite-point approximation to distance from a fitted point to the manifold. This is method-wise full-data oracle tuning. The displayed e_{fit} and e_{cov} are means over all fitted and evaluation points, respectively.

S5.2 Common search spaces

For STSP, MS-STSP, local PCA, and mean shift, the neighbourhood of each query consists of all fixed reference observations within radius h . The ordinary synthetic studies use

$$h/\sigma \in \{1.5, 2, 2.5, 3, 3.5, 4, 5, 6, 8\}.$$

Local PCA uses $L = 1$. STSP and mean shift stop at the first $t \geq 1$ for which the mean displacement divided by s is below 10^{-3} , or at T_{max} , whichever occurs first. In the synthetic studies, $s = \sigma$ and $T_{\text{max}} = 30$. MS-STSP applies the same rule to the largest normalised mean displacement among its constituent branches and searches finite scale templates from the same radius pool, with an ℓ_1 weight screen to exclude unstable extrapolation designs. STSP and every MS-STSP branch use the original noisy cloud as their fixed reference at every iteration. All branches are evaluated at the common stopping iteration. Theorem S5 applies to finite horizons that satisfy its transient and accumulated-drift conditions. The displacement rule used in the experiments is a computational stopping criterion; it does not by itself verify those theorem conditions for the selected stopping time. For the ordinary synthetic settings, MS-STSP uses the seven triples of adjacent values in the ordered h/σ grid:

$$(1.5, 2, 2.5), (2, 2.5, 3), (2.5, 3, 3.5), (3, 3.5, 4), \\ (3.5, 4, 5), (4, 5, 6), (5, 6, 8).$$

Each triple is multiplied by σ to obtain its three radii. The high-noise study uses

$$h/\sigma \in \{1, 1.25, 1.5, 2, 2.5, 3, 3.5, 4\}$$

and the six triples of adjacent values in this grid:

$$(1, 1.25, 1.5), (1.25, 1.5, 2), (1.5, 2, 2.5), (2, 2.5, 3), \\ (2.5, 3, 3.5), (3, 3.5, 4).$$

These triples are also multiplied by σ . In both studies, triples with $\sum_j |w_j| > 5$ are excluded.

As in the main paper, the estimator of Yao et al. (2023) is called the Yao–Yau (YY) estimator. YY-adapted retains the direction from each query to its local centre in the published YY construction. Within the orthogonal complement of that direction, local PCA estimates the known d -dimensional tangent subspace; the remaining directions form the normal subspace used by the update. The published averaging option is enabled, and the routine’s noise input is set to the simulation value, that is, $\text{sig} = \sigma$. Its internal radii are therefore the published $r = 5\sigma/\log_{10} N$ and $R = 10\sigma\sqrt{\log(1/\sigma)/\log_{10} N}$. For a query $\mathbf{y}$, let $\mathcal{I}_{YY}(\mathbf{y})$ be the union of the reference points within radius $2r$ and its nearest $\min\{N, \max(5, d+2)\}$ reference points. Up to floating-point precision, the local centre is

$$\mathbf{m}_{YY}(\mathbf{y}) = |\mathcal{I}_{YY}(\mathbf{y})|^{-1} \sum_{i \in \mathcal{I}_{YY}(\mathbf{y})} \mathbf{z}_i, \quad \mathbf{v}_{YY}(\mathbf{y}) = \mathbf{y} - \mathbf{m}_{YY}(\mathbf{y}).$$

For a nonzero displacement, choose an orthonormal basis matrix $\mathbf{U}_{YY}$ for $\mathbf{v}_{YY}^\perp$ and form the matrix with rows $(\mathbf{z}_i - \mathbf{m}_{YY})^\top \mathbf{U}_{YY}$, $i \in \mathcal{I}_{YY}(\mathbf{y})$. If $\mathbf{V}_{YY,d}$ contains its first d right singular vectors, the tangent projector is

$$\mathbf{P}_{YY}(\mathbf{y}) = \mathbf{U}_{YY} \mathbf{V}_{YY,d} \mathbf{V}_{YY,d}^\top \mathbf{U}_{YY}^\top.$$

The cylindrical averaging set is

$$\mathcal{C}_{YY}(\mathbf{y}) = \{i : \|\mathbf{P}_{YY}(\mathbf{y})(\mathbf{z}_i - \mathbf{y})\| < r, \quad \|(I_D - \mathbf{P}_{YY}(\mathbf{y}))(\mathbf{z}_i - \mathbf{y})\| < R\}.$$

The preliminary output is the mean over $\mathcal{C}_{YY}(\mathbf{y})$ when this set contains more than ten points, and is $\mathbf{m}_{YY}(\mathbf{y})$ otherwise. The final averaging pass takes the mean of preliminary outputs within radius $r/4$, augmented by the five nearest preliminary outputs (or all outputs if fewer than five are available).

The original YY construction uses the full space $\mathbf{v}_{YY}^\perp$ as its tangent space, with projector $I_D - \mathbf{v}_{YY} \mathbf{v}_{YY}^\top / \|\mathbf{v}_{YY}\|^2$. YY-adapted selects d directions within that space, so that the normal complement has dimension $D - d$. For $d = D - 1$, the published routine is used. For $d < D - 1$, local PCA selects the tangent directions while retaining the mean-displacement direction in the normal space. The radii, cylindrical averaging rule, and final averaging pass retain the published construction.

MF25 uses $L_{\max} = 50$, $\varepsilon_{\text{opt}} = 10^{-8}$, initial step size one, and ten backtracking steps. Only $h = c_{\text{MF25}} \sqrt{\sigma}$ is selected, with candidate multipliers $c_{\text{MF25}} \in \{.75, 1, 1.25, 1.5, 1.75, 2, 2.5\}$.

MF25 uses the complete noisy cloud as both the reference set and the initial query set; all N fitted outputs enter the evaluation.

Table S2: MF25 radius multipliers and radii selected for the first realisation displayed in the fitted-cloud figures. YY-adapted has no selected external parameter.

Study	Data	c_{MF25}	$h = c_{\text{MF25}} \sqrt{\sigma}$
Curves	Circle	1.00	0.2828
Curves	Tennis curve	1.00	0.2828
Curves	Trefoil knot	1.75	0.4950
Curves	Sixfolds	1.00	0.2828
Surfaces	Sphere	1.00	0.2828
Surfaces	Torus	0.75	0.2121
High noise	$\sigma = 0.1$	1.25	0.3953
High noise	$\sigma = 0.2$	1.25	0.5590
High noise	$\sigma = 0.3$	1.25	0.6847

S5.3 SCMS configurations

Both SCMS variants use isotropic Gaussian weights over the complete fixed reference cloud, with all noisy observations as initial queries. SCMS-KDE takes the eigenvectors associated with the smallest $D - d$ eigenvalues of the density Hessian; SCMS-log-KDE uses the corresponding eigenvectors of the log-density Hessian. Each update adds the projection of the Gaussian mean-shift vector onto this subspace. The intrinsic dimension is supplied in the synthetic experiments. The ordinary synthetic bandwidth pool is

$$h/\sigma \in \{.35, .5, .75, 1, 1.25, 1.5, 2, 2.5, 3, 3.5, 4, 5, 6, 8\},$$

and the high-noise pool is

$$h/\sigma \in \{.25, .35, .5, .75, 1, 1.25, 1.5, 2, 2.5, 3, 3.5, 4\}.$$

The full-data geometric criterion selects the bandwidth separately for each variant. The reported comparisons use $T_{\max} = 30$ and the common mean-displacement threshold. Some SCMS-KDE configurations reach the iteration cap.

S5.4 Ordinary simulation repetitions

For each of the six ordinary manifold settings, ten independent noisy samples were generated. The evaluation cloud is fixed within each setting and is independent of the noisy samples. The method-specific search described above was repeated on each noisy sample. The main-paper tables report the mean and standard deviation of each error across the ten repetitions.

Table S3: Configurations selected for the first realisation of the six synthetic settings. The four curves use $N = 5000$ and $\sigma = 0.08$; the sphere and torus use $N = 20000$ and $\sigma = 0.08$. A list in the radius column gives the three MS-STSP scales.

Data	Method	h/σ	L	Radius or scales
Circle	Local PCA	4	1	0.32
Circle	Mean shift	1.5	28	0.12
Circle	SCMS-KDE	2	30	0.16
Circle	SCMS-log-KDE	1.25	9	0.1
Circle	STSP	2.5	8	0.2
Circle	MS-STSP	–	5	0.32, 0.4, 0.48
Tennis curve	Local PCA	4	1	0.32
Tennis curve	Mean shift	1.5	25	0.12
Tennis curve	SCMS-KDE	2	30	0.16
Tennis curve	SCMS-log-KDE	1.25	9	0.1
Tennis curve	STSP	2.5	8	0.2
Tennis curve	MS-STSP	–	5	0.32, 0.4, 0.48
Trefoil knot	Local PCA	5	1	0.4
Trefoil knot	Mean shift	2	22	0.16
Trefoil knot	SCMS-KDE	2.5	14	0.2
Trefoil knot	SCMS-log-KDE	2	6	0.16
Trefoil knot	STSP	4	5	0.32
Trefoil knot	MS-STSP	–	4	0.4, 0.48, 0.64
Sixfolds	Local PCA	3.5	1	0.28
Sixfolds	Mean shift	1.5	22	0.12
Sixfolds	SCMS-KDE	1.25	30	0.1
Sixfolds	SCMS-log-KDE	1	13	0.08
Sixfolds	STSP	2	12	0.16
Sixfolds	MS-STSP	–	6	0.24, 0.28, 0.32
Sphere	Local PCA	3.5	1	0.28
Sphere	Mean shift	1.5	25	0.12
Sphere	SCMS-KDE	1.25	30	0.1
Sphere	SCMS-log-KDE	1	11	0.08
Sphere	STSP	2	11	0.16
Sphere	MS-STSP	–	5	0.4, 0.48, 0.64
Torus	Local PCA	3	1	0.24
Torus	Mean shift	1.5	27	0.12
Torus	SCMS-KDE	0.75	30	0.06
Torus	SCMS-log-KDE	1	13	0.08
Torus	STSP	2	12	0.16
Torus	MS-STSP	–	8	0.24, 0.28, 0.32




Figure S2: Fitting and coverage errors for the circle, tennis curve, and trefoil knot, from top to bottom, over ten independent realisations. Each row pairs a scatter plot of method means on log-log axes with horizontal bars showing the two mean errors. Error bars show one standard deviation; the grey diagonal marks equal errors in the scatter plot. Smaller errors are preferable.




Figure S3: Fitting and coverage errors for Sixfolds, the sphere, and the torus, from top to bottom, continuing Figure S2. Each setting uses ten independent realisations; graphical conventions are the same as in Figure S2.




Figure S4: Circle fitted clouds. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.




Figure S5: Tennis-curve fitted clouds. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.




Figure S6: Trefoil-knot fitted clouds. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.




Figure S7: Sixfolds fitted clouds. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.




Figure S8: Sphere fitted clouds. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.




Figure S9: Torus fitted clouds. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.

S5.5 High-noise circle experiment

The high-noise study uses $N = 5000$ and $\sigma \in \{0.1, 0.2, 0.3\}$. We compare STSP, MS-STSP, YY-adapted, MF25, local PCA, mean shift, SCMS-KDE, and SCMS-log-KDE. At each noise level each tunable method is selected by the geometric criterion using the high-noise candidate sets specified above.

Table S4: Configurations selected for the first realisation at each high-noise level.

Data	Method	h/σ	L	Radius or scales
$\sigma = 0.1$	Local PCA	4	1	0.4
$\sigma = 0.1$	Mean shift	2	30	0.2
$\sigma = 0.1$	SCMS-KDE	2	30	0.2
$\sigma = 0.1$	SCMS-log-KDE	1.25	9	0.125
$\sigma = 0.1$	STSP	2.5	8	0.25
$\sigma = 0.1$	MS-STSP	—	6	0.3, 0.35, 0.4
$\sigma = 0.2$	Local PCA	3	1	0.6
$\sigma = 0.2$	Mean shift	1.5	30	0.3
$\sigma = 0.2$	SCMS-KDE	1.25	30	0.25
$\sigma = 0.2$	SCMS-log-KDE	0.75	17	0.15
$\sigma = 0.2$	STSP	1.5	18	0.3
$\sigma = 0.2$	MS-STSP	—	6	0.6, 0.7, 0.8
$\sigma = 0.3$	Local PCA	3	1	0.9
$\sigma = 0.3$	Mean shift	1.25	30	0.375
$\sigma = 0.3$	SCMS-KDE	0.5	30	0.15
$\sigma = 0.3$	SCMS-log-KDE	0.75	17	0.225
$\sigma = 0.3$	STSP	1.5	17	0.45
$\sigma = 0.3$	MS-STSP	—	9	0.75, 0.9, 1.05




Figure S10: Fitting and coverage errors over ten independent realisations of the high-noise circle. Each row gives a scatter plot of method means and paired horizontal bars for e_{fit} and e_{cov} . Error bars show one standard deviation; the grey diagonal marks equal errors in the scatter plot.




Figure S11: Circle fitted clouds at $\sigma = 0.1$. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.




Figure S12: Circle fitted clouds at $\sigma = 0.2$. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.




Figure S13: Circle fitted clouds at $\sigma = 0.3$. Panels (a)–(h) show local PCA, mean shift, SCMS-KDE, SCMS-log-KDE, YY-adapted, MF25, STSP, and MS-STSP, respectively.

S5.6 Rate, convergence, and radius diagnostics

The expanded rate diagnostic uses hard-radius STSP with $N = 5000$, $h = 3\sigma$, $T_{\max} = 30$, the displacement stopping rule above, 10 replicates, and

$$\sigma \in \{.01, .02, .04, .08, .10, .15, .20, .25, .30, .35\}$$

on the circle and sphere. The iteration diagnostic uses the same two manifolds, $N = 5000$, $\sigma = .08$, $h = 3\sigma$, 30 iterations, and 10 replicates. The radius-sensitivity study uses the circle, 15 geometrically spaced radii from $.2\sigma$ to $.5$, 10 iterations, and ten replicates at each of the two displayed noise levels.

S5.7 Dentate gyrus configuration

The supplied count matrix contains 2,930 cells and 14 annotations: astrocytes (120), Cajal–Retzius cells (34), Cck–Tox cells (27), endothelial cells (87), GABAergic cells (61), immature granule cells (785), mature granule cells (1,070), microglia (81), mossy cells (75), neuroblasts (417), oligodendrocytes (50), oligodendrocyte precursor cells (53), radial glia-like cells (51), and neuronal intermediate progenitor cells (19). Counts are normalised to a library size of 10^4 and log-transformed. The 2,000 genes with the largest variance are retained, standardised coordinatewise, and clipped to $[-10, 10]$. A randomised PCA with seed 20260925 produces one nested 100-component representation. For each displayed value of p , the first p components are centred and divided by their root-mean-square cloud radius.

The principal analysis uses $p = 30$. The dimension comparison uses $p \in \{20, 30, 40, 50\}$. Mean shift, local PCA, STSP, MS-STSP, both SCMS variants, YY-adapted, and MF25 use the common candidate scale grid formed by the upper 20 values of a 30-point logarithmically spaced sequence from 0.025 to 2. Equivalently, its unrounded values are $h_j = 0.025(2/0.025)^{(j+10)/29}$ for $j = 0, \dots, 19$; to six decimal places, this gives

$$\begin{aligned} h \in \{ & 0.113286, 0.131766, 0.153259, 0.178258, 0.207335, \\ & 0.241155, 0.280492, 0.326245, 0.379462, 0.441359, \\ & 0.513353, 0.597090, 0.694486, 0.807769, 0.939531, \\ & 1.092785, 1.271038, 1.478368, 1.719516, 2\}. \end{aligned}$$

Mean shift, local PCA, and STSP use the reference observations within radius h of each query. The SCMS variants use Gaussian weights with bandwidth h ; YY-adapted and MF25 retain their method-specific neighbourhoods. MS-STSP uses three distinct radii; all $\binom{20}{3} = 1140$ triples from the 20-scale grid satisfy $\sum_j |w_j| \leq 5$ and are evaluated at each p . The triple with the largest full-cell average silhouette is selected. Each branch applies the fixed-reference STSP update, with multi-scale weights as in Algorithm 2. Methods requiring an intrinsic dimension use $d = 1$. Mean shift and STSP retain the original PCA cloud as the reference throughout their iterations. With the normalised input radius $s = 1$, mean shift, STSP, and both SCMS variants stop at the first iteration satisfying

$$\frac{1}{N} \sum_{i=1}^N \|\mathbf{y}_i^{(t)} - \mathbf{y}_i^{(t-1)}\| < 10^{-3},$$

or at $T_{\max} = 30$. MS-STSP uses the same displacement criterion for each of its three branches, stopping when all branches satisfy it or at $T_{\max} = 30$. Local PCA applies one projection. YY-adapted uses its published stopping rule. MF25 uses 10 initialization iterations, 50 maximum outer iterations and tolerance 10^{-8} . At each p , the full-cell average Euclidean silhouette with respect to the 14 supplied annotations selects the radius or, for MS-STSP, the radius triple.

At $p = 30$, the selected radii are 0.513353 for mean shift, 0.939531 for local PCA, 0.441359 for SCMS-KDE, 0.326245 for SCMS-log-KDE, 0.326245 for YY-adapted, 0.939531 for MF25, and 0.694486 for STSP. Mean shift stops after 18 updates and STSP after 11. At $p = (20, 30, 40, 50)$, the selected MS-STSP radius triples are (0.694486, 1.478368, 1.719516), (0.694486, 1.271038, 1.719516), (0.807769, 1.271038, 1.719516), and (0.807769, 1.271038, 1.719516), respectively. Their stopping times are (23, 16, 14, 19). At the same dimensions, the selected STSP radii are (0.694486, 0.694486, 0.807769, 0.807769) and the corresponding stopping times are (10, 11, 9, 11). The selected mean-shift radii are (0.513353, 0.513353, 0.597090, 0.694486), with stopping times (17, 18, 19, 16). Both STSP and mean shift reach the displacement threshold before the iteration cap at these dimensions.

For each selected $p = 30$ output, k -means uses 14 clusters, random seeds 0–9, five k -means++ initializations per seed, and at most 500 iterations. ARI and geometric NMI compare each partition with the supplied annotations; accuracy uses an optimal one-to-one matching between cluster and annotation labels. The main-paper table reports averages over the 10 seeds.

For the main-paper t-SNE figure, the selected STSP and MS-STSP outputs are separately centred and divided by their root-mean-square radii. Each is embedded with perplexity 80, early exaggeration 24, PCA initialisation, automatic learning rate, 1,500 optimisation iterations, and random seed 20260927. Both embeddings are aligned by a global orthogonal transformation to the independently computed input embedding; this allows rotation and reflection but no local deformation. The annotations determine colours in the t-SNE display and are not supplied to the embedding algorithm; they also enter the SIL criterion used to select the fitted representation.

References

- Olivier Bousquet. A Bennett concentration inequality and its application to suprema of empirical processes. *Comptes Rendus Mathématique*, 334(6):495–500, 2002.
- Herm Jan Brascamp and Elliott H. Lieb. On extensions of the Brunn–Minkowski and Prékopa–Leindler theorems, including inequalities for log concave functions, and with an application to the diffusion equation. *Journal of Functional Analysis*, 22(4):366–389, 1976.
- Ming-Yen Cheng and Hau-Tieng Wu. Local linear regression on manifolds and its geometric interpretation. *Journal of the American Statistical Association*, 108(504):1421–1434, 2013. doi: 10.1080/01621459.2013.827984.
- Dorin Comaniciu and Peter Meer. Mean shift: A robust approach toward feature space analysis. *IEEE Transactions on Pattern Analysis and Machine Intelligence*, 24(5):603–619, 2002. doi: 10.1109/34.1000236.
- Manfredo P. do Carmo. *Riemannian Geometry*. Birkhäuser, Boston, 1992.
- Charles Fefferman, Sanjoy Mitter, and Hariharan Narayanan. Testing the manifold hypothesis. *Journal of the American Mathematical Society*, 29(4):983–1049, 2016.
- Charles Fefferman, Sergei Ivanov, Yaroslav Kurylev, Matti Lassas, and Hariharan Narayanan. Fitting a putative manifold to noisy data. In *Proceedings of the 31st Conference on Learning Theory*, volume 75 of *Proceedings of Machine Learning Research*, pages 688–720, 2018. URL <https://proceedings.mlr.press/v75/fefferman18a.html>.
- Christopher Genovese, Marco Perone-Pacifico, Isabella Verdinelli, and Larry Wasserman. Minimax manifold estimation. *Journal of Machine Learning Research*, 13(May):1263–1291, 2012a.
- Christopher R. Genovese, Marco Perone-Pacifico, Isabella Verdinelli, and Larry Wasserman. Manifold estimation and singular deconvolution under Hausdorff loss. *The Annals of Statistics*, 40(2):941–963, 2012b. doi: 10.1214/12-AOS994.
- Christopher R. Genovese, Marco Perone-Pacifico, Isabella Verdinelli, and Larry Wasserman. Nonparametric ridge estimation. *The Annals of Statistics*, 42(4):1511–1545, 2014. doi: 10.1214/14-AOS1218.
- Hannah Hochgerner, Amit Zeisel, Peter Lönnerberg, and Sten Linnarsson. Conserved properties of dentate gyrus neurogenesis across postnatal development revealed by single-cell RNA sequencing. *Nature Neuroscience*, 21:290–299, 2018. doi: 10.1038/s41593-017-0056-2.
- Partha Niyogi, Stephen Smale, and Shmuel Weinberger. Finding the homology of submanifolds with high confidence from random samples. *Discrete & Computational Geometry*, 39(1-3):419–441, 2008.
- Umut Ozertem and Deniz Erdogmus. Locally defined principal curves and surfaces. *Journal of Machine Learning Research*, 12(34):1249–1286, 2011.
- Nikita Puchkin and Vladimir Spokoiny. Structure-adaptive manifold estimation. *Journal of Machine Learning Research*, 23(40):1–62, 2022. URL <https://www.jmlr.org/papers/v23/21-0338.html>.
- Aad W. van der Vaart and Jon A. Wellner. *Weak Convergence and Empirical Processes*. Springer, New York, 1996.

- Nick Whiteley, Annie Gray, and Patrick Rubin-Delanchy. Statistical exploration of the manifold hypothesis. *Journal of the Royal Statistical Society Series B: Statistical Methodology*, 88(2):353–385, 2026. doi: 10.1093/jrsssb/qkag055.
- Zhigang Yao and Jiaji Su. Manifold fitting: A review of methods and applications. *Harvard Data Science Review*, 8(3), 2026. doi: 10.1162/99608f92.686fa68c. URL <https://hdsr.mitpress.mit.edu/pub/1bucoqwm>.
- Zhigang Yao and Yuqing Xia. Manifold fitting under unbounded noise. *Journal of Machine Learning Research*, 26(45):1–55, 2025. URL <https://www.jmlr.org/papers/v26/21-0039.html>.
- Zhigang Yao, Jiaji Su, Bingjie Li, and Shing-Tung Yau. Manifold fitting, 2023. URL <https://arxiv.org/abs/2304.07680>. arXiv:2304.07680.
- Yikun Zhang and Yen-Chi Chen. Linear convergence of the subspace constrained mean shift algorithm: From Euclidean to directional data. *Information and Inference: A Journal of the IMA*, 12(1):210–311, 2023. doi: 10.1093/imaiai/iaac005.