

Lecture 2: Paths, Cycles, Connectivity

Dr. George Mertzios

`george.mertzios@durham.ac.uk`

Reminder from last lecture

Reminder from last lecture

- A **graph** G is a pair $(V(G), E(G))$, where
 - $V(G)$ is a **nonempty** set of **vertices** (or **nodes**),
 - $E(G)$ is a set of **unordered pairs** uv with $u, v \in V(G)$ and $u \neq v$, called the **edges** of G .

Reminder from last lecture

- A **graph** G is a pair $(V(G), E(G))$, where
 - $V(G)$ is a **nonempty** set of **vertices** (or **nodes**),
 - $E(G)$ is a set of **unordered pairs** uv with $u, v \in V(G)$ and $u \neq v$, called the **edges** of G .
- When G is clear from the context, we write simply V and E .

Reminder from last lecture

- A **graph** G is a pair $(V(G), E(G))$, where
 - $V(G)$ is a **nonempty** set of **vertices** (or **nodes**),
 - $E(G)$ is a set of **unordered pairs** uv with $u, v \in V(G)$ and $u \neq v$, called the **edges** of G .
- When G is clear from the context, we write simply V and E .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.

Reminder from last lecture

- A **graph** G is a pair $(V(G), E(G))$, where
 - $V(G)$ is a **nonempty** set of **vertices** (or **nodes**),
 - $E(G)$ is a set of **unordered pairs** uv with $u, v \in V(G)$ and $u \neq v$, called the **edges** of G .
- When G is clear from the context, we write simply V and E .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.
- Many problems can be modelled using graphs.

Reminder from last lecture

- A **graph** G is a pair $(V(G), E(G))$, where
 - $V(G)$ is a **nonempty** set of **vertices** (or **nodes**),
 - $E(G)$ is a set of **unordered pairs** uv with $u, v \in V(G)$ and $u \neq v$, called the **edges** of G .
- When G is clear from the context, we write simply V and E .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.
- Many problems can be modelled using graphs.
- We can draw graphs:
 - represent each vertex u by a point, and
 - each edge uv by a line or curve connecting u and v .

Reminder from last lecture

- A **graph** G is a pair $(V(G), E(G))$, where
 - $V(G)$ is a **nonempty** set of **vertices** (or **nodes**),
 - $E(G)$ is a set of **unordered pairs** uv with $u, v \in V(G)$ and $u \neq v$, called the **edges** of G .
- When G is clear from the context, we write simply V and E .
- Graphs can be directed, multi-, pseudo-, weighted. By default they are not.
- Many problems can be modelled using graphs.
- We can draw graphs:
 - represent each vertex u by a point, and
 - each edge uv by a line or curve connecting u and v .
- Paths, cycles, bipartite graphs, complete graphs, hypercubes

Contents for today's lecture

- Paths and directed paths;
- The shortest path problem;
- Connectivity and connected components;
- Eulerian and Hamiltonian cycles;
- Examples and exercises.

Walks, paths, cycles, and distances

Walks, paths, cycles, and distances

- A **walk** in a graph G is a sequence of edges $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$. In this case we also say that v_0, v_1, \dots, v_n is a walk in G .
- A walk v_0, v_1, \dots, v_n in G is a **path** if all v_i 's are distinct. In this case we also say that v_0, v_1, \dots, v_n is a path in G .

Walks, paths, cycles, and distances


- A **walk** in a graph G is a sequence of edges $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$. In this case we also say that v_0, v_1, \dots, v_n is a walk in G .
- A walk v_0, v_1, \dots, v_n in G is a **path** if all v_i 's are distinct. In this case we also say that v_0, v_1, \dots, v_n is a path in G .
- A walk v_0, v_1, \dots, v_n with $v_0 = v_n$ is called a **circuit** or **closed walk**.

Walks, paths, cycles, and distances

- A **walk** in a graph G is a sequence of edges $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$. In this case we also say that v_0, v_1, \dots, v_n is a walk in G .
- A walk v_0, v_1, \dots, v_n in G is a **path** if all v_i 's are distinct. In this case we also say that v_0, v_1, \dots, v_n is a path in G .
- A walk v_0, v_1, \dots, v_n with $v_0 = v_n$ is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all v_i 's in it are distinct except $v_0 = v_n$.

Walks, paths, cycles, and distances

- A **walk** in a graph G is a sequence of edges $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$. In this case we also say that v_0, v_1, \dots, v_n is a walk in G .
- A walk v_0, v_1, \dots, v_n in G is a **path** if all v_i 's are distinct. In this case we also say that v_0, v_1, \dots, v_n is a path in G .
- A walk v_0, v_1, \dots, v_n with $v_0 = v_n$ is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all v_i 's in it are distinct except $v_0 = v_n$.
- If G is a directed graph then the directed paths and directed cycles are defined in a natural way, with each edge being directed from v_i to v_{i+1} .



Walks, paths, cycles, and distances

- A **walk** in a graph G is a sequence of edges $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$. In this case we also say that v_0, v_1, \dots, v_n is a walk in G .
- A walk v_0, v_1, \dots, v_n in G is a **path** if all v_i 's are distinct. In this case we also say that v_0, v_1, \dots, v_n is a path in G .
- A walk v_0, v_1, \dots, v_n with $v_0 = v_n$ is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all v_i 's in it are distinct except $v_0 = v_n$.
- If G is a directed graph then the **directed paths** and **directed cycles** are defined in a natural way, with each edge being directed from v_i to v_{i+1} .
- The **length** of a path or a cycle is the number of edges in it.


Walks, paths, cycles, and distances

- A **walk** in a graph G is a sequence of edges $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$. In this case we also say that v_0, v_1, \dots, v_n is a walk in G .
- A walk v_0, v_1, \dots, v_n in G is a **path** if all v_i 's are distinct. In this case we also say that v_0, v_1, \dots, v_n is a path in G .
- A walk v_0, v_1, \dots, v_n with $v_0 = v_n$ is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all v_i 's in it are distinct except $v_0 = v_n$.
- If G is a directed graph then the **directed paths** and **directed cycles** are defined in a natural way, with each edge being directed from v_i to v_{i+1} .
- The **length** of a path or a cycle is the number of edges in it.
- The **distance** between vertices u and v in a graph, denoted $\text{dist}(u, v)$, is the length of a shortest path from u to v if such a path exists, and ∞ otherwise.

Walks, paths, cycles, and distances


- A **walk** in a graph G is a sequence of edges $v_0v_1, v_1v_2, v_2v_3, \dots, v_{n-1}v_n$. In this case we also say that v_0, v_1, \dots, v_n is a walk in G .
- A walk v_0, v_1, \dots, v_n in G is a **path** if all v_i 's are distinct. In this case we also say that v_0, v_1, \dots, v_n is a path in G .
- A walk v_0, v_1, \dots, v_n with $v_0 = v_n$ is called a **circuit** or **closed walk**.
- A closed walk is a **cycle** (or **simple circuit**) if all v_i 's in it are distinct except $v_0 = v_n$.
- If G is a directed graph then the **directed paths** and **directed cycles** are defined in a natural way, with each edge being directed from v_i to v_{i+1} .
- The **length** of a path or a cycle is the number of edges in it.
- The **distance** between vertices u and v in a graph, denoted $dist(u, v)$, is the length of a shortest path from u to v if such a path exists, and ∞ otherwise.
- The **diameter** of a graph is the largest distance between two vertices in it

Exercise




- Does this graph contain a path of length 7?

Exercise




- Does this graph contain a path of length 7?

Exercise




- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?

Exercise




- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?

Exercise




- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?

Exercise



- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?

Exercise



- Does this graph contain a path of length 7?
- Does it contain a cycle of length 7?
- What is the distance from 2 to 6?
- What is the diameter of this graph?

The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted



The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices.

The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices. (Can you put this in plain words?)

The acquaintance graph and six degrees of separation

The **acquaintance graph**:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices. (Can you put this in plain words?)
- This is known as the “small world phenomenon”.

The acquaintance graph and six degrees of separation

The acquaintance graph:

- The vertices are all people
- There is an edge between two of them if they are acquainted



- Many sociologists conjectured/observed that this graph has diameter 6, after removing a tiny fraction of vertices. (Can you put this in plain words?)
- This is known as the “small world phenomenon”.
- There is a popular play (and a film) based on this, called “Six degrees of separation”.

The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper

The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper



- Paul Erdős (1913-1996) was a great and extremely prolific mathematician.

The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper



- Paul Erdős (1913-1996) was a great and extremely prolific mathematician.
- The Erdős number of a person is the distance to Paul Erdős in the collaboration graph.

The collaboration graph and the Erdős number

The **collaboration graph**:

- The vertices are all people (or only scientists)
- There is an edge between two of them if they have written a joint paper



- Paul Erdős (1913-1996) was a great and extremely prolific mathematician.
- The Erdős number of a person is the distance to Paul Erdős in the collaboration graph.
- Btw, my Erdős number is 2. (Can you put this in plain words?)

The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.

The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.



- Kevin Bacon is a famous Hollywood actor.

The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.



- Kevin Bacon is a famous Hollywood actor.
- The Bacon number of a person is the distance to Kevin Bacon in the Hollywood graph.

The Hollywood graph and the Bacon number

The **Hollywood graph**:

- The vertices are all people (or only actors).
- There is an edge between two of them if they appeared in the same film.



- Kevin Bacon is a famous Hollywood actor.
- The Bacon number of a person is the distance to Kevin Bacon in the Hollywood graph.
- About 90% of actors have a Bacon number (i.e. the distance is not ∞)

The Erdős-Bacon number

Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

The Erdős-Bacon number

Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

The Erdős-Bacon number

Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.

The Erdős-Bacon number

Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.



The Erdős-Bacon number

Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.



The Erdős-Bacon number

Definition

For any person, their Erdős-Bacon number is the sum of their Erdős number and their Bacon number.

There are not many people with a small Erdős-Bacon number.

For example, the following people have Erdős-Bacon number 7 or less.



- Btw, my Erdős-Bacon number is ∞ , but I have a colleague who has a co-author (Hubie Chen) with Erdős-Bacon number 5 (3+2)

Shortest-path problems

In a graph (possibly with **edge weights**), the problem of computing a path from a given vertex u (“source”) to a given vertex v (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.

Shortest-path problems


In a graph (possibly with **edge weights**), the problem of computing a path from a given vertex u (“source”) to a given vertex v (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.

We all often (use software applications that) compute such paths. Example?

Shortest-path problems

In a graph (possibly with **edge weights**, the problem of computing a path from a given vertex u (“source”) to a given vertex v (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.


We all often (use software applications that) compute such paths. Example?



Shortest-path problems

In a graph (possibly with **edge weights**, the problem of computing a path from a given vertex u (“source”) to a given vertex v (“target”) with the smallest total length (or weight) is known as the **shortest-path problem**.

We all often (use software applications that) compute such paths. Example?




We will learn about algorithms for the (unweighted) problem in a few lectures.

Connectivity

Definition

A graph $G = (V, E)$ is called **connected** if, between every pair of vertices u, v , there exists at least one path in G .

A **connected component** of G is a **maximal** connected subgraph of G .




- Is this graph connected?

Connectivity

Definition

A graph $G = (V, E)$ is called **connected** if, between every pair of vertices u, v , there exists at least one path in G .

A **connected component** of G is a **maximal** connected subgraph of G .




- What about this graph?

Connectivity

Definition

A graph $G = (V, E)$ is called **connected** if, between every pair of vertices u, v , there exists at least one path in G .

A **connected component** of G is a **maximal** connected subgraph of G .




- What about this graph?

Connectivity

Definition

A graph $G = (V, E)$ is called **connected** if, between every pair of vertices u, v , there exists at least one path in G .

A **connected component** of G is a **maximal** connected subgraph of G .



- How many connected components does this graph have?

Connected components

Theorem

Every graph $G = (V, E)$ contains at least $|V| - |E|$ connected components.

Connected components

Theorem

Every graph $G = (V, E)$ contains at least $|V| - |E|$ connected components.

Proof.

By induction over $m = |E|$.



Connected components

Theorem

Every graph $G = (V, E)$ contains at least $|V| - |E|$ connected components.

Proof.

By induction over $m = |E|$.

Induction basis: $m = 0$. No edges, so only isolated vertices, i.e. $|V|$ connected components.



Connected components

Theorem

Every graph $G = (V, E)$ contains at least $|V| - |E|$ connected components.

Proof.

By induction over $m = |E|$.

Induction basis: $m = 0$. No edges, so only isolated vertices, i.e. $|V|$ connected components.

Induction step: Let $G = (V, E)$ with $|E| = m + 1$. Consider an arbitrary $e \in E$, and define $E' = E \setminus \{e\}$. By induction hypothesis: $G' = (V, E')$ has at least $|V| - |E'| = |V| - m$ connected components. Two cases:

- ① With e , the number of connected components does not change.
- ② With e , the number of connected components ???



Connected components

Theorem

Every graph $G = (V, E)$ contains at least $|V| - |E|$ connected components.

Proof.

By induction over $m = |E|$.

Induction basis: $m = 0$. No edges, so only isolated vertices, i.e. $|V|$ connected components.

Induction step: Let $G = (V, E)$ with $|E| = m + 1$. Consider an arbitrary $e \in E$, and define $E' = E \setminus \{e\}$. By induction hypothesis: $G' = (V, E')$ has at least $|V| - |E'| = |V| - m$ connected components. Two cases:

- 1 With e , the number of connected components does not change.
- 2 With e , the number of connected components decreases by 1.



Connected components

Theorem

Every graph $G = (V, E)$ contains at least $|V| - |E|$ connected components.

Proof.

By induction over $m = |E|$.

Induction basis: $m = 0$. No edges, so only isolated vertices, i.e. $|V|$ connected components.

Induction step: Let $G = (V, E)$ with $|E| = m + 1$. Consider an arbitrary $e \in E$, and define $E' = E \setminus \{e\}$. By induction hypothesis: $G' = (V, E')$ has at least $|V| - |E'| = |V| - m$ connected components. Two cases:

- 1 With e , the number of connected components does not change.
- 2 With e , the number of connected components decreases by 1.

In both cases, G has at least $|V| - m - 1 = |V| - |E|$ connected components. \square

Connected components

Theorem

Every graph $G = (V, E)$ contains at least $|V| - |E|$ connected components.

Proof.

By induction over $m = |E|$.

Induction basis: $m = 0$. No edges, so only isolated vertices, i.e. $|V|$ connected components.

Induction step: Let $G = (V, E)$ with $|E| = m + 1$. Consider an arbitrary $e \in E$, and define $E' = E \setminus \{e\}$. By induction hypothesis: $G' = (V, E')$ has at least $|V| - |E'| = |V| - m$ connected components. Two cases:

- ① With e , the number of connected components does not change.
- ② With e , the number of connected components decreases by 1.

In both cases, G has at least $|V| - m - 1 = |V| - |E|$ connected components. \square

Corollary (useful in various algorithmic proofs)

If $G = (V, E)$ is connected then $|E| \geq |V| - 1$.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

- Proof **by contradiction**: Assume G is not connected, derive a contradiction.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

- Proof **by contradiction**: Assume G is not connected, derive a contradiction.
- Take two vertices u and v in different connected components.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

- Proof **by contradiction**: Assume G is not connected, derive a contradiction.
- Take two vertices u and v in different connected components.
- We have that $\deg(u) \geq (n-1)/2$ and $\deg(v) \geq (n-1)/2$.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

- Proof **by contradiction**: Assume G is not connected, derive a contradiction.
- Take two vertices u and v in different connected components.
- We have that $\deg(u) \geq (n-1)/2$ and $\deg(v) \geq (n-1)/2$.
- The set $N(u)$ contains none of the neighbours of v nor v itself.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

- Proof **by contradiction**: Assume G is not connected, derive a contradiction.
- Take two vertices u and v in different connected components.
- We have that $\deg(u) \geq (n-1)/2$ and $\deg(v) \geq (n-1)/2$.
- The set $N(u)$ contains none of the neighbours of v nor v itself.
- Similarly, $N(v)$ contains none of the neighbours of u nor u itself.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

- Proof **by contradiction**: Assume G is not connected, derive a contradiction.
- Take two vertices u and v in different connected components.
- We have that $\deg(u) \geq (n-1)/2$ and $\deg(v) \geq (n-1)/2$.
- The set $N(u)$ contains none of the neighbours of v nor v itself.
- Similarly, $N(v)$ contains none of the neighbours of u nor u itself.
- Hence G contains at least $\frac{n-1}{2} + 1 + \frac{n-1}{2} + 1 = n + 1$ vertices, a contradiction.

Exercise

Exercise 1: Prove that if G is a graph on n vertices and $\delta(G) \geq (n-1)/2$ then G is connected.

- Proof **by contradiction**: Assume G is not connected, derive a contradiction.
- Take two vertices u and v in different connected components.
- We have that $\deg(u) \geq (n-1)/2$ and $\deg(v) \geq (n-1)/2$.
- The set $N(u)$ contains none of the neighbours of v nor v itself.
- Similarly, $N(v)$ contains none of the neighbours of u nor u itself.
- Hence G contains at least $\frac{n-1}{2} + 1 + \frac{n-1}{2} + 1 = n + 1$ vertices, a contradiction.
- In fact, we even proved more: any two vertices in G are at distance at most 2 (so the diameter of G is at most 2).

Exercise

Exercise 2: Let G be a connected graph. Let v a vertex of degree 1 in G . Prove that the graph $G - v$ (obtained from G by deleting v and its incident edge) is also connected.

Exercise 3: Let $G = (V, E)$ be graph on n vertices. The complement graph \overline{G} is obtained from K_n by deleting all the edges that belong to E . Prove that at least one G and \overline{G} must be connected.


Strong connectivity

Definition

A directed graph G is called (weakly) connected if the graph obtained from G by forgetting directions is connected.

A directed graph is called strongly connected if any two distinct vertices are connected by directed paths in both directions.

A strongly connected component (or simply strong component) of a digraph G is a maximal strongly connected subgraph of G .



- Is this graph strongly connected?


Strong connectivity

Definition

A directed graph G is called (weakly) connected if the graph obtained from G by forgetting directions is connected.

A directed graph is called strongly connected if any two distinct vertices are connected by directed paths in both directions.

A strongly connected component (or simply strong component) of a digraph G is a maximal strongly connected subgraph of G .



- Is this graph strongly connected?

Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC

Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC

Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC
- Wikipedia talk graph: 48% SCC, 100% WCC

Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC
- Wikipedia talk graph: 48% SCC, 100% WCC
- Facebook graph: 100% WCC

Connectivity properties of commercially important graphs

Important large graphs tend to have similar connectivity properties:

See the Stanford Large Networks Dataset Collection:

<https://snap.stanford.edu/data/index.html>

Key: WCC = weakly connected component, SCC = strongly ...

- Web graph: 50% SCC, 91% WCC
- Amazon co-purchase graph: 91% SCC, 100% WCC
- Wikipedia talk graph: 48% SCC, 100% WCC
- Facebook graph: 100% WCC
- Google+ graph: 61% SCC, 100% WCC
- ...

Special circuits/cycles in graphs


- Can we travel along the edges of a given graph G so that we start and finish at the same vertex and traverse **each edge exactly once**?

Special circuits/cycles in graphs

- Can we travel along the edges of a given graph G so that we start and finish at the same vertex and traverse **each edge exactly once**?
 - Such a circuit in G is called a **Eulerian** circuit, after Leonhard Euler (1707-83).


Special circuits/cycles in graphs

- Can we travel along the edges of a given graph G so that we start and finish at the same vertex and traverse **each edge exactly once**?
 - Such a circuit in G is called a **Eulerian** circuit, after Leonhard Euler (1707-83).



Special circuits/cycles in graphs


- Can we travel along the edges of a given graph G so that we start and finish at the same vertex and traverse **each edge exactly once**?
 - Such a circuit in G is called a **Eulerian** circuit, after Leonhard Euler (1707-83).



- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?

Special circuits/cycles in graphs


- Can we travel along the edges of a given graph G so that we start and finish at the same vertex and traverse **each edge exactly once**?
 - Such a circuit in G is called a **Eulerian** circuit, after Leonhard Euler (1707-83).




- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?
 - Such a cycle is called a **Hamiltonian** cycle, after William Hamilton (1805-65).

Special circuits/cycles in graphs

- Can we travel along the edges of a given graph G so that we start and finish at the same vertex and traverse **each edge exactly once**?
 - Such a circuit in G is called a **Eulerian** circuit, after Leonhard Euler (1707-83).




- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?
 - Such a cycle is called a **Hamiltonian** cycle, after William Hamilton (1805-65).




Special circuits/cycles in graphs

- Can we travel along the edges of a given graph G so that we start and finish at the same vertex and traverse **each edge exactly once**?
 - Such a circuit in G is called a **Eulerian** circuit, after Leonhard Euler (1707-83).



- Can we travel along the edges of a given graph so that we start and finish at the same vertex and visit **each vertex exactly once**?
 - Such a cycle is called a **Hamiltonian** cycle, after William Hamilton (1805-65).



- Detecting one of these two types of circuits is easy, while detecting the other is not easy at all. Which is which?