

Simulating the transport and dispersal of volcanic ash clouds with initial conditions created by a 3D plume model

Zhixuan Cao^{1,2} Marcus Bursik³ Qingyuan Yang^{4,5} Abani Patra^{*,1,6}

¹ Mechanical and Aerospace Engineering Department, SUNY Buffalo, Buffalo, NY, USA

² Fluids Business Unit, ANSYS Inc, Lebanon, NH, USA

³ Center for Geohazards Studies, SUNY Buffalo, Buffalo, NY, USA

⁴ Earth Observatory of Singapore, Singapore, Singapore

⁵ The Asian School of the Environment, Nanyang Technological University, Singapore, Singapore

⁶ Data Intensive Studies Center, Tufts University, Medford, MA, USA

Correspondence*:

Abani Patra

abani.patra@tufts.edu

2 ABSTRACT

volcanic ash transport and dispersion (VATD) models simulate atmospheric transport of ash starting from a source originating at the volcano represented by concentration of ash with height. Most VATD models use a source of some prescribed shape calibrated against an empirical expression for the height-mass eruption rate (MER) relation. The actual vertical ash distribution in volcanic plumes usually varies from case to case and have complex dependencies on eruption source parameters and atmospheric conditions. We present here for the first time the use of a three-dimensional (3D) plume model to represent the ash cloud source without any assumption regarding plume geometry. By eliminating assumed behavior associated with a semiempirical plume geometry, the predictive skill of VATD simulations is greatly improved. To date, no VATD simulation adopts initial conditions created from first principles based on a 3D plume simulation. We use our recently developed volcanic plume model based on a 3D smoothed-particle hydrodynamic Lagrangian method, and couple the output to a standard Lagrangian VATD model. We apply the coupled model to the Pinatubo eruption in 1991 to illustrate the effectiveness of the approach. Our investigation reveals that initial particle distribution in the vertical direction has more impact on transport of ash clouds than does the horizontal distribution. Comparison with satellite data indicates that ash particles are concentrated through the depth of the volcanic umbrella cloud, and much lower than the observed maximum plume height.

Keywords: VATD, volcano, 3D plume model, initial conditions, numerical simulation, SPH, Pinatubo, ash transport, ash dispersal

1 INTRODUCTION

Volcanic ash, the fine-grained fraction of tephra, can be widely dispersed to synoptic and global scales, and can lead to a degradation of air quality and pose threats to aviation (Tupper et al., 2007). Identification, tracking and modeling the future movement of volcanic ash help route and schedule flights to avoid ash

24 clouds. Numerical estimation of ash distribution using known and forecast wind fields is necessary if we
25 are to accurately predict ash cloud propagation and spread. Numerous VATD models have been developed
26 by both civil and military aviation, and meteorological agencies to provide forecasts of ash cloud motion
27 (Witham et al., 2007), such as Puff (Tanaka, 1991; Searcy et al., 1998), VAFTAD (Heffter and Stunder,
28 1993), Tephra (Bonadonna et al., 2005), HYSPLIT(Stein et al., 2015; Rolph et al., 2017) and Ash3d
29 (Schwaiger et al., 2012). New techniques have been integrated into VATDs to satisfy increasing demands
30 for different types of output, model accuracy and forecast reliability. This contribution explores a method
31 for creating initial conditions for VATD simulations, which promises to reduce user inversion and hence
32 improve prediction capability.

33 Fero et al. (2009) and Stohl et al. (2011) showed that initial source conditions have significant effects on
34 simulation of volcanic ash transport. Constantinescu et al. (2021) proven that an enhanced initial condition
35 provides an overall better fit of the deposit than models without a disk source, demonstrating the significant
36 impact of initial condition on ash dispersion. Besides location of the eruption vent and timings of the release,
37 traditional VATD simulation requires key global descriptors of the volcanic plume, especially plume height,
38 grain size, eruption duration and mass loading, or alternatively, a mass eruption rate (MER). No matter
39 how these global descriptors are obtained, they are used to furnish the initial conditions for VATDs in the
40 form of a line-source term of a spatio-temporal distribution of particle mass. It is a common practice to
41 pick values for these global descriptors using an empirical expression for the height-MER relation. The
42 values for the descriptors can also be found by parameter calibration (e.g. Fero et al., 2008, 2009; Stohl
43 et al., 2011; Zidikheri et al., 2017). One-dimensional (1D) plume models serve as an alternative option to
44 provide these values. For example, Bursik et al. (2012) used the 1D model puffin (Bursik, 2001) to generate
45 estimates of mass eruption rate and grain size. In some cases, an extra step is adopted to spread ash particles
46 from the line source horizontally, resulting in an initial ash cloud in 3D space. The horizontal spreading
47 depends on an empirical expression as well. For example, the VATD model Puff spreads particles from the
48 line source uniformly in the horizontal direction within a given radius. Considering the complexities of
49 volcanic eruptions, the actual ash distribution in the initial ash cloud should vary from case to case and with
50 time, making it difficult to find one general expression that is suitable for all cases. It is useful therefore to
51 investigate alternative ways for creating initial ash clouds without assumptions regarding plume geometry,
52 or numerical inversion. This provides the major motivation of this paper.

53 VATD models can be categorized into Lagrangian particle tracking and Eulerian advection-diffusion types.
54 Among several available particle tracking models, such as, Hypact (Walko et al., 1995), Puff (Searcy et al.,
55 1998), CANERM (D'amus, 1998), and HYSPLIT (Draxler and Hess, 1998) and advection-diffusion
56 models, such as, Tephra (Bonadonna and Houghton, 2005), Fall3D (Folch et al., 2009), and Ash3D
57 (Schwaiger et al., 2012), we adopt a particle tracking model, Puff, as the primary VATD model. Puff
58 can accept a 3D point cloud description of the starting ash cloud as an initial condition, which makes
59 it technically easier to couple with 3D plume models. Puff initializes a discrete number of tracers that
60 represent a sample of the eruption cloud, and calculates transport, turbulent dispersion, and fallout for
61 each representative tracer. A cylinder emanating vertically from the volcano summit to a specified plume
62 height is the standard approach to provide a simple model of the geometry of a typical ash column. Puff
63 minimally requires horizontal wind field data. The “restart” feature of Puff makes it technically feasible to
64 accommodate the hand-off between a plume simulation and the Puff simulation in terms of time and length
65 scales. We also use the Hybrid Single-Particle Lagrangian Integrated Trajectory model (HYSPLIT) (Stein
66 et al., 2015; Rolph et al., 2017) to better understand simulation results from Puff in this study.

67 Besides parameter calibration, 1D plume models have been used to obtain global descriptors of volcanic
68 plumes. 1D plume models (e.g. Woods, 1988; Bursik, 2001; Mastin, 2007; de'Michieli Vitturi et al., 2015;
69 Folch et al., 2016; Pouget et al., 2016b) solve the equations of motion in 1D using simplifying assumptions,
70 and hence depend on estimation of certain parameters, especially those related to the entrainment of air,
71 which is evaluated based on two coefficients: a coefficient due to turbulence in the rising buoyant jet, and
72 one due to the crosswind field. Different 1D models adopt different entrainment coefficients based on a
73 specific formulation or calibration against well-documented case studies. The feedback from plume to
74 atmosphere is usually ignored in 1D models. While these 1D models generated well-matched results with
75 3D models for plumes that are dominated by wind (often called weak plumes) much greater variability
76 is observed for strong plume scenarios (Bursik et al., 2009; Costa et al., 2016). On the other hand, 3D
77 numerical models for volcanic plumes based on first principles and having few parametrized coefficients
78 (Oberhuber et al., 1998; Neri et al., 2003; Suzuki et al., 2005; Cerminara et al., 2016a; Cao et al., 2018)
79 naturally create a 3D ash cloud, which could serve directly as an initial state of the volcanic material for
80 VATDs. However, there is no VATD simulation using such 3D ash clouds as initial conditions. In this paper,
81 we will carry out VATD simulations using an initial state for the ash cloud based on 3D plume simulations,
82 generated with Plume-SPH (Cao et al., 2018, 2017). The implementation techniques described in this paper
83 can be applied to any combination of VATD model and 3D plume model even though our investigation is
84 based on a specific VATD model and plume model.

85 The 1991 eruption of Pinatubo volcano is used as a case study. Pinatubo erupted between June 12 and 16,
86 1991, after weeks of precursory activity. The climactic phase started on June 15 at 0441 UTC and ended
87 around 1341 UTC (Holasek et al., 1996a). The climactic phase generated voluminous pyroclastic flows,
88 and sent Plinian and co-ignimbrite ash and gas columns to great altitudes (Scott et al., 1996). The evolution
89 of the Pinatubo ash and SO₂louds was tracked using visible (Holasek et al., 1996a), ultraviolet (Total
90 Ozone Mapping Spectrometer; TOMS) (Guo et al., 2004a) and infrared sensors, including the Advanced
91 Very High-Resolution Radiometer (AVHRR) (Guo et al., 2004b). There is sufficient observational data to
92 estimate the eruption conditions for the climactic phase of the eruption (Suzuki and Koyaguchi, 2009). The
93 availability of calibrated eruption conditions and extensive observational data regarding ash cloud transport
94 make the Pinatubo eruption an ideal case study.

2 MATERIALS AND METHODS

95 2.1 Plume-SPH Model

96 Plume-SPH (Cao et al., 2018) is designed to describe an injection of well mixed solid and volcanic gas
97 from a circular vent above a flat surface into a stratified stationary atmosphere. The basic assumptions of
98 the model are:

- 99 1. Molecular viscosity and heat conduction is neglected since turbulent energy and momentum exchange
100 are dominant.
- 101 2. Erupted material consisting of solid with different size and mixture of gases is assumed to be well
102 mixed and behave like a single phase fluid (phase 2) which is valid for eruptions with fine particles and
103 ash.
- 104 3. Air, which is assumed to be well mixed mixture of different gases, is assumed to be another phase
105 (phase 1).
- 106 4. Assume thermodynamic equilibrium and dynamic equilibrium between the two phases. As a result,
107 both phases share the common energy equation and momentum equations.

- 108 5. All other microphysical processes (such as the phase changes of H₂O aggregation, disaggregation,
 109 absorption of gas on the surface of solids, solution of gas into a liquid) and chemical processes are not
 110 considered in this model.
 111 6. The effect of wind, is also not yet considered in this model.

Based on above assumptions, the governing equations of our model are given as:

$$\frac{\partial \rho}{\partial t} + \nabla \cdot (\rho \mathbf{v}) = 0 \quad (1)$$

$$\frac{\partial \rho \xi}{\partial t} + \nabla \cdot (\rho \xi \mathbf{v}) = 0 \quad (2)$$

$$\frac{\partial \rho \mathbf{v}}{\partial t} + \nabla \cdot (\rho \mathbf{v} \mathbf{v} + p \mathbf{I}) = \rho \mathbf{g} \quad (3)$$

$$\frac{\partial \rho E}{\partial t} + \nabla \cdot [(\rho E + p) \mathbf{v}] = \rho \mathbf{g} \cdot \mathbf{v} \quad (4)$$

112 where ρ is the density, \mathbf{v} is the velocity, ξ is the mass fraction of ejected material, \mathbf{g} is the gravitational
 113 acceleration, \mathbf{I} is a unit tensor. $E = e + K$ is the total energy which is a summation of kinetic energy K
 114 and internal energy e . An additional equation is required to close the system. In this model, the equation
 115 for closing the system is the following equation of state (EOS).

$$p = (\gamma_m - 1) \rho e \quad (5)$$

116 where

$$\gamma_m = R_m / C_{vm} + 1 \quad (6)$$

$$R_m = \xi_g R_g + \xi_a R_a \quad (7)$$

$$C_{vm} = \xi_s C_{vs} + \xi_g C_{vg} + \xi_a C_{va} \quad (8)$$

$$\xi_a = 1 - \xi \quad (9)$$

$$\xi_g = \xi \cdot \xi_{g0} \quad (10)$$

$$\xi_s = \xi - \xi_g \quad (11)$$

122 where, C_v is the specific heat with constant volume, R is the gas constant. ξ is the mass fraction of erupted
 123 material. The subscript m represents mixture of ejected material and air, s represents solid portion in the
 124 ejected material, g represents gas portion in the ejected material, a represents air, 0 represents physical
 125 properties of erupted material. ξ_{g0} is the mass fraction of vapor in the erupted material.

126 Three different boundary conditions are applied in this model. At the vent, temperature of erupted
 127 material T , eruption velocity \mathbf{v} , the mass fraction of vapor in erupted material ξ_{g0} and mass discharge
 128 rate \dot{M} are given. The pressure of erupted material p is assumed to be the same as ambient pressure for
 129 pressure-balanced eruption. The radius of vent is determined from ρ , \dot{M} and \mathbf{v} . Non-slip wall boundary
 130 condition is applied to the flat ground, where we enforce the velocity to be zero. With further assumption
 131 that the ground is adiabatic, internal energy flux, which consists of heat flux and energy flux carried by
 132 mass flux, vanishes on the wall boundary. Pressure outlet boundary condition is applied to the surrounding
 133 atmosphere where the pressure is given. Except for the pressure, boundary values for density, velocity, and

134 energy are determined by numerical calculation naturally. The initial condition for Plume-SPH is created
 135 based on atmosphere profile before the eruption.

136 The governing equations, EOS, boundary conditions, and initial conditions establish a complete mathema-
 137 tic model. The model is then discretized using smoothed particle hydrodynamics (SPH) method (Gingold
 138 and Monaghan, 1977). The computational domain is discretized by SPH particles. The current version,
 139 Plume-SPH (Cao et al., 2018) uses two types of SPH particles: 1) particles of phase 1 to represent ambient
 140 air, and 2) particles of phase 2 to represent erupted material. So before the eruption, the computational
 141 domain is fully occupied by particles of phase 1. During the eruption, particles of phase 2 are injected
 142 into the computational domain. The discretized model is then converted into computational software
 143 (Plume-SPH) based on a parallel data management framework (Cao et al., 2017).

144 The input parameters for Plume-SPH include the eruption condition at vent, the material properties, and
 145 atmosphere profile. The eruption parameters, material properties and atmosphere for the “Strong plume–no
 146 wind” case in the recent comparison study on eruptive column models (Costa et al., 2016) are adopted.
 147 Eruption conditions and material properties are listed in Table 1. Note that the density of erupted material
 148 at the vent and radius of the vent can be computed from the given parameters. The eruption pressure
 149 is assumed to be the same as the atmospheric pressure at the vent, hence is not given in the table. The
 150 vertical profiles of atmospheric properties were based on the reanalysis data from European Centre for
 151 Medium-Range Weather Forecasts (ECMWF) for the period corresponding to the climactic phase of the
 152 Pinatubo eruption.

153 Running of Plume-SPH essentially updates physics quantities, such as temperature, velocity, and the
 154 position of SPH particles in each time step. During Plume-SPH simulation, SPH particles of phase 2, which
 155 represent the erupted material, are injected from the eruption vent into the compuation domain with an
 156 initial injection velocity. As they moving upwards, these particles will get mixed with SPH particles of
 157 phase 1, which represent the air, during the whole simulation. Their physics quantities get updated as well.
 158 After the simulation, the computation domain will be filled with SPH particles of both phase 1 and phase 2.
 159 Removing all SPH particles of phase 1 from the computation domain, all of the left SPH particles are these
 160 particles that represent the erupted material, which naturely forms a plume (see Fig. 1).

161 2.2 Puff and Initial Ash Cloud

162 Puff (Tanaka, 1991; Searcy et al., 1998) is a dynamic pollutant tracer model. the model is based on a 3D
 163 Lagrangian form of the fluid mechanics, in which the material transport is represented by the fluid motion,
 164 and diffusion is parameterized by a stochastic process of random walk. Here, the model is constructed by a
 165 sufficiently large number of Lagrangian tracer particles with a random variables $\mathbf{R}_i(t) = (x(t), y(t), z(t))$,
 166 where $i = 1 \sim M$, which represent position vectors of particles from the origin of the ash source at the
 167 time t . M is total number of Lagrangian tracer particles, a sample of all the ash particles.

$$\mathbf{R}_i(t + \Delta t) = \mathbf{R}_i(t) + \mathbf{W}(t)\Delta t + \mathbf{Z}(t)\Delta t + \mathbf{S}_i(t)\Delta t \quad (12)$$

168 Here, \mathbf{W} accounts for local wind advection, \mathbf{Z} is generated by Gaussian random numbers and accounts
 169 for turbulent dispersion, and \mathbf{S} is the terminal gravitational fallout velocity or settling speed, which depends
 170 on a tracer's size.

171 To start a Puff simulation, it requires a collection of tracer particles as the initial condition. The tracer
 172 particles has three basic properties, age, size and position. The age of each particle is the elapsed time
 173 from when they were released from the site. Ash particles in the initial ash cloud has zero ages. Ash size

174 distribution is initialized using a Gaussian shape on a logarithmic scale. According to mean and standard
 175 deviation provided by user, Puff assigns size to each particle. Puff initialize the position of each particle
 176 according to semiempirical expressions. The height of each particle is determined according to specified
 177 distribution from the surface (1000 mbar \cong 0 m) to the top of the plume height, H_{max} , which is given
 178 by user. Puff also supports reading predefined initial ash cloud from a file, contains the coordinates of all
 179 tracer particles.

180 The commonly used vertical particle distribution in Puff is the Poisson distribution. For the Poisson
 181 distribution, the vertical height of ash particles is given by Eq. (13):

$$H = H_{max} - 0.5H_{width}P + H_{width}R \quad (13)$$

182 where P is an integral value drawn from a Poisson distribution of unit mean, R is a uniformly distributed
 183 random number between 0 and 1, H_{max} is the maximum plume height, H_{width} represents an approximate
 184 vertical range over which the ash will be distributed. So for Poisson distribution, user can specify two
 185 parameters, H_{max} and H_{width} . Another commonly used vertical ash distribution in VATD simulation is
 186 Suzuki. For the Suzuki plume shape (Suzuki et al., 1983), the ash mass vertical distribution is assumed to
 187 follow the Eq. (Eq. (14)):

$$Q(z) = Q_m \frac{k^2(1 - z/H_{max})\exp(k(z/H_{max} - 1))}{H_{max} [1 - (1 + k)\exp(-k)]} \quad (14)$$

188 Where Q_m is the total mass of erupted material, k is shape factor, which is an adjustable constant that
 189 controls ash distribution with height. A low value of k gives a roughly uniform distribution of mass with
 190 elevation, while high values of k concentrate mass near the plume top. So for Suzuki distribution, besides
 191 the plume height H_{max} , there is another user specified parameter, k .

192 Puff initialize the horizontal distribution of ash particles according to semiempirical expression as well.
 193 Puff uses a uniformly distributed random process to determine ash particle locations in a circle centered
 194 on the volcano site. The maximum radius (at plume top) at which a particle can be located is given as
 195 “horizontal spread”. The horizontal displacement from a vertical line above the volcano is a random value
 196 within a circle of which the radius equals the “horizontal spread” multiplied by the ratio of the particle
 197 height H to the maximum H_{max} , see Eq. 15. So the resulting shape of the particle distribution within the
 198 plume is an inverted cone in which particles are located directly over the volcano at the lowest level and
 199 extend out further horizontally with increasing plume height.

$$r(H) = r_{max}H/H_{max}R \quad (15)$$

200 where $r(H)$ is the radius of the horizontal circle, whithin which, all particles at the height of H locate.
 201 r_{max} is the horizontal spread. H is the height, R is an uniformly distributed random number between 0 and
 202 1.

203 In summary, particle distributions in the initial ash cloud generated by semiempirical expressions are
 204 controlled by several parameters, for example, H_{max} , H_{width} , and r_{max} if user choose to use semiempirical
 205 expressions Eq. (13) and (15). User can optimize or calibrate these parameters to adjust the initial condition
 206 for Puff so that the simulated results match better with observations. Besides the initial ash cloud, other
 207 input parameters for Puff are diffusivity in the vertical and horizontal directions, start and end time of the
 208 eruption, and eruption duration. When creating initial conditions from output of Plume-SPH, the total

209 number of Lagrangian tracers is the count of all SPH particles of phase 2 in the plume. The same total
210 number of Lagrangian tracers are used when creating initial ash cloud based on semiempirical expressions.
211 All input parameters for Puff are listed in Table 2.

212 2.3 Creation of Initial Ash Cloud From Plume-SPH Output

213 In this study, we convert the output of Plume-SPH into initial ash cloud which serves as the initial
214 condition for Puff. The method proposed consists in generating the initial ash cloud directly from Plume-
215 SPH, foregoing assumptions and estimates, or inverse modeling, regarding ash injection height and timing.
216 The steps to create an initial ash cloud based on the raw output of Plume-SPH are shown in Fig. 1. The
217 initial ash cloud is created from SPH particles of phase 2, which represents the erupted material in the
218 model. After reaching the maximum rise height and starting to spread horizontally, particles of phase 2
219 form an initial umbrella cloud (Fig. 2). The 3D plume simulation is considered complete once the umbrella
220 cloud begins to form. Parcels that will be transported by the ambient wind are those above the “corner”
221 region, where mean plume motion is horizontal rather than vertical. With such consideration, we introduce
222 an elevation threshold, which is the lower elevation limit of the ash that will be transported by the VATD.
223 All SPH particles with elevation lower than the threshold are excluded when creating the initial ash cloud.
224 The inflection point from vertical raising to horizontal spreading happens around 15 km according to the
225 averaged vertical velocity ((d) in Fig. 2) and horizontal velocity ((e) in Fig. 2)). Below this inflection
226 point, particle trajectories are primarily vertical in the stalk-like eruption column. Above this level, particle
227 trajectories are primarily horizontal, as they flow into the umbrella cloud gravity current. So we choose 15
228 km to be the elevation threshold in this study.

229 Considering that SPH particles are only discretization points, each is assigned a grain size according to a
230 given total grain size distribution (TGSD) (Paladio-Melosantos et al., 1996), and a concentration according
231 to the mass and volumetric eruption rate. The Plume-SPH discretization points are thus switched to Puff
232 Lagrangian tracer particles having grain sizes and concentrations. The coordinates of these tracer particles,
233 which are initially in the local Cartesian coordinate system of Plume-SPH, are converted into Puff’s global
234 coordinate system, which is given in terms of (*longitude, latitude, height*). Puff takes the initial ash
235 cloud, consisting of the collection of Lagrangian tracer particles with grain size and concentration, and
236 propagates from time t to time $t + \Delta t$ via solution to an advection/diffusion equation (Eq. (12)).

237 To summarize, there are four steps to create an initial ash cloud from the raw output of Plume-SPH:

- 238 1. filter by SPH particle type to select SPH particles that represent erupted material (phase 2)
- 239 2. filter by a mean velocity threshold to select the upper part (above the “corner” region) dominated by
240 horizontal transport
- 241 3. switch SPH discretization points to Lagrangian tracer particles, by assigning grain size to each particle
- 242 4. convert coordinates of the SPH Lagrangian tracers into the VATDs’ geographic coordinate system

243 The features of the volcanic plume and resulting initial ash cloud used in the case study are shown in Fig. 2.
244 It is important to point out that since both Plume-SPH and Puff are based on the Lagrangian method, there
245 is no extra step of conversion between an Eulerian grid and Lagrangian particles.

246 2.4 Puff Restart

247 The plume and ash transport models are run at different time scales and length scales. The spatial and
248 temporal resolutions of the plume simulations are much finer than those of the ash transport model. It takes
249 tens of minutes (600 s in this case) for the Pinatubo plume to reach a steady height. However the eruption
250 persisted for a few hours (9 hours for the climactic phase of Pinatubo eruption), and it may be necessary to

251 track ash transport for days following an eruption. At present, it is too computationally expensive to do 3D
252 plume simulations of several hours in real time. In order to handle the difference in time scale, we mimic
253 a continuing eruption with intermittent pulses releasing ash particles. In particular, we restart Puff at an
254 interval of 600 s, i.e., the physical time of the plume simulation to reach a steady height. At every Puff
255 restart, we integrate the output of the last Puff simulation and Plume-SPH into a new ash cloud. This new
256 ash cloud serves as a new initial condition with which to restart a Puff simulation. A sketch demonstrating
257 the overall restart process is shown in Fig. (3). The total number of Lagrangian tracer particles used in Puff
258 thus equals the summed number of particles in all releases. The total number of tracer particles is therefore
259 no longer a user-selected parameter. Fero et al. (2008) proposed using more realistic time-dependent plume
260 heights. We do not adopt that strategy here for simplicity, although the idea would be straightforward in
261 execution, given time-dependent eruption conditions.

3 RESULTS

262 Transport of volcanic ash resulting from the Pinatubo eruption on June 15, 1991, is simulated using two
263 different initial conditions. The first type of initial condition is created in a traditional way according to user
264 specified parameters (H_{max} , H_{width} and r_{max}) and the semiempirical plume shape expressions (Eq. (13)
265 and (15)). We use the observed plume height (40 km) and two other parameters assigned semiempirically
266 (Bursik et al., 2012) (TO DO: Add more details). The second type of initial condition is created by the new
267 method proposed in this paper. To create initial conditions using the new method described in this paper,
268 the plume rise is simulated first by Plume-SPH. Then the initial ash cloud is obtained by processing the raw
269 output of Plume-SPH following steps described in Sec. 2.3. Except for initial conditions, the simulation
270 parameters that control the VATD simulation are the same for both simulations. Simulated ash transport
271 results are compared against observations.

272 The simulation results using different initial conditions are compared with TOMS images and AVHRR
273 BTD ash cloud map imagery (Fig. 5). The differences between simulated ash transport by the “Semiempirical
274 initial cloud + Puff” and “Plume-SPH+ Puff” conditions are significant. The simulated ash concentration
275 based on the initial conditions created from Plume-SPH is qualitatively closer to observation than that
276 based on the semiempirical plume shape expression. At 23 hours and 31 hours after the beginning of the
277 climactic phase, the “Plume-SPH + Puff” simulation generates ash footprints that are closer to observations,
278 especially in forecasting the location where there is a high concentration of ash. The “Semiempirical initial
279 cloud + Puff” simulation, however, forecasts an ash distribution more spatially restricted than observation.
280 The location of the high concentration region is far northwest of observation. This is confirmed by the
281 Figure of Merit in Space (FMS), which is 0.148 when for Plume-SPH results while 0.124 for Semiempirical
282 initial cloud. This must mean (TO DO: Add more analysis), Around 55 hours after the beginning of the
283 climactic phase, the disparity between observation and simulation becomes more obvious. Ash in the
284 “Semiempirical initial cloud + Puff” simulation is located too far west of the observation, with a FMS
285 value equals to . The high concentration area of the “Plume-SPH + Puff” simulation, even though closer to
286 observation, has also propagated further down the wind direction than the observation. As expected, the
287 FMSs go down to 1 and 2 for the “Semiempirical initial cloud + Puff” and “Plume-SPH+ Puff” respectively.
288 The decreasing of FMSs as a function of time indicates the accumulation of simulation errors during the
289 simulation.

290 (TO DO: Add more analysis regarding the elevation plot, analysis the results by considering the effect of
291 wind field)

292 Except for the initial conditions, both simulations adopt the same parameters and wind field data. That is
293 to say, the only difference between these two simulations is the initial distribution of ash parcels. The main
294 difference between simulation results from the “Plume-SPH + Puff” and the “Semiempirical initial cloud +
295 Puff” runs can be directly attributed to the initial ash particle distribution, which we discuss in detail in the
296 following section.

297 3.1 Importance of plume height (H_{max})

298 In this section, we discuss the vertical distribution of ash particles in the initial ash cloud. The majority
299 of volcanic ash particles are usually injected at an elevation lower than the plume height. For instance,
300 Holasek et al. (1996a,b) reported the maximum Pinatubo plume height as ~ 39 km while the cloud heights
301 were estimated at $\sim 20 - 25$ km. Self et al. (1996) reported that the maximum plume height could have
302 been > 35 km, but that plume heights were $23 \sim 28$ km after $\sim 15 - 16$ hours. The neutral buoyancy
303 height of the Pinatubo aerosol cloud was estimated with different methods at: $\sim 17 - 26$ km (lidar) by
304 DeFoor et al. (1992), $\sim 20 - 23$ km (balloon) by Deshler et al. (1992), $\sim 17 - 28$ km (lidar) by Jäger (1992),
305 and $\sim 17 - 25$ km (lidar) by Avdyushin et al. (1993). Based on comparison between simulated clouds
306 with early infrared satellite imagery of Pinatubo, Fero et al. (2008) reported that the majority of ash was
307 transported between 16 km and 18 km. These observations make good physical sense, as particles are
308 concentrated near the intrusion height of the umbrella cloud, not near the plume top, because the plume top
309 is due to momentum overshoot. However, the empirical expressions for the height-MER relation, which
310 are commonly adopted to create initial conditions for VATD simulations, tend to place the majority of ash
311 particles closer to the top if one uses observed plume height in the empirical expressions.

312 Here we investigate two commonly used plume shapes, the Poisson (see Eq. (13)) and Suzuki (see Eq.
313 (14)). Particle distributions (in terms of mass percentage or particle number percentage) in the vertical
314 direction in the initial ash cloud are shown in Fig. 6. In that figure, the vertical particle distribution based on
315 Plume-SPH output is compared with the vertical particle distribution created based on semiempirical shape
316 expressions. Both Poisson and Suzuki distributions in Fig. 6 take $H_{max} = 40$ km, which is close to the
317 reported observation of plume height. When adopting the Poisson distribution, see (c) in Fig. 6, the majority
318 of the particles are between 30 km ~ 40 km. Obviously, the Poisson function distributes the majority of ash
319 at a higher elevation than was observed (e.g. Fero et al., 2008). As for the Suzuki distribution, (d) in Fig. 6,
320 the majority of ash particles also occur in a range that is significantly higher than 25 km. Note that in the
321 plot (d), the Suzuki constant k is set to 4, which is commonly used for sub-plinian and plinian eruption
322 columns (Pfeiffer et al., 2005). As for initial ash clouds based on Plume-SPH simulation, most ash particles
323 are distributed between $\sim 17 - 28$ km, which matches well with observations. The plume height is also
324 consistent with observation.

325 For the Poisson distributions, the ash particles cannot be lower without changing the plume height. To
326 distribute the majority of ash particles at a lower elevation, the plume height must be reduced to a value
327 smaller than the observed plume height. Adjusting parameters such as plume height in the empirical
328 expression is actually the traditional source term calibration method. A set of initial ash clouds using
329 different plume heights based on the Poisson distribution is shown in Fig. 7. The plume heights adopted
330 in plume shape expressions are not obtained from any plume model or observation of plume height, but
331 by *a posteriori* calibration to later-observed ash cloud transport heights. For Suzuki, adjusting the Suzuki
332 constant can adjust the distribution of ash particles in vertical direction. As shown in Fig. 6, when k is
333 equals to 1 (see (e)), the majority of ash particles are at a lower elevation than observation. With $k = 3$ and
334 $k = 6$ (figure (g) and (h)), the majority of ash particles are at a higher elevation than observation. When k
335 is set to 2 (see (f)), we can see that the majority of ash particles are roughly distributed in the range 17 -

28 km. But the shape, does not look like a typical plume, as particles are more uniformly distributed in the vertical direction. In addition, the Suzuki constant is different from typical value for sub-plinian and plinian eruption columns (Pfeiffer et al., 2005), means we can not apply previously experiences into the semiempirical expression here.

The ash clouds created by the Poisson distribution with different plume heights are used as initial conditions in Puff simulations, whose results are shown in Fig. 10. Except for the plume height, all other parameters for creating an initial ash cloud are the same as those in Table 2. Of course, the range over which the majority of ash particles is located is lower when using lower plume heights. Figure 10 thus shows that the plume height has a significant influence on the ash transport simulation. When the plume height is 10 km, the high concentration area lags behind that observed. If the designated plume height is 35 km, the high concentration area propagates faster and is more spatially confined than observed. When using a plume height of ~ 41 km, the high concentration area propagates faster and the footprint is narrower than in both observation and “Plume-SPH + Puff” simulation results (see Fig. 5). The simulated high concentration area is closest to “Plume-SPH + Puff” simulation results when assigning a plume height of 30 km. The low-concentration front of the volcanic ash cloud propagates faster than observed, and is located far west of the high concentration areas. A low concentration tail area also appears in the simulation results while there is no such tail in the observed imagery, although this could be the result of imagery calibration or sensitivity. Simulation results based on a calibrated plume height of 30 km show a footprint similar to those of “Plume-SPH + Puff”, although smaller in terms of area. However, the initial ash cloud created by a Poisson distribution with plume height around 20 km generates the best match ash with observation. (TO DO: More analysis here) That is to say, a plume height lower than the real plume height is required by the Poisson plume shape to distribute ash particles at elevations comparable to the “true” ash distribution. Our hypothesis regarding the disparity between the “Semiempirical initial cloud + Puff” simulations and observation is confirmed. Since the initial condition of vertical ash distribution has such a dominant effect on VATD simulation, it is critical for the forecast capability of VATD simulations to explore more accurate and adaptive ways for establishing the initial ash distribution, especially methods that do not rely on *a posteriori* parameter calibration or inversion.

3.2 Effect of Vertical Spread (H_{width})

In the previous section, we explored the effects of adjusting the plume height to change the vertical ash distribution at the source. In this section, we investigate the importance of another parameter in the semiempirical Poisson expression (Eq. (13)). We vary the “vertical spread”, H_{width} , in the range $\sim 3 - 10$ km. A set of initial ash clouds with different vertical spreads are shown in Fig. 7. Except for vertical spread, all other parameters for creating an initial ash cloud are the same as those in Table 2. The vertical width of the region within which the majority of ash particles are located becomes narrower when a smaller value for the vertical spread parameter is used, but changing it has no obvious effect on the height at which the largest fraction of ash particles is injected (essentially the height of a mode in ash distribution). The ash clouds based on different vertical spread parameters are then used as initial conditions in Puff simulations.

The VATD results are shown in Fig. 10. Adjusting of the vertical spread changes particle distribution in the vertical direction, and thus, not surprisingly, affects the VATD simulation results. None of the VATD simulations based on initial ash clouds with vertical spreads equal to 3, 5 or 10 km yield better results than do VATD simulations based on initial conditions created by Plume-SPH (see Fig. 10).

The calibration tests on vertical spread, carried out here, are certainly not exhaustive. One could do a more comprehensive calibration throughout the multi-dimensional parameter space (for Poisson distribution, the parameter space is two dimensional) and find better results. In addition, with a more complicated

380 semiempirical plume shape expression, one could have more control over plume shape and might be able
381 to get an initial condition that yields a more accurate ash transport forecast. However, more complicated
382 and adaptable plume shape expressions imply a higher dimensional parameter space, which requires more
383 effort in calibration, even though the degrees of freedom to adjust plume shape are still limited. Creating
384 initial conditions based on 3D plume simulations is more adaptive to various cases and yields results as
385 good as or better than calibration of the poorly-constrained semiempirical parameter, vertical spread.

386 3.3 Horizontal Ash Distribution

387 The differences between the semiempirical plume particle distribution and actual (or simulated by the
388 3D plume model) are not only in the vertical direction. The importance of the horizontal distance of each
389 initial ash particle from a line extending upward from the volcano is investigated in this section. Puff uses a
390 uniformly distributed random process to determine ash particle locations in a circle centered on the volcano
391 site as described in section 2.2. For the output of Plume-SPH, an effective (maximum) radius is determined
392 according to a given threshold of ash concentration, following Cerminara et al. (2016b). A time averaged,
393 spatial integration of the dynamic 3D flow field is conducted to remove significant fluctuations in time and
394 space. Fig. 8 compares radius of the initial ash clouds created by 3D plume simulations with that assumed
395 in the semiempirical plume shape expression adopted in Puff. It is impossible for the simple, assumed
396 plume shapes to capture the complex and more realistic shapes developed by Plume-SPH. Additional
397 parameterization may generate more reasonable shapes, but these would continue to be *ad hoc*, none would
398 likely to have the potential fidelity of the 3D simulation to reality, and adding a temporally changing
399 distribution would be difficult.

400 Comparison between cross-sectional views of the initial ash clouds is shown in Fig. 9. The cross-sectional
401 view of horizontal particle distribution using the semiempirical method (last figure in Fig. 9) is similar to
402 a cross-sectional view of a simulated 3D plume, in a general sense. However, for simulated 3D plumes,
403 the ash particle distribution in cross section varies with height, which factor would become increasingly
404 important with increasing wind speed, were wind speed to be included in the estimate of initial plume
405 shape. It is difficult for the semiempirical expressions to accommodate such a complex distribution.

406 Despite the obvious difficulty of correctly estimating ash distribution near the vent, or for short propagation
407 times, assigning different values for the horizontal spread has a negligible effect on VATD simulation
408 results at large time. We investigated horizontal spread values between 50 km and 1600 km to create initial
409 ash clouds; all of them generated similar results at large propagation times (> 1 day). Figure 10 shows
410 two different simulation results based on initial ash clouds with horizontal spread equal to 50 km and 600
411 km, respectively. No visible differences are apparent between them. (TO DO: add FMS comparison) This
412 implies that horizontal distribution has a less significant influence on VATD simulation results than does
413 vertical distribution for long distance or large time. Perhaps the most important ramification of this result
414 is that it means the time at which the “handshake” is made between Plume-SPH and the VATD does not
415 affect results significantly for relatively large distances and times.

4 DISCUSSION

416 4.1 Sensitivity of Other Inputs Parameters

417 Besides the initial ash cloud, other parameters for Puff simulations are: horizontal diffusivity, vertical
418 diffusivity, mean grain size, grain size standard deviation and total number of tracers. We present in this
419 subsection informal sensitivity studies on these parameters. We also investigate the influence of eruption
420 duration. The sensitivity analyses will serve as the basis for identifying possible sources of disparities
421 between simulation and observation.

422 Fero et al. (2008) simulated the volcanic ash transportation of Pinatubo eruption in 1991, he carried out
423 systematic sensitivity analysis with respect to input parameters of Puff and found that all other parameters
424 except for the plume height have negligible effect on long term ash transportation of Pinatubo. Inspired by
425 Fero et al. (2008), we carried out similar informal sensitivity analysis with much fewer sample points in
426 the parameter space and get similar results. Among the parameters explored, the eruption duration and
427 beginning time show the most obvious influence on simulated ash distribution, although the effect is still
428 small. To show the differences in an intuitive way, (a) - (c) in Fig. 11 shows simulated ash distribution
429 corresponding to 4.9 hours duration, 9 hours duration and 11 hours duration, respectively. After 72 hours,
430 relative to the simulation starting time, these three cases generate very similar results with tiny visible
431 differences (TO DO: The FMS values are , respectively). Daniele et al. (2009) did sensitivity analysis with
432 respect to the input parameters of Puff on different volcanos and found that for eruptive eruptions, the most
433 dominant factors are the wind field and plume height, while all other input parameters are relatively less
434 important. The significance of wind field has been confirmed by other researchers (Stefanescu et al., 2014,
435 e.g) as well.

436 We conducted several simulations with eruption duration varying in the range of [5, 11] hours with
437 slightly different starting time of climactic phase. Table 3 lists all these simulations. However, only slight
438 visible differences are observed among the simulated ash transport outputs. (TO DO: Add FMS value here).
439 We can see that the eruption duration has negligible effects on long-term ash transport.

440 The new methodology for generating initial ash clouds introduces a new parameter: elevation threshold,
441 which were specified based on averaged vertical velocity and horizontal velocity. We carry out a separate,
442 informal sensitivity analysis on this parameter by varying the elevation threshold from 1.5 km (the height
443 of the vent) to 25 m. The simulated ash distributions show obvious visible differences. Such influence
444 is especially obvious when the elevation threshold is either very high or very low. However, varying the
445 elevation threshold in the range of [12, 18] km generates relatively small differences in ash transport
446 simulation results. Figure 11 (d) and (e) compare the simulated ash distributions corresponding to elevation
447 thresholds of 1.5 km and 15 km. Compared with the ash distribution for a threshold of 15 km, an extra long
448 tail appears when using an elevation threshold of 1.5 km. Adopting lower elevation thresholds adds more
449 tracer particles at lower elevation. As the winds at different elevations are different, the tracers at lower
450 elevations propagate in different directions. The HYSPLIT (Stein et al., 2015; Rolph et al., 2017) forward
451 trajectory tracking starting at 1624 UTC on June 15, indicates that the wind between elevations of 10 km
452 and 15 km blew from north-east to south-west, while winds of higher elevation blew from east to west (see
453 Fig. 4). The results confirms that the elevation threshold is best estimated from the height the inflection
454 point in the vertical distribution (*cf.* Figure 11(d) and (e)).

455 4.2 Other Sources of Disparities

456 The full range of research issues raised by numerical forecasting of volcanic clouds is diverse. We focused
457 on the effect of initial conditions in this paper. During the plume modeling, secondary factors, such as
458 microphysical processes, even though they play lesser roles, likely need to be included to improve accuracy
459 for a particular eruption. Wind field are not considered in current version of Plume-SPH, but for weak
460 plume, wind plays such an important role that it has to be considered in the plume model. In addition,
461 eruption conditions are subject to change with time, even during the climactic phase of an eruption. In the
462 future, time-dependent initial conditions for VATDs can be created from 3D plume simulations based on
463 time-dependent eruption conditions. However, the eruption conditions at the vent are usually inferred from
464 observable information based on 1D plume models. Using 3D plume model won't reduce uncertainties
465 from the eruption conditions.

466 Assumption made in each VATD model is another source of errors. For example, a recent study by Osman
467 et al. (2020) demonstrated the great impact of GSD on modelled ash mass loadings using NAME(Jones
468 et al., 2007) to simulate historical eruptions of various VEIs. Other researchers (Beckett et al., 2015; Scollo
469 et al., 2008) have proven the significance of GSD using other VATD models. Their conclusion, however, is
470 different from our informal sensitivity study and other more comprehensive sensitivity studies using Puff.
471 This might be because of different assumptions made in different VATD models. These assumptions may
472 lead to underestimation or overestimation of certain factor, such as GSD.

473 One implicit assumption in the current method is that ash transportation is dominated by wind advection
474 (the passive dispersion approximation). However, during the growth of volcanic umbrella, the dominant
475 factors are various in different regimes (Pouget et al., 2016a) depending on characteristics of a particular
476 eruption. Webster et al. (2020) suggested that the lateral spread by the intrusive gravity current dominates
477 the transport of the ash cloud in this stage. A few studies by Larry Mastin (Mastin et al., 2014; Mastin
478 and Van Eaton, 2020) also showed that neglecting the umbrella cloud formation for larger eruptions led
479 to significantly different footprints for the resulting VATD fallout maps. Their studies imply that including
480 mapped velocities of the plume as a perturbation on the winds can better capture the radial spreading
481 of umbrella. In current method, the 3D plume model generated initial ash cloud has a radius around 25
482 km. For the Pinatubo 1991 eruption, the passive dispersion approximation can be reasonably applied
483 when radius is greater than 450 km, and can be fully valid only when the radius is greater than 1800 km
484 (Costa et al., 2013). It is computationally too expensive for Plume-SPH model to simulate an volcanic
485 umbrella. An umbrella model, with much coarse resolution, in between plume model and VATD model
486 would presumably better model the whole ash transportation process.

487 Besides the errors from assumptions of the model, errors are also introduced from the reanalysis wind
488 field data and the satellite observations, which are retrievals other than the “truth”.

489 4.3 Summary

490 This paper presents, for the first time, VATD simulations using initial source conditions created by a 3D
491 plume model. Traditional VATD simulations use initial conditions created according to a semiempirical
492 plume shape expression. A case study of the 1991 Pinatubo eruption demonstrates that a 3D plume model
493 can create more realistic initial ash cloud and ash parcel positions, and therefore improve the accuracy of
494 ash transport forecasts. Informal sensitivity analyses suggest that initial conditions, as expressed in the
495 disposition of initial ash parcel positions in the vertical, have a more significant effect on a volcanic ash
496 transport forecast than most other parameters. Comparison of initial ash parcel distributions among the
497 3D plume model, semiempirical expressions, and observations suggests that a major subpopulation of ash
498 parcels should be placed at a much lower elevation than plume height to obtain a better VATD forecast.
499 For the Pinatubo case study, “well-matched” simulation results are observed when using a plume height
500 of around 30 km in semiempirical expressions, which is much lower than the observed plume height of
501 40 km. Comparing the effects of the plume height, vertical spread and horizontal spread shows that ash
502 particle distribution in the vertical direction has the strongest effect on VATD simulation.

503 To summarize, we have presented a novel method for creating *a priori* initial source conditions for
504 VATD simulations. We have shown that it might be possible to obtain initial positions of ash parcels
505 with deterministic forward modeling of the volcanic plume, potentially obviating or lessening the need to
506 attempt to somehow observe initial positions, or *a posteriori* create a history of release heights via inversion
507 (Stohl et al., 2011). Although the method now suffers from the high computational cost associated with
508 3D forward modeling, there is the possibility that in future it might not only help overcome shortcomings
509 of existing methods used to generate *a priori* input parameters, but also overcome the need to do the

510 thousands of runs associated with inverse modeling. In addition, computational cost will continue to
511 diminish as computing speed increases. As they are forward numerical models based on first principles,
512 3D plume models need little if any parameterization, and user intervention should not be required to
513 improve forecast power; no assumption about the initial position of ash parcels is needed. Generation of the
514 initial cloud of ash parcels directly by 3D simulation is potentially adaptable to a variety of volcanic and
515 atmospheric scenarios. In contrast, semiempirical expressions used to determine initial conditions require
516 several parameters to control ash particle distribution along a vertical line source or some simplified shape
517 of the initial ash cloud, making it difficult in some cases to generate initial conditions that closely resemble
518 a complex reality.

CONFLICT OF INTEREST STATEMENT

519 The authors declare that the research was conducted in the absence of any commercial or financial
520 relationships that could be construed as a potential conflict of interest.

AUTHOR CONTRIBUTIONS

521 The idea of using 3D plume model to start a VATD simulation originated from a conservation between
522 AP and MB. ZC carried out the Plume-SPH simulations, Puff simulations, initial results analysis, and
523 prepared the first draft. All authors worked together for further revisions. MB carried out the HYSPLIT
524 simulation. QY post processed the Puff simulation results, overlapped the simulation results with satellite
525 observation, calculated the FMS values. All authors contributed equally to the manuscript writing. AP and
526 MB obtained funding to financially support the work.

FUNDING

527 This work was supported by National Science Foundation awards 1521855, 1621853, and 1821311,
528 1821338 and 2004302 and by the National Research Foundation Singapore and the Singapore Ministry
529 of Education under the Research Centres of Excellence initiative (project number: NRF2018NRF-
530 NSFC003ES-010).

ACKNOWLEDGMENTS

531 We are grateful to the two anonymous reviewers of the paper for their constructive comments and suggestions
532 that improved the paper. Support for the Twentieth Century Reanalysis Project dataset is provided by
533 the U.S. Department of Energy, Office of Science Innovative and Novel Computational Impact on Theory
534 and Experiment (DOE INCITE) program, and Office of Biological and Environmental Research (BER),
535 and by the National Oceanic and Atmospheric Administration Climate Program Office.

REFERENCES

- 536 Avdyushin, S., Tulinov, G., Ivanov, M., Kuzmenko, B., Mezhuev, I., Nardi, B., et al. (1993). 1. spatial and
537 temporal evolution of the optical thickness of the pinatubo aerosol cloud in the northern hemisphere
538 from a network of ship-borne and stationary lidars. *Geophysical research letters* 20, 1963–1966
539 Beckett, F., Witham, C., Hort, M., Stevenson, J., Bonadonna, C., and Millington, S. (2015). Sensitivity of
540 dispersion model forecasts of volcanic ash clouds to the physical characteristics of the particles. *Journal
541 of Geophysical Research: Atmospheres* 120, 11–636
542 Bonadonna, C., Connor, C. B., Houghton, B., Connor, L., Byrne, M., Laing, A., et al. (2005). Probabilistic
543 modeling of tephra dispersal: Hazard assessment of a multiphase rhyolitic eruption at tarawera, new
544 zealand. *Journal of Geophysical Research: Solid Earth* 110

- 545 Bonadonna, C. and Houghton, B. (2005). Total grain-size distribution and volume of tephra-fall deposits.
546 *Bulletin of Volcanology* 67, 441–456
- 547 Bursik, M. (2001). Effect of wind on the rise height of volcanic plumes. *Geophys. Res. Lett* 28, 3621–3624
- 548 Bursik, M., Jones, M., Carn, S., Dean, K., Patra, A., Pavoloni, M., et al. (2012). Estimation and
549 propagation of volcanic source parameter uncertainty in an ash transport and dispersal model: application
550 to the eyjafjallajokull plume of 14–16 april 2010. *Bulletin of volcanology* 74, 2321–2338
- 551 Bursik, M., Kobs, S., Burns, A., Braitseva, O., Bazanova, L., Melekestsev, I., et al. (2009). Volcanic
552 plumes and wind: Jetstream interaction examples and implications for air traffic. *Journal of Volcanology*
553 and *Geothermal Research* 186, 60–67
- 554 Cao, Z., Patra, A., Bursik, M., Pitman, E. B., and Jones, M. (2018). Plume-sph 1.0: a three-dimensional,
555 dusty-gas volcanic plume model based on smoothed particle hydrodynamics. *Geoscientific Model
556 Development* 11, 2691–2715
- 557 Cao, Z., Patra, A., and Jones, M. (2017). Data management and volcano plume simulation with parallel
558 sph method and dynamic halo domains. *Procedia Computer Science* 108, 786–795
- 559 Cerminara, M., Esposti Ongaro, T., and Berselli, L. (2016a). Ashee-1.0: a compressible, equilibrium-
560 eulerian model for volcanic ash plumes. *Geoscientific Model Development* 9, 697–730
- 561 Cerminara, M., Esposti Ongaro, T., and Neri, A. (2016b). Large eddy simulation of gas–particle kinematic
562 decoupling and turbulent entrainment in volcanic plumes. *Journal of Volcanology and Geothermal
563 Research*
- 564 Constantinescu, R., Hopulele-Gligor, A., Connor, C. B., Bonadonna, C., Connor, L. J., Lindsay, J. M.,
565 et al. (2021). The radius of the umbrella cloud helps characterize large explosive volcanic eruptions.
566 *Communications Earth & Environment* 2, 1–8
- 567 Costa, A., Folch, A., and Macedonio, G. (2013). Density-driven transport in the umbrella region of volcanic
568 clouds: Implications for tephra dispersion models. *Geophysical Research Letters* 40, 4823–4827
- 569 Costa, A., Suzuki, Y., Cerminara, M., Devenish, B., Esposti Ongaro, T., Herzog, M., et al. (2016). Results
570 of the eruptive column model inter-comparison study. *Journal of Volcanology and Geothermal Research*
- 571 D'amours, R. (1998). Modeling the etex plume dispersion with the canadian emergency response model.
572 *Atmospheric Environment* 32, 4335–4341
- 573 Daniele, P., Lirer, L., Petrosino, P., Spinelli, N., and Peterson, R. (2009). Applications of the puff model to
574 forecasts of volcanic clouds dispersal from etna and vesuvio. *Computers & Geosciences* 35, 1035–1049
- 575 DeFoor, T. E., Robinson, E., and Ryan, S. (1992). Early lidar observations of the june 1991 pinatubo
576 eruption plume at mauna loa observatory, hawaii. *Geophysical research letters* 19, 187–190
- 577 de'Michieli Vitturi, M., Neri, A., and Barsotti, S. (2015). Plume-mom 1.0: A new integral model of
578 volcanic plumes based on the method of moments. *Geoscientific Model Development* 8, 2447–2463
- 579 Deshler, T., Hofmann, D., Johnson, B., and Rozier, W. (1992). Balloonborne measurements of the pinatubo
580 aerosol size distribution and volatility at laramie, wyoming during the summer of 1991. *Geophysical
581 research letters* 19, 199–202
- 582 Draxler, R. R. and Hess, G. (1998). An overview of the hysplit_4 modelling system for trajectories.
583 *Australian meteorological magazine* 47, 295–308
- 584 Fero, J., Carey, S. N., and Merrill, J. T. (2008). Simulation of the 1980 eruption of mount st. helens using
585 the ash-tracking model puff. *Journal of Volcanology and Geothermal Research* 175, 355–366
- 586 Fero, J., Carey, S. N., and Merrill, J. T. (2009). Simulating the dispersal of tephra from the 1991
587 pinatubo eruption: implications for the formation of widespread ash layers. *Journal of Volcanology and
588 Geothermal Research* 186, 120–131

- 589 Folch, A., Costa, A., and Macedonio, G. (2009). Fall3d: A computational model for transport and
590 deposition of volcanic ash. *Computers & Geosciences* 35, 1334–1342
- 591 Folch, A., Costa, A., and Macedonio, G. (2016). Fplume-1.0: An integral volcanic plume model accounting
592 for ash aggregation. *Geoscientific Model Development* 9, 431
- 593 Gingold, R. A. and Monaghan, J. J. (1977). Smoothed particle hydrodynamics: theory and application to
594 non-spherical stars. *Monthly notices of the royal astronomical society* 181, 375–389
- 595 Guo, S., Bluth, G. J., Rose, W. I., Watson, I. M., and Prata, A. (2004a). Re-evaluation of so₂ release
596 of the 15 june 1991 pinatubo eruption using ultraviolet and infrared satellite sensors. *Geochemistry,*
597 *Geophysics, Geosystems* 5
- 598 Guo, S., Rose, W. I., Bluth, G. J., and Watson, I. M. (2004b). Particles in the great pinatubo volcanic cloud
599 of june 1991: The role of ice. *Geochemistry, Geophysics, Geosystems* 5
- 600 Heffter, J. L. and Stunder, B. J. (1993). Volcanic ash forecast transport and dispersion (vaftad) model.
601 *Weather and forecasting* 8, 533–541
- 602 Holasek, R., Self, S., and Woods, A. (1996a). Satellite observations and interpretation of the 1991 mount
603 pinatubo eruption plumes. *Journal of Geophysical Research: Solid Earth* 101, 27635–27655
- 604 Holasek, R. E., Woods, A. W., and Self, S. (1996b). Experiments on gas-ash separation processes in
605 volcanic umbrella plumes. *Journal of volcanology and geothermal research* 70, 169–181
- 606 Jäger, H. (1992). The pinatubo eruption cloud observed by lidar at garmisch-partenkirchen. *Geophysical*
607 *research letters* 19, 191–194
- 608 Jones, A., Thomson, D., Hort, M., and Devenish, B. (2007). The uk met office's next-generation
609 atmospheric dispersion model, name iii. In *Air pollution modeling and its application XVII* (Springer).
610 580–589
- 611 Mastin, L. G. (2007). A user-friendly one-dimensional model for wet volcanic plumes. *Geochemistry,*
612 *Geophysics, Geosystems* 8
- 613 Mastin, L. G. and Van Eaton, A. R. (2020). Comparing simulations of umbrella-cloud growth and ash
614 transport with observations from pinatubo, kelud, and calbuco volcanoes. *Atmosphere* 11, 1038
- 615 Mastin, L. G., Van Eaton, A. R., and Lowenstern, J. B. (2014). Modeling ash fall distribution from a
616 yellowstone supereruption. *Geochemistry, Geophysics, Geosystems* 15, 3459–3475
- 617 Neri, A., Esposti Ongaro, T., Macedonio, G., and Gidaspow, D. (2003). Multiparticle simulation of
618 collapsing volcanic columns and pyroclastic flow. *Journal of Geophysical Research: Solid Earth*
619 (1978–2012) 108
- 620 Oberhuber, J. M., Herzog, M., Graf, H.-F., and Schwanke, K. (1998). Volcanic plume simulation on large
621 scales. *Journal of Volcanology and Geothermal Research* 87, 29–53
- 622 Osman, S., Beckett, F., Rust, A., and Snee, E. (2020). Sensitivity of volcanic ash dispersion modelling to
623 input grain size distribution based on hydromagmatic and magmatic deposits. *Atmosphere* 11, 567
- 624 Paladio-Melosantos, M. L. O., Solidum, R. U., Scott, W. E., Quiambao, R. B., Umbal, J. V., Rodolfo, K. S.,
625 et al. (1996). Tephra falls of the 1991 eruptions of mount pinatubo. *Fire and mud* 12000, 12030
- 626 Pfeiffer, T., Costa, A., and Macedonio, G. (2005). A model for the numerical simulation of tephra fall
627 deposits. *Journal of Volcanology and Geothermal Research* 140, 273–294
- 628 Pouget, S., Bursik, M., Johnson, C. G., Hogg, A. J., Phillips, J. C., and Sparks, R. S. J. (2016a).
629 Interpretation of umbrella cloud growth and morphology: implications for flow regimes of short-lived
630 and long-lived eruptions. *Bulletin of Volcanology* 78, 1–19
- 631 Pouget, S., Bursik, M., Singla, P., and Singh, T. (2016b). Sensitivity analysis of a one-dimensional model
632 of a volcanic plume with particle fallout and collapse behavior. *Journal of Volcanology and Geothermal*
633 *Research*

- 634 Rolph, G., Stein, A., and Stunder, B. (2017). Real-time environmental applications and display system:
635 Ready. *Environmental Modelling & Software* 95, 210–228
- 636 Schwaiger, H. F., Denlinger, R. P., and Mastin, L. G. (2012). Ash3d: A finite-volume, conservative
637 numerical model for ash transport and tephra deposition. *Journal of Geophysical Research: Solid Earth*
638 117
- 639 Scollo, S., Folch, A., and Costa, A. (2008). A parametric and comparative study of different tephra fallout
640 models. *Journal of Volcanology and Geothermal Research* 176, 199–211
- 641 Scott, W. E., Hoblitt, R. P., Torres, R. C., Self, S., Martinez, M. M. L., and Nillos, T. (1996). Pyroclastic
642 flows of the june 15, 1991, climactic eruption of mount pinatubo. *Fire and Mud: eruptions and lahars of*
643 *Mount Pinatubo, Philippines*, 545–570
- 644 Searcy, C., Dean, K., and Stringer, W. (1998). Puff: A volcanic ash tracking and prediction model. *Journal*
645 *of Volcanology and Geothermal Research* 80, 1–16
- 646 Self, S., Zhao, J.-X., Holasek, R. E., Torres, R. C., and King, A. J. (1996). The atmospheric impact of the
647 1991 mount pinatubo eruption
- 648 Stefanescu, E., Patra, A. K., Bursik, M., Jones, M., Madankan, R., Pitman, E. B., et al. (2014). Fast
649 construction of surrogates for uq central to dddas—application to volcanic ash transport. *Procedia*
650 *Computer Science* 29, 1227–1235
- 651 Stein, A., Draxler, R., Rolph, G., Stunder, B., Cohen, M., and Ngan, F. (2015). Noaa's hysplit atmospheric
652 transport and dispersion modeling system. *Bulletin of the American Meteorological Society* 96, 2059–
653 2077
- 654 Stohl, A., Prata, A., Eckhardt, S., Clarisse, L., Durant, A., Henne, S., et al. (2011). Determination of
655 time-and height-resolved volcanic ash emissions and their use for quantitative ash dispersion modeling:
656 the 2010 eyjafjallajökull eruption. *Atmospheric Chemistry and Physics* 11, 4333–4351
- 657 Suzuki, T. et al. (1983). A theoretical model for dispersion of tephra. *Arc volcanism: physics and tectonics*
658 95, 113
- 659 Suzuki, Y. and Koyaguchi, T. (2009). A three-dimensional numerical simulation of spreading umbrella
660 clouds. *Journal of Geophysical Research: Solid Earth (1978–2012)* 114
- 661 Suzuki, Y. J., Koyaguchi, T., Ogawa, M., and Hachisu, I. (2005). A numerical study of turbulent mixing
662 in eruption clouds using a three-dimensional fluid dynamics model. *Journal of Geophysical Research:*
663 *Solid Earth* 110
- 664 Tanaka, H. (1991). Development of a prediction scheme for the volcanic ash fall from redoubt volcano. In
665 *First Int'l. Symp. on Volcanic Ash and Aviation Safety*. vol. 58
- 666 Tupper, A., Itikarai, I., Richards, M., Prata, F., Carn, S., and Rosenfeld, D. (2007). Facing the challenges of
667 the international airways volcano watch: the 2004/05 eruptions of manam, papua new guinea. *Weather*
668 and Forecasting 22, 175–191
- 669 Walko, R., Tremback, C., and Bell, M. (1995). Hypact: The hybrid particle and concentration transport
670 model. *User's guide*
- 671 Webster, H. N., Devenish, B. J., Mastin, L. G., Thomson, D. J., and Van Eaton, A. R. (2020). Operational
672 modelling of umbrella cloud growth in a lagrangian volcanic ash transport and dispersion model.
673 *Atmosphere* 11, 200
- 674 Witham, C., Hort, M., Potts, R., Servranckx, R., Husson, P., and Bonnardot, F. (2007). Comparison of
675 vaac atmospheric dispersion models using the 1 november 2004 grimsvötn eruption. *Meteorological*
676 *Applications* 14, 27–38
- 677 Woods, A. (1988). The fluid dynamics and thermodynamics of eruption columns. *Bulletin of Volcanology*
678 50, 169–193




Figure 1. Steps to create initial condition for Puff based on raw output of Plume-SPH (Cao et al., 2018). First row: raw output of Plume-SPH. Blue particles are phase 1 (ambient air), red particles are phase 2 (erupted material). Second row: plume after removing SPH particles of phase 1. Picture at right is colored according to the mass fraction of erupted material. Third row: volcanic plume above the “corner” region after cutting off the lower portion. Fourth row: assign sizes to particles converting numerical discretization points into tracers. Fifth row: switch coordinates in local coordinate system into (*longitude, latitude, height*)

679 Zidikheri, M. J., Lucas, C., and Potts, R. J. (2017). Estimation of optimal dispersion model source
 680 parameters using satellite detections of volcanic ash. *Journal of Geophysical Research: Atmospheres*
 681 122, 8207–8232

FIGURE CAPTIONS




Figure 2. Volcano plume from 3D plume model. All particles in the pictures are of phase 2 (particle of phase 1 has been removed) at 600s after eruption, at which time, the plume has already reached the plume height and started spreading radially. (a) is front view of the whole plume. (b) is top view of the plume. (c) is front view of the initial ash cloud, which is essentially a portion of the whole plume whose elevation is higher than a given threshold (in this picture is 15 km). Particles are colored according to mass fraction of erupted material. Red represents high mass fraction while blue represents low mass fraction. (d) is averaged vertical velocity of the plume. At elevations below 15 km, the average vertical velocity decreases. At elevations higher 15 km, the averaged vertical velocity starts increasing. (e) is averaged horizontal velocity of the plume. The averaged horizontal velocity becomes obviously larger when elevation is higher than 15 km. So the reflection point is somewhere around 15 km.

Table 1. List of eruption condition and material properties for plume simulation

Parameters	Units	Plume
Vent velocity	$\text{m} \cdot \text{s}^{-1}$	275
Vent gas mass fraction		0.05
Vent Temperature	K	1053
Vent height	m	1500
Mass discharge rate	$\text{kg} \cdot \text{s}^{-1}$	1.5×10^9
Specific heat of gas at constant volume	$\text{J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$	717
Specific heat of air at constant volume	$\text{J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$	1340
Specific heat of solid	$\text{J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$	1100
Specific heat of gas at constant pressure	$\text{J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$	1000
Specific heat of air at constant pressure	$\text{J} \cdot \text{kg}^{-1} \cdot \text{K}^{-1}$	1810
Density of air at vent height	$\text{kg} \cdot \text{m}^{-3}$	1.104
Pressure at vent height	Pa	84363.4




Figure 3. Mimic successive eruption with intermittent pulsed releasing of ash particles. t_I is the period of pulsing release. t_I equals the physical time of 3D plume simulation.

Table 2. Parameters used in VATD simulation of the climactic phase of Pinatubo eruption on June 15 1991. The first six parameters are used by semiempirical expression to create an initial ash cloud. When creating an initial condition based on the Plume-SPH model, these parameters are extracted from output of Plume-SPH model.

Parameters	Unit	Semiempirical	Plume-SPH
plume height (H_{max})	km	40	-
Horizontal Spread (r_{max})	km	103.808	-
Vertical Spread (H_{width})	km	6.662	-
Plume Shape	-	Poisson	-
Total Ash Particles	-	1768500	1768500
Elevation Threshold	m	-	15000
Horizontal Diffusivity	m^2/s	10000	10000
Vertical Diffusivity	m^2/s	10	10
Grain Size Distribution	-	Gaussian	Gaussian
Mean of Grain Size (Radius)	mm	3.5×10^{-2}	3.5×10^{-2}
Standard Deviation of Grain Size	-	1.0	1.0
Start Time	UT	0441	0441
End time	UT	1341	1341
Simulation Duration	hour	72	72

Table 3. The starting and ending time (UT) for simulating the climactic phase of Pinatubo eruption on June 15 1991. Observed plume height (Holasek et al., 1996a) at different time are also listed in the table.

Eruption duration	4.9 hours	9 hours	10 hours	11.1 hours
Start time	0441	0441	0441	0334
Height at start time	37.5 km	37.5 km	37.5 km	24.5 km
End time	0934	1341	1441	1441
Height at end time	35 km	26.5 km	22.5	22.5 km




Figure 4. Trajectories of particles starting from different heights indicating the wind directions of different evaluations. The trajectories are chosen to start at points that were on the perimeter of the umbrella cloud in x , y and z , and in its center, right before it became affected by the wind to give an idea of the maximum possible spread of the trajectories from that initial condition.




Figure 5. Comparison between “Semiempirical initial cloud + Puff” and “Plume-SPH + Puff”. Pictures to the left are: Puff simulation based on initial condition created according to semiempirical plume shape expression. Pictures to the right are Puff simulation based on initial condition generated by Plume-SPH. TOMS or AVHRR image of Pinatubo ash cloud are overlapped with the simulation results. Ash clouds at different hours after eruption are on different rows. From top to bottom, the images are corresponding to around 23 hours after eruption (UT 199106160341), 31 hours after eruption (UT 199106161141), 55 hours after eruption (UT 199106171141). The observation data on the first row are TOMS ash and ice map. The observation data on the second and third row are AVHRR BTD ash cloud map with atmospheric correction method applied (Guo et al., 2004b).




Figure 6. First row, comparison of particle distribution of initial ash cloud in vertical direction. (a) is corresponding to the initial ash cloud obtained from Plume-SPH output. (b) is (a) truncated by a elevation threshold of 15 km. (c) is for vertical ash distribution based on Poisson distribution (Eq. (13)) with H_{max} equals to 40 km. Another parameter, H_{width} is 6662 m. (d) is corresponding to Suzuki distribution (Eq. (14)) with H_{max} equals to 40 km and k equals to 4(Pfeiffer et al., 2005). The second row, Suzuki distribution with H_{max} equals to 40 km but different values for k . The x axis is the percentage of particle numbers for Plume-SPH and Poisson. For Suzuki the x axis is the mass percentage of erupted material.




Figure 7. Initial particle distribution in vertical direction based on Poisson plume shape (Eq. (13)). The first row varies plume heights. (a) to (d) are corresponding to plume height of 10 km, 20 km, 30 km, 35 km. Another parameter, H_{width} is 6662 m for all four figures in the first row. The second row varies “vertical spread”, H_{width} . (e) to (g) are corresponding to vertical spread of 3 km, 5 km and 10 km. The plume height, H_{max} is set to 40 km for all three figures. The x axis is the percentage of particle numbers. See Fig. 6 for vertical ash distribution of Plume-SPH output.




Figure 8. Comparison between radius of initial ash clouds created by 3D plume model (Plume-SPH) and assumed initial ash cloud shape (Eq. 15) in Puff. The plume shape expression used in Puff defines an inverted cone whose actual shape changes when “horizontal spread” takes different values. $R = 25$ km is corresponding to “horizontal spread” equals to 50 km. $R = 50$ km is corresponding to “horizontal spread” equals to 100 km




Figure 9. Horizontal distribution of ash particles (tracers) on a cross section of initial ash cloud. Puff assumes a randomly uniform distribution of ash particles within a circle, as shown by blue dots in (f). All other figures show the ash particle distribution of initial ash clouds created by Plume-SPH at different elevations.




Figure 10. Ash transport simulated by Puff using different initial ash clouds created according the empirical expressions. Initial ash cloud for (a) to (d) are created according to Poisson distribution with maximum plume heights of 10 km, 20 km, 30 km and 35 km respectively. Initial ash cloud for (e) to (g) are created with vertical spread equals to 3 km, 5 km and 10 km, respectively. Initial ash cloud for (h) - (i) are created with “horizontal spread” equals to 50 km and 600 km respectively. All images are for simulated ash transport around 55 hours after eruption (UT 199106171141). See the observed cloud image in Fig. 5.




Figure 11. Sensitivity of Puff simulation with respect to eruption durations and initial ash cloud cutoff heights .(a) to (c) are simulated ash distribution with different starting and ending time. They corresponding to eruption duration of 4.9 hours, 9 hours and 11.1 hours respectively. Starting and ending time for each case is in Table 3. (d) and (e) are simulated ash distribution taking initial ash clouds obtained using different elevation thresholds (1500 m and 15000 km) from output of Plume-SPH. The starting and ending time are corresponding to 9 hours duration case in Table 3. The contours correspond to ash concentration at 72 hours after eruption.