

***Homework #3***

***ECE661***

***Name: Chengzhang Zhong***

***Email: zhongc@purdue.edu***

# ***Method introduction***

## ***Two step method:***

### ***1. Remove projective distortion***

To compute the homogenous to remove projective distortion, we need to find the vanishing line first. Thus, we first need to find two pairs of lines in the image, which suppose to be parallel in the world coordinate. For example, the four points can be represented as  $p_1, p_2, p_3, p_4$ . Two pairs of parallel lines are  $l_1 = p_1 \times p_2$ ,  $l_2 = p_3 \times p_4$ ,  $l_3 = p_1 \times p_3$  and  $l_4 = p_2 \times p_4$ . Then, the two points A and B on the vanishing line are  $A = l_1 \times l_2$  and  $B = l_3 \times l_4$ . Now the vanishing line can be written as:

$$V = A \times B$$

Where  $V = [k_1 \ k_2 \ k_3]^T$ . Now the Homography that maps this vanishing line to infinity can be written as:

$$H_p = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ k_1 & k_2 & k_3 \end{bmatrix}$$

Apply this Homography to the image, we can get rid of the projective distortion from the image.

### ***2. Remove affine distortion***

To remove affine distortion, two pairs of orthogonal lines are required.

For example, if we have orthogonal lines  $l = [l_1 \ l_2 \ l_3]^T$  and  $m = [m_1 \ m_2 \ m_3]^T$

The angle  $\theta$  between them can be represent as:

$$\cos(\theta) = \frac{l_1 m_1 + l_2 m_2}{\sqrt{(l_1^2 + l_2^2)(m_1^2 + m_2^2)}}$$

Now, represent it by the dual degenerate conic, where:

$$C_{\infty}^* = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 0 \end{bmatrix}$$

$$\cos(\theta) = \frac{l^T C_{\infty}^* m}{\sqrt{(l^T C_{\infty}^* l)(m^T C_{\infty}^* m)}}$$

After further analysis, we have:

$$l^T H_a C_{\infty}^* H_a^T m = [l_1 \ l_2 \ l_3] \begin{bmatrix} A A^T & 0 \\ 0 & 0 \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \\ m_3 \end{bmatrix} = 0$$

where  $H_a$  is the homography we want to use:

$$H_a = \begin{bmatrix} A & 0 \\ 0 & 1 \end{bmatrix}$$

Now, we define  $S = A A^T$ , and the equation equals to:

$$[l_1 \ l_2] \begin{bmatrix} s_{11} & s_{12} \\ s_{12} & s_{22} \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \end{bmatrix} = 0$$

We assume  $s_{22} = 1$ , and then:

$$H_a (l_1 m_1 \quad l_1 m_2 + l_2 m_1) \begin{bmatrix} s_{11} \\ s_{12} \end{bmatrix} = -l_2 m_2$$

The relation between S and A are:

$$A = V D V^T$$

$$S = V D^2 V^T$$

Finally, we can use  $H_a^{-1} H_p$  to remove both projective and affine distortion.

## ***One step method:***

This method is based on orthogonal lines. For example, two orthogonal lines  $l =$

$[l_1 \ l_2 \ l_3]^T$  and  $m = [m_1 \ m_2 \ m_3]^T$ . The format can be written as:

$$[l_1 \ l_2 \ l_3] \begin{bmatrix} a & b/2 & d/2 \\ b/2 & c & e/2 \\ d/2 & e/2 & f \end{bmatrix} \begin{bmatrix} m_1 \\ m_2 \\ m_3 \end{bmatrix} = 0$$

where:

$$C_{\infty}^* = \begin{bmatrix} a & b/2 & d/2 \\ b/2 & c & e/2 \\ d/2 & e/2 & f \end{bmatrix}$$

In here, we assume  $f = 1$  and there left  $a, b, c, d, e$  five unknowns, which requires 5 pairs of orthogonal lines to solve this problem.

After solving  $C_{\infty}^*$ , which is also:

$$C_{\infty}^* = \begin{bmatrix} AA^T & Av \\ v^T A^T & v^T v \end{bmatrix}$$

The Homography needs to use here is:

$$H = \begin{bmatrix} A & 0 \\ v^T & 1 \end{bmatrix}$$

Now, we define  $S = \begin{bmatrix} a & b/2 \\ b/2 & c \end{bmatrix}$ :

$$A = VDV^T$$

$$S = VD^2V^T$$

and  $v^T A^T = [d/2 \quad e/2]$  to solve  $v^T$ .

## ***Compare one step method and two step method:***

For both one step and two step methods, it is hard to implement the procedure of selecting points for orthogonal lines. However, one step method requires more attempts than two step method to generate undistorted image. Thus, I consider two step method is more robust than one step method.

Also, two step methods requires only 4 points as input, and I picked 7 points for the one point method to work. From the size of input perspective, two step method is better than one step method.

Although one step method is less robust than two step method, sometimes it gives a better result. For example, the flatiron image, two step method's result looks more decline than

one step method.

***Basic result demonstration(no lines illustration):***

***See images with lines marked in the next session***

***1. flatiron.jpg***



Figure 1.a original image




Figure 1.b two-step method, projective distortion removed



Figure 1.c two-step method, affine distortion removed





Figure 1.d one-step method

## 2. monalisa.jpg




Figure 2.a original image



Figure 2.b two-step method, projective distortion removed



Figure 2.c two-step method, affine distortion removed





Figure 2.d one-step method

### ***3. wideangle.jpg***



Figure 3.a original image





Figure 3.b two-step method, projective distortion removed



Figure 3.c two-step method, affine distortion removed





Figure 3.d one-step method

## 4. *my first image*



Figure 4.a original image




Figure 4.b two step method, projective distortion removed




Figure 4.c two step method, affine distortion removed






Figure 4.d one-step method

## 5. *my second image*



Figure 5.a original image





Figure 5.b two-step method, projective distortion removed



Figure 5.c two-step method, affine distortion removed



Figure 5.d one-step method



## ***Result demonstration:***

### ***1. flatiron.jpg***



Figure 1.a original image




Figure 1.b two-step method, projective distortion removed




Figure 1.c two-step method, affine distortion removed




Figure 1.d one-step method

## 2. *monalisa.jpg*




Figure 2.a original image





Figure 2.b two-step method, projective distortion removed



Figure 2.c two-step method, affine distortion removed




Figure 2.d one-step method

### ***3. wideangle.jpg***




Figure 3.a original image





Figure 3.b two-step method, projective distortion removed



Figure 3.c two-step method, affine distortion removed






Figure 3.d one-step method

## 4. *my first image*




Figure 4.a original image




Figure 4.b two step method, projective distortion removed




Figure 4.c two step method, affine distortion removed






Figure 4.d one-step method

## 5. *my second image*



Figure 5.a original image





Figure 5.b two-step method, projective distortion removed



Figure 5.c two-step method, affine distortion removed



Figure 5.d one-step method