




Figure 1. Six study sites in Hong Kong with three types (MPA : Cape D'Aguilar [CDA, west]/Tung Ping Chau [TPC, east]; mariculture : Lamma [LM, west]/Sai Kung [SK, east]; sewage : Peng Chau [PC, west]/Center Island [CI, east]) and their environmental impacts. (a) Phase I (Seeding): 28 ARMS deployed (14/site) in MPAs for 12 months; 2 ARMS/site (dashed outlines) sampled. (b) Phase II (resistance): 4 ARMS translocated from MPAs to impacted sites (mariculture: LM/SK, sewage: PC/CI). After 12 months, half of the ARMS (dashed outlines) were sampled. (c) Phase III (resilience): remaining ARMS returned to original MPAs; all ARMS sampled after 6 months. (d) Heatmap of water quality during Phase II, normalized by regional means (east/west). Red/blue indicate values above/below the mean (dissolved oxygen inverted).







Figure 2. PCoA of all ARMS (Bray-Curits metric). Each point represents one community in different sites (MPA: blue, mariculture: yellow, sewage: red) from different phase (phase I: circle, phase II: triangle, phase III: square) across the east (hollow shape) and the west (solid shape). **Succession** was indicated by blue shapes in the top part of the plot where it started in the middle (circles) in the seeding phase, developing sideward through the resistance phase (triangles), and reached to both top corners (squares); communities translocated to the stressed sites during **resistance** phase moved downward with sewage communities on the bottom; after all communities were translocated back to the two MPAs in the **resilience** phase, all communities from the same seeding site clustered together on the two top corners. Note that in the west, phase II MPA communities (green triangles) were clustered together with all the resilience communities (top right) but the pattern is different in the east (top left).

a



b




Figure 3. Percentage total and phyla-specific richness loss (red) and gain (blue) of (a) the resistance communities over the seeding communities and (b) the resilience communities over the resistance communities.




Figure S1. Post-disassembling, individual ARMS plates (top) were rinsed with filtered sea water to remove any motile organism and remaining debris. Dislodged materials were then sieved through a filter set of three (2 mm, 500 µm, 106 µm). Organisms over 2mm were not analyzed in this work. Motile fraction I and II were then preserved in 90% ethanol for later downstream processing. Sessile organisms retained on the plate were then scraped off and put through a blender to homogenize (sessile fraction), and then preserved with 95% ethanol.