

Supplementary Materials for

Low Resistance but High Resilience of Marine Invertebrate Communities to Urban Stressors

Wan & McIlroy *et al.*

Corresponding author: dmbaker@hku.hk

The PDF file includes:

1. Materials and Methods
2. References

1. Materials and Methods

1.1 Site Description

Marine Park:

Tung Ping Chau Marine Park (TPC, 22°32'34.5"N 114°26'13.8"E) is a 270 ha MPA surrounding the northeasternmost island of Hong Kong (HKAFC, 2024b). Sheltered by Mirs Bay and exposed to the oceanic South China Sea, it hosts a robust coral community of over 10 genera of scleractinian coral (Yeung et al., 2021). Cape D'Aguilar Marine Reserve (CDA, 22°12'24.6"N 114°15'24.2"E) is a much smaller MPA of only 20 ha located on the southeast tip of Hong Kong Island. Compared to TPC, it has a higher anthropogenic footprint due to the proximity to urban areas. Regardless, it is rich in biodiversity with different habitats including rocky and cobble beaches, rocky shore, intertidal pools, and coral communities (Xu et al., 2015).

Mariculture:

Sai Kung Tai Tau Chau (SK, 22°22'11.4"N 114°19'26.5"E) and Lamma Lo Tik Wan (LM, 22°13'12.9"N 114°07'39.5"E) are two of the 28 fish culture zones (HKAFC, 2024a) governed by the Hong Kong Agriculture, Fisheries and Conservation Department (HKAFC). Most of the farms are small family-run fish rafts, which collectively produce fewer than 500 tonnes of live fish – accounting for only 2% of local demand (HKAFC, 2024a). Although mariculture is not a major source of seafood, the chemical, nutrient, and bacterial pollution produced by this industry impose considerable stress on the surrounding ecosystems (Lai et al., 2016). Both SK and LM are well sheltered by neighbouring bays and islands. LM on the west is more susceptible to

inconsistent water quality influenced by the Pearl River while SK on the east is supplied by marine water from the South China Sea.

Sewage:

Center Island (CI, 22°26'14.1"N 114°13'16.6"E) is in the heart of Tolo Harbor, a slim coastal inlet of about 52 km² in area with a narrow (~1 km) bottleneck. As a semi-enclosed embayment, water circulation is dictated by tidal movement which makes the area naturally prone to eutrophication (Chau, 2007). From the 80s to the late 90s, secondary treated sewage from an expanding population over 500 thousand was discharged into the harbor. In 1988 alone, there were over 40 episodes of harmful algal blooms recorded in the cove (HKEPD, 2007). Although the government had redirected the sewage discharge away since the late 90s, this area continues to record high levels of ammonia nitrogen, *Escherichia coli*, orthophosphate phosphorus, and biochemical oxygen demand compared to neighboring water bodies (HKEPD, 2023). Peng Chau (PC, 22°17'23.7"N 114°02'03.9"E) is better connected to the open sea by water currents than CI. However, Peng Chau Sewage Treatment Works, a secondary wastewater treatment facility with a design capacity of 3,250 m³/day, discharges into this area via submarine outfall (HKEPD, 2001).

1.2 ARMS processing and molecular workflow

A typical set of data retrieved from an ARMS includes plate photos, taxonomic inventories of motile organisms (>2mm), and molecular data from three bulk fractions (motile 500µm, motile

106µm, sessile). We focused on the data from the bulk samples. There are two motile fractions coming from sieving which followed the standard protocols (NMNH, n.d.-b, Fig. S1). In brief, ARMS and ~60 L of surrounding seawater were collected and transported in a tarpaulin waterproof bag to a disassembly station. After all ARMS plates were disassembled, we filtered all of the seawater together with the sediment and the motile organisms from the disassembly bin with a sieve set of 2mm/500µm and then a set of 500µm/106µm. Organisms were hence separated into two size fractions based on the nominal mesh opening: one retained on the 500 µm sieve (motile 500 µm), and the other retained on the 106 µm sieve (motile 106 µm). Both fractions were rinsed with 90% ethanol before storage in a -20 °C freezer in 95% ethanol. The one sessile fraction came from organic materials that were collected from plate scraping following a different protocol (NMNH, n.d.-b): all sessile organisms were removed from the ARMS plate using a metal scraper, transferred into a blender, and homogenized. Similar to the motile fraction, ~ 15ml sessile fraction was then preserved in 95% ethanol before storing in a -20 °C freezer.

DNA extraction and the downstream molecular workflow were identical for all three fractions except for an initial decanting step for the two motile fractions to remove sediments from the targeted organic matter. After this step, ~10 g material from each sample was weighed out and extracted from using the DNeasy PowerMax Soil Kit (Qiagen) using a slightly modified protocol (NMNH, n.d.-c) with an overnight chemical lysis with proteinase K rather than the standard physical lysis with glass beads aiming to enhance integrity of the DNA fragments. All DNA extracts were then purified using the DNeasy PowerClean Pro Cleanup Kit (Qiagen) with the standard protocol (QIAGEN, 2024).

We then targeted a 313 bp mitochondrial Cytochrome c oxidase I (COI/CO1) region using the “mlCOIintF/jgHCO2198” primer set for PCR amplification (Leray et al., 2013). All DNA samples were amplified in triplicate to minimize PCR bias. Each PCR reaction mixture contained 2 µl 10X Advantage 2 PCR Buffer (Takara Bio, USA), 1 µl of “mlCOIintF” and 1 µl of “jgHCO2198” (10 µM), 1.4 µl of 50X dNTP Mix (10 mM ea.), 0.4 µl 50X Advantage 2 Polymerase Mix, and 14.2 µl of a mixture of DNA and PCR grade of water (20 µl in total). Owing to natural variation of DNA quality, the final DNA input ranged from 10 ng ~50 ng for all samples. All samples were put through the same two-steps denaturing at 95°C for 60s, followed by 16 cycles of 10s denaturing at 95°C -> 30s annealing at 62°C -> 60s elongation at 72°C, then followed by 20 cycles of 10s denaturing at 95°C -> 30s annealing at 46°C -> 60s elongation at 72°C, then an extending elongation of 7min at 72 °C, ending with a storage temperature of 4°C. Triplicate amplifications were pooled as one sample before sequencing by the Genomics Core Centre for PanorOmics Science (CPOS) of the University of Hong Kong.

Two separate sequencing runs were conducted on the illumina MiSeq System using the MiSeq Reagent Kit v2 with 250-bp pair-end readings aiming for a minimum 100k pair-end reads per sample. All motile samples of the baseline/resistance phase ARMS were sequenced in one round (32 samples in total), and the remaining samples were sequenced in another round (52 samples in total). All 84 samples (2 motile and 1 sessile fraction per ARMS, 28 ARMS total) were successfully sequenced.

1.3 Bioinformatic pipeline

Sequences were first imported into QIIME2 (Bolyen et al., 2019) followed by adaptor removal using cutadapt (Martin, 2011) to clean the adaptor sequence used in PCR and NGS sequencing. Subsequently, DADA2 (Callahan et al., 2016) was used to first truncate sequences to 220 bp to filter out low quality reads, and then pair the forward and reverse reads, followed by denoising – a process removing potential errors, artifacts, and chimeras to construct amplicon sequence variants (ASVs). We removed ASVs detected in negative controls unless their sample abundance exceeded control levels by ≥ 10 -fold. Since the targeted region (COI/CO1) is a protein coding region, we further filtered sequences based on amino acid translation. More specifically, MASCE (Ranwez, 2011) was used to first build an amino acid alignment based on the Moorea BIOCDE library. Then the invertebrate protein translation code was used to translate the remaining sequences to amino acid sequences. Afterwards, all translated sequences were aligned to the Moorea BIOCDE alignment removing any sequences that have stop codons, insertions, and more than three deletions (Leray et al., 2013). Lastly, VSEARCH (Rognes et al., 2016) was used to cluster the filtered ASVs into operational taxonomic units (OTUs) based on 97% sequence similarity as proxy for species.

Taxonomic assignment was conducted using BLAST within QIIME2 to a locally curated database (McIlroy et al., 2024). A minimum of 80% over at least 300 bp was considered a match to the phylum level. For all the unassigned OTUs, we performed a second round of BLAST against MIDORI (Leray et al., 2022), a regional COI database following the same criteria.

2. Reference

- Álvarez-Noriega, M., Burgess, S. C., Byers, J. E., Pringle, J. M., Wares, J. P., & Marshall, D. J. (2020). Global biogeography of marine dispersal potential. *Nature Ecology & Evolution*, 4(9), 1196–1203. <https://doi.org/10.1038/s41559-020-1238-y>
- Angelini, C., Altieri, A. H., Silliman, B. R., & Bertness, M. D. (2011). Interactions among foundation species and their consequences for community organization, biodiversity, and conservation. *BioScience*, 61(10), 782–789. <https://doi.org/10.1525/bio.2011.61.10.8>
- Benjamini, Y., & Hochberg, Y. (1995). Controlling the false discovery rate: a practical and powerful approach to multiple testing. *Journal of the Royal statistical society: series B (Methodological)*, 57(1), 289–300. <https://doi.org/10.1111%2Fj.2517-6161.1995.tb02031.x>
- Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J.E., Bittinger, K., Brejnrod, A., Brislawn, C.J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M., Chase, J., Cope, E. K., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. *Nature Biotechnology*, 37, 852–857. <https://doi.org/10.1038/s41587-019-0209-9>
- Browning, T. J., & Moore, C. M. (2023). Global analysis of ocean phytoplankton nutrient limitation reveals high prevalence of co-limitation. *Nature Communications*, 14(1), 5014. <https://doi.org/10.1038/s41467-023-40774-0>
- Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, 13(7), 581–583. <https://doi.org/10.1038/nmeth.3869>
- Chau, K. W. (2007). Integrated water quality management in Tolo Harbour, Hong Kong: A case study. *Journal of Cleaner Production*, 15(16), 1568–1572. <https://doi.org/10.1016/j.jclepro.2006.07.047>
- Cornell, H. V., & Harrison, S. P. (2014). What are species pools and when are they important? *Annual Review of Ecology, Evolution, and Systematics*, 45, 45–67. <https://doi.org/10.1146/annurev-ecolsys-120213-091759>
- Cosby, A. G., Lebakula, V., Smith, C. N., Wanik, D. W., Bergene, K., Rose, A. N., Swanson, D., & Bloom, D. E. (2024). Accelerating growth of human coastal populations at the global and continent levels: 2000–2018. *Scientific Reports*, 14(1), 22489. <https://doi.org/10.1038/s41598-024-73287-x>
- Côté, I. M., & Darling, E. S. (2010). Rethinking ecosystem resilience in the face of climate change. *PLoS biology*, 8(7), e1000438. <https://doi.org/10.1371/journal.pbio.1000438>

- Couëdel, M., Dettai, A., Guillaume, M. M., Bruggemann, F., Bureau, S., Frattini, B., Azie, J.L., & Bruggemann, J. H. (2023). New insights into the diversity of cryptobenthic Cirripectes blennies in the Mascarene Archipelago sampled using Autonomous Reef Monitoring Structures (ARMS). *Ecology and Evolution*, 13(3), e9850. <https://doi.org/10.1111/mec.15692>
- Cowie, R. H., Bouchet, P., & Fontaine, B. (2022). The Sixth Mass Extinction: Fact, fiction or speculation? *Biological Reviews*, 97(2), 640–663. <https://doi.org/10.1111/brv.12816>
- Devol, A. H. (2015). Denitrification, anammox, and N₂ production in marine sediments. *Annual Review of Marine Science*, 7, 403–423. <https://doi.org/10.1146/annurev-marine-010213-135040>
- Donohue, I., Hillebrand, H., Montoya, J. M., Petchey, O. L., Pimm, S. L., Fowler, M. S., Healy, K., Jackson, A. L., Lurgi, M., McClean, D., O'Connor, N. E., O'Gorman, E. J., & Yang, Q. (2016). Navigating the complexity of ecological stability. *Ecology Letters*, 19(9), 1172–1185. <https://doi.org/10.1111/ele.12648>
- Donohue, I., Petchey, O. L., Montoya, J. M., Jackson, A. L., McNally, L., Viana, M., Healy, K., Nessa, M. L., & Emmerson, M. C. (2013). On the dimensionality of ecological stability. *Ecology Letters*, 16(4), 421–429. <https://doi.org/10.1111/ele.12086>
- Eddy, T. D., Lam, V. W. Y., Reygondeau, G., Cisneros-Montemayor, A. M., Greer, K., Palomares, M. L. D., Bruno, J. F., Ota, Y., & Cheung, W. W. L. (2021). Global decline in capacity of coral reefs to provide ecosystem services. *One Earth*, 4(9), 1278–1285. <https://doi.org/10.1016/j.oneear.2021.08.016>
- Fetzer, I., Johst, K., Schäwe, R., Banitz, T., Harms, H., & Chatzinotas, A. (2015). The extent of functional redundancy changes as species' roles shift in different environments. *Proceedings of the National Academy of Sciences*, 112(48), 14888–14893. <https://doi.org/10.1073/pnas.1505587112>
- Free, C. M., Cabral, R. B., Froehlich, H. E., Battista, W., Ojea, E., O'Reilly, E., Palardy, J. E., García Molinos, J., Siegel, K. J., Arnason, R., Juinio-Meñez, M. A., Fabricius, K., Turley, C., & Gaines, S. D. (2022). Expanding ocean food production under climate change. *Nature*, 605(7910), 490–496. <https://doi.org/10.1038/s41586-022-04674-5>
- Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., Howarth, R. W., Cowling, E. B., & Cosby, B. J. (2003). The nitrogen cascade. *BioScience*, 53(4), 341–356. [https://doi.org/10.1641/0006-3568\(2003\)053\[0341:TNC\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2003)053[0341:TNC]2.0.CO;2)
- Garland, G., Banerjee, S., Edlinger, A., Miranda Oliveira, E., Herzog, C., Wittwer, R., Philippot, L., Maestre, F. T., & van der Heijden, M. G. A. (2021). A closer look at the functions behind ecosystem multifunctionality: A review. *Journal of Ecology*, 109(2), 600–613. <https://doi.org/10.1111/1365-2745.13511>

- Google. (2025). Google Earth (Version 10.79.0.2). Retrieved May 8, 2025, from <https://www.google.com/earth/>
- Grall, J., & Chauvaud, L. (2002). Marine eutrophication and benthos: the need for new approaches and concepts. *Global Change Biology*, 8(9), 813-830. <https://doi.org/10.1046/j.1365-2486.2002.00519.x>
- Grimm, V., & Wissel, C. (1997). Babel, or the ecological stability discussions: An inventory and analysis of terminology and a guide for avoiding confusion. *Oecologia*, 109, 323–334. <https://doi.org/10.1007/s004420050090>
- HKAFC. Hong Kong Agriculture, Fisheries and Conservation Department. (2024a). Marine fish culture, pond fish culture and oyster culture. Retrieved December 25, 2024, from https://www.afcd.gov.hk/english/fisheries/fish_aqu/fish_aqu_mpo/fish_aqu_mpo.html
- HKAFC. Hong Kong Agriculture, Fisheries and Conservation Department. (2024b). Designated Marine Parks and Marine Reserve. Retrieved December 25, 2024, from https://www.afcd.gov.hk/english/fisheries/fish_aqu/fish_aqu_mpo/fish_aqu_mpo.html
- HKDS. Hong Kong Drainage Services Department. (2019). Sewerage. Retrieved December 25, 2024, from <https://www.dsd.gov.hk/EN/Sewerage/index.html>
- HKEPD. Hong Kong Environmental Protection Department. (2001). Peng Chau Sewage Treatment Works Upgrade. Retrieved December 26, 2024, from <https://www.epd.gov.hk/eia/register/study/latest/esb-093.htm>
- HKEPD. Hong Kong Environmental Protection Department. (2007). 20 Years of Marine Water Quality Monitoring in Hong Kong. Retrieved December 26, 2024, from <https://www.epd.gov.hk/epd/sites/default/files/epd/english/environmentinhk/water/hkwqrc/files/waterquality/annual-report/marinereport2005.pdf>
- HKEPD. Hong Kong Environmental Protection Department. (2023). Marine Water Quality in Hong Kong in 2022. Retrieved September 30, 2024, from <https://www.epd.gov.hk/epd/sites/default/files/epd/english/environmentinhk/water/hkwqrc/files/waterquality/annual-report/marinereport2022.pdf>
- HKEPD. Hong Kong Environmental Protection Department. (2024). Marine Water Quality Data. Retrieved September 30, 2024, from <https://cd.epic.epd.gov.hk/EPICRIVER/marine/?lang=en>
- Hock, K., Wolff, N. H., Ortiz, J. C., Condie, S. A., Anthony, K. R., Blackwell, P. G., & Mumby, P. J. (2017). Connectivity and systemic resilience of the Great Barrier Reef. *PLoS biology*, 15(11), e2003355. <https://doi.org/10.1371/journal.pbio.2003355>
- Hocke, K. (2023). Oxygen in the Earth System. *Oxygen*, 3(3), 287–299. <https://doi.org/10.3390/oxygen3030019>

- Hooper, D. U., Chapin, F. S., Ewel, J. J., Hector, A., Inchausti, P., Lavorel, S., Lawton, J. H., Lodge, D. M., Loreau, M., Naeem, S., Schmid, B., Setälä, H., Symstad, A. J., Vandermeer, J., & Wardle, D. A. (2005). Effects of biodiversity on ecosystem functioning: A consensus of current knowledge. *Ecological Monographs*, 75(1), 3–35. <https://doi.org/10.1890/04-0922>
- Jones, H. P., Jones, P. C., Barbier, E. B., Blackburn, R. C., Rey Benayas, J. M., Holl, K. D., ... & Mateos, D. M. (2018). Restoration and repair of Earth's damaged ecosystems. *Proceedings of the Royal Society B: Biological Sciences*, 285(1873), 20172577. <https://doi.org/10.1098/rspb.2017.2577>
- Keck, F., Peller, T., Alther, R., Barouillet, C., Blackman, R., Capo, E., ... & Altermatt, F. (2025). The global human impact on biodiversity. *Nature*, 1-6. <https://doi.org/10.1038/s41586-025-08752-2>
- Kraft, N. J. B., Adler, P. B., Godoy, O., James, E. C., Fuller, S., & Levine, J. M. (2015). Community assembly, coexistence and the environmental filtering metaphor. *Functional Ecology*, 29(5), 592–599. <https://doi.org/10.1111/1365-2435.12345>
- Krause-Jensen, D., Markager, S., & Dalsgaard, T. (2012). Benthic and pelagic primary production in different nutrient regimes. *Estuaries and coasts*, 35(2), 527–545. <https://doi.org/10.1007/s12237-011-9443-1>
- Lai, R. W. S., Perkins, M. J., Ho, K. K. Y., Astudillo, J. C., Yung, M. M. N., Russell, B. D., Williams, G. A., & Leung, K. M. Y. (2016). Hong Kong's marine environments: History, challenges and opportunities. *Regional Studies in Marine Science*, 8, 259–273. <https://doi.org/10.1016/j.rsma.2016.09.001>
- Leray, M., & Knowlton, N. (2015). DNA barcoding and metabarcoding of standardized samples reveal patterns of marine benthic diversity. *Proceedings of the National Academy of Sciences*, 112(7), 2076–2081. <https://doi.org/10.1073/pnas.1424997112>
- Leray, M., Knowlton, N., & Machida, R. J. (2022). MIDORI2: A collection of quality controlled, preformatted, and regularly updated reference databases for taxonomic assignment of eukaryotic mitochondrial sequences. *Environmental DNA*, 4(4), 894–907. <https://doi.org/10.1002/edn3.303>
- Leray, M., Yang, J. Y., Meyer, C. P., Mills, S. C., Agudelo, N., Ranwez, V., Boehm, J. T., & Machida, R. J. (2013). A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: Application for characterizing coral reef fish gut contents. *Frontiers in Zoology*, 10(1), 1–14. <https://doi.org/10.1186/1742-9994-10-34>
- Leung, H. M., Leung, S. K. S., Au, C. K., Cheung, K. C., Wong, Y. K., Leung, A. O. W., & Yung, K. K. L. (2015). Comparative assessment of water quality parameters of mariculture for fish production in Hong Kong Waters. *Marine Pollution Bulletin*, 94(1-2), 318–322. <https://doi.org/10.1016/j.marpolbul.2015.01.028>

- Liess, A., Lange, K., Schulz, F., Piggott, J. J., Matthaei, C. D., & Townsend, C. R. (2009). Light, nutrients and grazing interact to determine diatom species richness via changes to productivity, nutrient state and grazer activity. *Journal of Ecology*, 97(2), 326–336. <https://doi.org/10.1111/j.1365-2745.2008.01463.x>
- Lokatis, S., & Jeschke, J. M. (2022). Urban biotic homogenization: Approaches and knowledge gaps. *Ecological Applications*, 32(8), e2703. <https://doi.org/10.1002/eap.2703>
- Loreau, M., & de Mazancourt, C. (2013). Biodiversity and ecosystem stability: A synthesis of underlying mechanisms. *Ecology Letters*, 16, 106–115. <https://doi.org/10.1111/ele.12073>
- Mace, G. M., Norris, K., & Fitter, A. H. (2012). Biodiversity and ecosystem services: A multilayered relationship. *Trends in Ecology & Evolution*, 27(1), 19–26. <https://doi.org/10.1016/j.tree.2011.08.006>
- Martin, M. (2011). Cutadapt removes adapter sequences from high-throughput sequencing reads. *EMBnet journal*, 17(1), 10–12. <http://dx.doi.org/10.14806/ej.17.1.200>
- McIlroy, S. E., Guibert, I., Archana, A., Chung, W. Y. H., Duffy, J. E., Gotama, R., Hui, J., Knowlton, N., Leray, M., Meyer, C., Panagiotou, G., Paulay, G., Russell, B., Thompson, P. D., & Baker, D. M. (2024). Life goes on: Spatial heterogeneity promotes biodiversity in an urbanized coastal marine ecosystem. *Global Change Biology*, 30(4), e17248. <https://doi.org/10.1111/gcb.17248>
- Ng, T. P., Cheng, M. C., Ho, K. K., Lui, G. C., Leung, K. M., & Williams, G. A. (2017). Hong Kong's rich marine biodiversity: The unseen wealth of South China's megalopolis. *Biodiversity and Conservation*, 26(1), 23–36. <https://doi.org/10.1007/s10531-016-1224-5>
- NMNH. National Museum of Natural History. (n.d.-a). Global ARMS Program. Smithsonian Institution. Retrieved December 26, 2024, from <https://naturalhistory.si.edu/research/global-arms-program>
- NMNH. National Museum of Natural History. (n.d.-b). ARMS Protocols. Smithsonian Institution. Retrieved December 26, 2024, from <https://naturalhistory.si.edu/research/global-arms-program/protocols>
- NMNH. National Museum of Natural History. (n.d.-c). Bulk DNA extraction. Smithsonian National Museum of Natural History. <https://naturalhistory.si.edu/sites/default/files/media/file/arms-11dnaextraction.pdf>
- O'Hara, C. C., Frazier, M., & Halpern, B. S. (2021). At-risk marine biodiversity faces extensive, expanding, and intensifying human impacts. *Science*, 372(6537), 84–87. <https://doi.org/10.1126/science.abe6731>

- Obst, M., Exter, K., Allcock, A. L., Arvanitidis, C., Axberg, A., Bustamante, M., Cancio, I., Carreira-Flores, D., Chatzinikolaou, E., Chatzigeorgiou, G., Christmas, N., Clark, M. S., Comtet, T., Dailianis, T., Davies, N., Deneudt, K., Diaz de Cerio, O., Fortič, A., Gerovasileiou, V., Hablützel, P. I., Keklikoglou, K., Kotoulas, G., Lasota, R., Leite, B. R., Loisel, S., Lévêque, L., Levy, L., Malachowicz, M., Mavrič, B., Meyer, C., Mortelmans, J., Norkko, J., Pade, N., Power, A. M., Ramšak, A., Reiss, H., Solbakken, J., Staehr, P. A., Sundberg, P., Thyrring, J., Troncoso, J. S., Viard, F., Wenne, R., Yperifanou, E. I., Zbawicka, M., & Pavloudi, C. (2020). A marine biodiversity observation network for genetic monitoring of hard-bottom communities (ARMS-MBON). *Frontiers in Marine Science*, 7, 572680. <https://doi.org/10.3389/fmars.2020.572680>
- Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Stevens, M. H. H., Oksanen, M. J., & Suggests, M. A. S. S. (2007). The vegan package. Community Ecology Package, <https://doi.org/10.32614/CRAN.package.vegan>
- Pennekamp, F., Pontarp, M., Tabi, A., Altermatt, F., Alther, R., Choffat, Y., Fronhofer, E. A., Ganesanandamoorthy, P., Garnier, A., Griffiths, J. I., Greene, S., Horgan, K., Massie, T. M., Mächler, E., Palamara, G. M., Seymour, M., & Petchey, O. L. (2018). Biodiversity increases and decreases ecosystem stability. *Nature*, 563(7729), 109–112. <https://doi.org/10.1038/s41586-018-0627-8>
- Pirotta, E., Booth, C. G., Costa, D. P., Fleishman, E., Kraus, S. D., Lusseau, D., Moretti, D., New, L. F., Schick, R. S., Schwars, L. K., Simmons, S. E., Thomas, L., Tyack, P. L., Weise, M. J., Wells, R. S., & Harwood, J. (2018). Understanding the population consequences of disturbance. *Ecology and Evolution*, 8(19), 9934–9946. <https://doi.org/10.1002/ece3.4458>
- QIAGEN. (2024). DNeasy PowerClean Pro Cleanup Quick Start Protocol. QIAGEN. Retrieved December 26, 2024, from <https://www.qiagen.com/us/resources/resourcedetail?id=03d492da-7744-4f92-8e57-2abaca7265e2&lang=en>
- Ranwez, V., Harispe, S., Delsuc, F., & Douzery, E. J. (2011). MACSE: Multiple Alignment of Coding SEquences accounting for frameshifts and stop codons. *PLoS ONE*, 6(9), e22594. <https://doi.org/10.1371/journal.pone.0022594>
- Rockström, J., Steffen, W., Noone, K., Persson, Å., Chapin III, F. S., Lambin, E. F., ... & Foley, J. A. (2009). A safe operating space for humanity. *Nature*, 461(7263), 472–475. <https://doi.org/10.1038/461472a>
- Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: A versatile open source tool for metagenomics. *PeerJ*, 4, e2584. <https://doi.org/10.7717/peerj.2584>
- Sanderman, J., Hengl, T., Fiske, G., Solvik, K., Adame, M. F., Benson, L., Bukoski, J. J., Carnell, P., Cifuentes-Jara, M., Donato, D., Duncan, C., Eid, E. M., Zu ermgassen, P., Lewis, C. J. E., Macreadie, P. I., Glass, L., Gress, S., Jardine, S. L., Jones, T. G., Nsomobo, E. N., Rahman, M. M., Sanders, C. J., Spalding, M., & Landis, E. (2018). A global map of mangrove forest soil

- carbon at 30 m spatial resolution. *Environmental Research Letters*, 13(5), 055002. <https://doi.org/10.1088/1748-9326/aabe1c>
- Snelgrove, P. V. R., Thrush, S. F., Wall, D. H., & Norkko, A. (2014). Real world biodiversity–ecosystem functioning: A seafloor perspective. *Trends in Ecology & Evolution*, 29(7), 398–405. <https://doi.org/10.1016/j.tree.2014.05.002>
- Stein, A., Gerstner, K., & Kreft, H. (2014). Environmental heterogeneity as a universal driver of species richness across taxa, biomes and spatial scales. *Ecology Letters*, 17(7), 866–880. <https://doi.org/10.1111/ele.12277>
- Sutton, L., Mueter, F. J., Bluhm, B. A., & Iken, K. (2021). Environmental filtering influences functional community assembly of epibenthic communities. *Frontiers in Marine Science*, 8, 736917. <https://doi.org/10.3389/fmars.2021.736917>
- Thébault, E., & Loreau, M. (2005). Trophic interactions and the relationship between species diversity and ecosystem stability. *The American Naturalist*, 166(4), E95–E114. <https://doi.org/10.1086/444403>
- Tilman, D., & Isbell, F. (2015). Recovery as nitrogen declines. *Nature*, 528(7582), 336–337. <https://doi.org/10.1038/nature16320>
- Van den Bulcke, L., De Backer, A., Wittoeck, J., Beentjes, K., Maes, S., Christodoulou, M., Arbizu, P. M., Sapkota, R., Van der Hoorn, B., Winding, A., Hostens, K., & Derycke, S. (2023). DNA metabarcoding on repeat: Sequencing data of marine macrobenthos are reproducible and robust across labs and protocols. *Ecological Indicators*, 150, 110207. <https://doi.org/10.1016/j.ecolind.2023.110207>
- Van Meerbeek, K., Jucker, T., & Svenning, J. C. (2021). Unifying the concepts of stability and resilience in ecology. *Journal of Ecology*, 109(9), 3114–3132. <https://doi.org/10.1111/1365-2745.13651>
- Wei, X., Cai, S., Ni, P., & Zhan, W. (2020). Impacts of climate change and human activities on the water discharge and sediment load of the Pearl River, southern China. *Scientific Reports*, 10(1), 16743. <https://doi.org/10.1038/s41598-020-73939-8>
- Weidlich, E. W. A., Nelson, C. R., Maron, J. L., Callaway, R. M., Delory, B. M., & Temperton, V. M. (2021). Priority effects and ecological restoration. *Restoration Ecology*, 29(1), e13317. <https://doi.org/10.1111/rec.13317>
- Wickham, H. (2016). ggplot2: Elegant Graphics for Data Analysis. *Springer Cham*. ISBN 978-3-319-24277-4. <https://doi.org/10.1007/978-3-319-24277-4>
- Wiens, J. J. (2023). How many species are there on Earth? Progress and problems. *PLoS Biology*, 21(11), e3002388. <https://doi.org/10.1371/journal.pbio.3002388>

- Worm, B., Barbier, E. B., Beaumont, N., Duffy, J. E., Folke, C., Halpern, B. S., Jackson, J., B. C., Lotze, H. K., Michell, F., Palumbi, S. R., Sala, E., Selkoe, K. A., Stachowicz, J. J., & Watson, R. (2006). Impacts of biodiversity loss on ocean ecosystem services. *Science*, 314(5800), 787–790. <https://doi.org/10.1126/science.1132294>
- Wu, P. P. Y., Mengersen, K., McMahon, K., Kendrick, G. A., Chartrand, K., York, P. H., ... & Caley, M. J. (2017). Timing anthropogenic stressors to mitigate their impact on marine ecosystem resilience. *Nature communications*, 8(1), 1263. <https://doi.org/10.1038/s41467-017-01306-9>
- Xu, E. G. B., Leung, K. M. Y., Morton, B., & Lee, J. H. W. (2015). An integrated environmental risk assessment and management framework for enhancing the sustainability of marine protected areas: The Cape d'Aguilar Marine Reserve case study in Hong Kong. *Science of the Total Environment*, 505, 269–281. <https://doi.org/10.1016/j.scitotenv.2014.09.088>
- Xu, Q., Yang, X., Yan, Y., Wang, S., Loreau, M., & Jiang, L. (2021). Consistently positive effect of species diversity on ecosystem, but not population, temporal stability. *Ecology Letters*, 24(10), 2256–2266. <https://doi.org/10.1111/ele.13777>
- Yeung, Y. H., Xie, J. Y., Kwok, C. K., Kei, K., Ang Jr, P., Chan, L. L., Dellisanti, W., Cheang, C. C., Chow, W. K., & Qiu, J. W. (2021). Hong Kong's subtropical scleractinian coral communities: Baseline, environmental drivers and management implications. *Marine Pollution Bulletin*, 167, 112289. <https://doi.org/10.1016/j.marpolbul.2021.112289>