

1 Definitions, Classification of PDEs

Big $\mathcal{O}$: $f(t) = \mathcal{O}(g(t))$ as $t \rightarrow t_0$ if $\exists \varepsilon > 0, C > 0$ s.t. $\forall |t - t_0| < \varepsilon, |f(t)| \leq C|g(t)|$
 $\Leftrightarrow \lim_{t \rightarrow t_0} [f(t)/g(t)]$ bdd ps: If not say $t \rightarrow t_0$, assume $t \rightarrow 0$

More on Big $\mathcal{O}$: a. $F(x) = G(x) + \mathcal{O}(H(x))$ if $F(x) - G(x) = \mathcal{O}(H(x))$

b. $F(x) = \mathcal{O}(G(x)) + \mathcal{O}(H(x))$ if $F(x) = \mathcal{O}(\max\{|G(x)|, |H(x)|\})$

c. $F(x, y) = \mathcal{O}(x^m) + \mathcal{O}(y^n)$ if [Given $G(x) = \mathcal{O}(x^{\frac{m}{n}}) \Rightarrow F(x, G(x)) = \mathcal{O}(x^m)$], 且另一项同理

Classification: For $au_{xx} + bu_{xy} + cu_{yy} + du_x + eu_y + fu + g = 0$ ($a, b, c \neq (0, 0, 0)$)

- Hyperbolic:** $b^2 - 4ac > 0$ Admits non-trivial plane wave solutions
- Parabolic:** $b^2 - 4ac = 0$ (对变量 x, y 进行缩放后不会改变 PDE 类型)
- Elliptic:** $b^2 - 4ac < 0$ No non-trivial plane wave solutions

Weighted 2-Norm: $\|\mathbf{u}\|_{2, \Delta x} = \sqrt{(\Delta x)^d} \|u\|_2$ where d s.t. $\mathbf{u} : \mathbb{R}^d \rightarrow \mathbb{R}^n$

- For $d = 1$ and $\Delta x = \frac{D}{N}$, then: $\forall i, u_i = \mathcal{O}((\Delta x)^P) \Rightarrow \|\mathbf{u}\|_{2, \Delta x} = \mathcal{O}((\Delta x)^P)$

- For A symmetric: $\|A^{-1} \mathbf{w}\|_{2, \Delta x} \leq \frac{1}{|\lambda_1|} \|\mathbf{w}\|_{2, \Delta x} \quad 0 < |\lambda_1| \leq \dots \leq |\lambda_n|$

Linear Algebra: Positive Definite: $\lambda_i > 0, \forall i$ Negative Definite: $\lambda_i < 0, \forall i$

Spectral Theorem: If A symmetric, then $A = Q \Lambda Q^T$ where Q orthogonal, Λ diagonal with eigenvalues.

2 Global/Local Error, Convergence, Consistency, Stability

Exact & Approx. Sol: U_n : approx. n -th time-step sol; u_n : exact sol at n -th time-step.

Global Error ($e_n, \mathbf{e}(\Delta t)$): $e_n = U_n - u(t = n\Delta t)$ $\mathbf{e}(\Delta t) = [e_0, e_1, \dots, e_N]^T$

Convergence: A numerical method is convergent if $\lim_{\Delta t \rightarrow 0} \|\mathbf{e}(\Delta t)\| = 0$

If $\lim_{\Delta t \rightarrow 0} \frac{\|\mathbf{e}(\Delta t)\|}{(\Delta t)^P} = C$ (i.e. $\lim_{\Delta t \rightarrow 0} \|\mathbf{e}(\Delta t)\| = \mathcal{O}((\Delta t)^P)$) $\Rightarrow p$ -th order accurate

Local Truncation Error (τ_n): 数值分析的离散迭代公式, 将 U_n 替换为 u_n 后的误差 (LHS - RHS).

- e.g. For $\frac{U_{n+1} - U_n}{\Delta t} = f(U_n)$, $\tau_{n+1} = \frac{u_{n+1} - u_n}{\Delta t} - f(u_n)$ $\tau_0 = 0$

- e.g. 我们可以把线性数值方法写成 $Au = b$, A matrix, $u = [U_1, \dots, U_{N-1}]^T$.

Then: $\tau = A u_{\text{exact}} - b$ where $u_{\text{exact}} = [u_1, \dots, u_{N-1}]^T$, $\tau = [\tau_1, \dots, \tau_{N-1}]^T$

ps: $Ae = -\tau$ where $e = u - u_{\text{exact}}$ Define $\tau(\Delta t) = [\tau_1, \dots, \tau_{N-1}]^T$

Consistency: If $\lim_{\Delta t \rightarrow 0} \|\tau(\Delta t)\| = 0$, the method is consistent.

Linear Equation: Goal: Finally change ODE/PDE into $Au = b$ where A is matrix, $u = [U_1, \dots, U_{N-1}]^T$

Stability: A numerical method is stable if $\exists \Delta x_0$ s.t. For the Linear System $Au = b$:

- $A(\Delta x)$ is invertible $\forall \Delta x < \Delta x_0$
- $\|A^{-1}(\Delta x) \mathbf{w}\| \leq C \|\mathbf{w}\|$ for all $\mathbf{w} \in \mathbb{R}^{N-1}$

Property: Consistent + Stable $\Rightarrow$ Convergent (same norm!) and $\|\mathbf{e}(\Delta t)\| = \mathcal{O}(\|\tau(\Delta t)\|)$

3 Intro & Method of Lines

Taylor Series: $f(t + \Delta t) = f(t) + f'(t)\Delta t + \frac{1}{2}f''(t)(\Delta t)^2 + \dots + \mathcal{O}((\Delta t)^N)$ ps: $u_{n+1} = u(n\Delta t + \Delta t)$

$$f(t) = f(a) + f'(a)(t-a) + \frac{f''(a)}{2!}(t-a)^2 + \dots + \frac{f^{(N-1)}(a)}{(N-1)!}(t-a)^{N-1} + \mathcal{O}((t-a)^N)$$

Euler Methods: Forward: $\frac{du}{dt}|_n \Delta t \approx \frac{U_{n+1} - U_n}{\Delta t}$; Backward: $\frac{du}{dt}|_n \Delta t \approx \frac{U_n - U_{n-1}}{\Delta t}$

Leapfrog: $\frac{du}{dt}|_n \Delta t \approx \frac{U_{n+1} - U_{n-1}}{2\Delta t}$ **Trapezoidal:** $\frac{1}{2} [\frac{du}{dt}|_n \Delta t + \frac{du}{dt}|_{(n+1)\Delta t}] \approx \frac{U_{n+1} - U_n}{\Delta t}$

Centred Difference: $\frac{d^2 u}{dt^2}|_n \Delta t \approx \frac{U_{n-1} - 2U_n + U_{n+1}}{(\Delta t)^2}$

Method of Lines: Using $U_m(t) \approx u(a + m\Delta x, t)$, $U_m^n \approx u(a + m\Delta x, n\Delta t)$ a 为空间起点, 不重要

- For 1D:** a. 先对空间 x 离散化, 使用 U_m 表达, 此时 $\frac{dU}{dt} = \dots$ where $\mathbf{U} = [U_1(t), U_2(t), \dots, U_{M-1}(t)]^T$
b. 对时间 t 使用 ODE 离散化, 使用 U_m^n 表达, 得到 Linear Equation.

- Example: Crank-Nicolson for Heat Eq. $u_t = \kappa u_{xx}$: 对 x 用 centred difference; 对 t 用 trapezoidal:

$$\frac{U_m^{n+1} - U_m^n}{\Delta t} = \kappa \left(\frac{U_{m+1}^{n+1} - 2U_m^{n+1} + U_{m-1}^{n+1}}{(\Delta x)^2} + \frac{U_{m+1}^n - 2U_m^n + U_{m-1}^n}{(\Delta x)^2} \right)$$

Solve it: By iteration: e.g. $A \mathbf{U}^{n+1} = \mathbf{B} \mathbf{U}^n$, where $\mathbf{U}^n = [U_1^n, U_2^n, \dots, U_{M-1}^n]^T$.

Truncation for MoL: 对于 τ_m , 离散 x 后 U_m 替换为 u_m ; 对于 τ_m^n , 再离散 t 后 U_m^n 替换为 u_m^n .

e.g. For Crank-Nicolson above: $\tau_m = \frac{\partial u_m}{\partial t} - \kappa \frac{u_{m+1} - 2u_m + u_{m-1}}{(\Delta x)^2}$

e.g. For Crank-Nicolson above: $\tau_m^n = \frac{u_m^{n+1} - u_m^n}{\Delta t} - \frac{\kappa}{2} (\dots)$

New Norm: The norm $\|\mathbf{u}\| := \max_n \|\mathbf{u}^n\|_{2, \Delta x}$ where $\mathbf{u}^n = [u_1^n, u_2^n, \dots, u_{M-1}^n]^T$

Which means: Infinity norm in time, weighted 2-norm in space.

Stability for MoL:

5 Appendix

Prototype PDEs: Laplace Eq.: $u_{xx} + u_{yy} = 0$ (E) Poisson Eq.: $u_{xx} + u_{yy} = f(x, y)$ (E)

Helmholtz Eq.: $u_{xx} + u_{yy} + \alpha u = 0, \alpha \neq 0$ (E) Heat Eq.: $u_t = \kappa(u_{xx} + u_{yy}), \kappa > 0$ (P)

Wave Eq.: $u_{xx} - u_{yy} = f(x, y)$ (H) Advection Eq.: $u_t + v u_x + w u_y = 0$ (H)

Taylor Series: $f(t + \Delta t) = f(t) + f'(t)\Delta t + \frac{1}{2}f''(t)(\Delta t)^2 + \dots + \mathcal{O}((\Delta t)^N)$ ps: $u_{n+1} = u(n\Delta t + \Delta t)$

$u_{n+1}^n = u_m^n + \Delta t(u_t)_m^n + \frac{(\Delta t)^2}{2}(u_{tt})_m^n + \dots$ where $(u_t)_m^n = \frac{\partial u}{\partial t}(a + m\Delta x, n\Delta t)$

$u_{m+1}^n = u_m^n + \Delta x(u_x)_m^n + \frac{(\Delta x)^2}{2}(u_{xx})_m^n + \dots$ where $(u_x)_m^n = \frac{\partial u}{\partial x}(a + m\Delta x, n\Delta t)$