

## 1 Definitions, Classification of PDEs

### 1.1 Big O, Classification

**Big O:**  $f(t) = \mathcal{O}(g(t))$  as  $t \rightarrow t_0$  if  $\exists \varepsilon > 0, C > 0$  s.t.  $\forall |t - t_0| < \varepsilon, |f(t)| \leq C|g(t)|$   
 $\Leftrightarrow \lim_{t \rightarrow t_0} [f(t)/g(t)]$  bdd ps: If not say  $t \rightarrow t_0$ , assume  $t \rightarrow 0$

**More on Big O:** a.  $F(x) = G(x) + \mathcal{O}(H(x))$  if  $F(x) - G(x) = \mathcal{O}(H(x))$

- b.  $F(x) = \mathcal{O}(G(x)) + \mathcal{O}(H(x))$  if  $F(x) = \mathcal{O}(\max\{|G(x)|, |H(x)|\})$   
 c.  $F(x, y) = \mathcal{O}(x^m) + \mathcal{O}(y^n)$  [if Given  $G(x) = \mathcal{O}(x^{\frac{m}{n}}) \Rightarrow F(x, G(x)) = \mathcal{O}(x^m)$ ], 且另一项同理

**Classification:** For  $a u_{xx} + b u_{xy} + c u_{yy} + d u_x + e u_y + f u + g = 0$  ( $a, b, c \neq (0, 0, 0)$ )

- Hyperbolic:**  $b^2 - 4ac > 0$  Admits non-trivial plane wave solutions (Propagation)
- Parabolic:**  $b^2 - 4ac = 0$  and we can get  $2cd = be$  or  $2ae = bd$
- Elliptic:**  $b^2 - 4ac < 0$  No non-trivial plane wave solutions (No Propagation)

**Remark:** 对变量  $x, y$  进行缩放后不会改变 PDE 类型

### 1.2 Normal

**Weighted 2-Norm:**  $\|\mathbf{u}\|_{2, \Delta x} = \sqrt{(\Delta x)^d \|\mathbf{u}\|_2}$  where  $d$  s.t.  $\mathbf{u} : \mathbb{R}^d \rightarrow \mathbb{R}^n$

- For  $d = 1$  and  $\Delta x = \frac{D}{N}$ , then:  $\forall i, u_i = \mathcal{O}((\Delta x)^p) \Rightarrow \|\mathbf{u}\|_{2, \Delta x} = \mathcal{O}((\Delta x)^p)$
- For  $A$  symmetric:  $\|A^{-1} \mathbf{w}\|_2 \leq \frac{1}{|\lambda_1|} \|\mathbf{w}\|_2$   $0 < |\lambda_1| \leq \dots \leq |\lambda_n|$

**Property:** If  $\|\mathbf{u}\|_2 \leq \|\mathbf{w}\|_2$ , then  $\|\mathbf{u}\|_{2, \Delta x} \leq \|\mathbf{w}\|_{2, \Delta x}$

**New Norm:** The norm  $\|\mathbf{u}\| := \max_n \|\mathbf{u}^n\|_{2, \Delta x}$  where  $\mathbf{u}^n = [u_1^n, u_2^n, \dots, u_{M-1}^n]^\top$   
 Which means: Infinity norm in time, weighted 2-norm in space. 主要用于双变量 PDE

For  $\tau_m^n = \mathcal{O}((\Delta x)^\alpha) + \mathcal{O}((\Delta t)^b)$ ,  $\Delta x = \frac{b-\Delta t}{M-\Delta t}$ ,  $\Delta t = \frac{\tau}{N}$ , assume  $\Delta t = K(\Delta x)^\beta$ , then:  
 $\max_n \|\mathbf{x}^n\|_{2, \Delta x} = \mathcal{O}((\Delta x)^{\min\{\alpha, b\}})$

### 1.3 Linear Algebra

**Definite:** Positive Definite:  $\lambda_i > 0, \forall i$  Negative Definite:  $\lambda_i < 0, \forall i$

**Symmetric:** If  $A$  symmetric, then  $A = Q \Lambda Q^\top$  where  $Q$  orthogonal,  $\Lambda$  diagonal with eigenvalues.

**Normal:** Real matrix  $A$  is normal if  $A A^\top = A^\top A$  The following are properties of it:

- $A$  is orthogonally diagonalizable, i.e.  $A = Q \Lambda Q^\top$  where  $Q$  orthogonal,  $\Lambda$  diagonal with eigenvalues.
- Exists an eigenvectors of  $A$ :  $\{\mathbf{v}_i\}$  s.t.  $\mathbf{v}_i^\top \mathbf{v}_j = \delta_{ij}$  and Symmetric  $\Rightarrow$  Normal

## 2 Global/Local Error, Convergence, Consistency, Stability

**Exact & Approx. Sol:**  $U_n$ : approx.  $n$ -th time-step sol;  $u_n$ : exact sol at  $n$ -th time-step.

**Global Error**  $(e_n, \mathbf{e}(\Delta t))$ :  $e_n = U_n - u(t = n\Delta t)$   $\mathbf{e}(\Delta t) = [e_0, e_1, \dots, e_N]^\top$

**Convergence:** A numerical method is convergent if  $\lim_{\Delta t \rightarrow 0} \|\mathbf{e}(\Delta t)\| = 0$

If  $\lim_{\Delta t \rightarrow 0} \frac{\|\mathbf{e}(\Delta t)\|}{(\Delta t)^p} = C$  (i.e.  $\|\mathbf{e}(\Delta t)\| = \mathcal{O}((\Delta t)^p)$ )  $\Rightarrow$   $p$ -th order accurate

**Local Truncation Error**  $(\tau_n)$ : 数值分析的离散迭代公式, 将  $U_n$  替换为  $u_n$  后的误差 (LHS - RHS).

- e.g. For  $\frac{U_{n+1} - U_n}{\Delta t} = f(U_n)$ ,  $\tau_{n+1} = \frac{U_{n+1} - u_{n+1}}{\Delta t} - f(u_n)$   $\tau_0 = 0$
- e.g. 我们可以把线性数值方法写成  $Au = b$ ,  $A$  matrix,  $u = [U_1, \dots, U_{N-1}]^\top$ .

Then:  $\tau = A u_{\text{exact}} - b$  where  $u_{\text{exact}} = [u_1, \dots, u_{N-1}]^\top$ ,  $\tau = [\tau_1, \dots, \tau_{N-1}]^\top$

ps:  $Ae = -\tau$  where  $e = u - u_{\text{exact}}$  Define  $\tau(\Delta t) = [\tau_1, \dots, \tau_{N-1}]^\top$

**Consistency:** If  $\lim_{\Delta t \rightarrow 0} \|\tau(\Delta t)\| = 0$ , the method is consistent.

**Linear Equation:** Goal: Finally change ODE/PDE into  $Au = b$  where  $A$  is matrix,  $u = [U_1, \dots, U_{N-1}]^\top$

**Stability:** A numerical method is stable if  $\exists \Delta x_0$  s.t. For the linear system  $Au = b$ :

- $A(\Delta x)$  is invertible  $\forall \Delta x < \Delta x_0$
- $\|A(\Delta x)^{-1} \mathbf{w}\| \leq C \|\mathbf{w}\|$  for all  $\mathbf{w} \in \mathbb{R}^{N-1}$

**Property:** Consistent + Stable  $\Rightarrow$  Convergent (same norm!) and  $\|\mathbf{e}(\Delta t)\| = \mathcal{O}(\|\tau(\Delta t)\|)$

## 3 Intro & Method of Lines

**Taylor Series:**  $f(t + \Delta t) = f(t) + f'(t)\Delta t + \frac{1}{2}f''(t)(\Delta t)^2 + \dots + \mathcal{O}((\Delta t)^N)$  ps:  $u_{n+1} = u(n\Delta t + \Delta t)$

$f(t) = f(a) + f'(a)(t-a) + \frac{f''(a)}{2!}(t-a)^2 + \dots + \frac{f^{(N-1)}(a)}{(N-1)!}(t-a)^{N-1} + \mathcal{O}((t-a)^N)$

**Euler Methods:** Forward:  $\frac{du}{dt}|_n \Delta t \approx \frac{U_{n+1} - U_n}{\Delta t}$ ; Backward:  $\frac{du}{dt}|_n \Delta t \approx \frac{U_n - U_{n-1}}{\Delta t}$

**Leapfrog:**  $\frac{du}{dt}|_n \Delta t \approx \frac{U_{n+1} - U_{n-1}}{2\Delta t}$  **Trapezoidal:**  $\frac{1}{2} \left[ \frac{du}{dt}|_n \Delta t + \frac{du}{dt}|_{n+1} \Delta t \right] \approx \frac{U_{n+1} - U_n}{\Delta t}$

**Centred Difference:**  $\frac{d^2 u}{dt^2}|_n \Delta t \approx \frac{U_{n+1} - 2U_n + U_{n-1}}{(\Delta t)^2}$

**Method of Lines:** Using  $U_m(t) \approx u(a + m\Delta x, t)$ ,  $U_m^n \approx u(a + m\Delta x, n\Delta t)$   $a$  为空间起点, 不重要

- For 1D:** a. 先对空间  $x$  离散化, 使用  $U_m$  表达, 此时  $\frac{d}{dt} \mathbf{U} = \dots$  where  $\mathbf{U} = [U_1(t), U_2(t), \dots, U_{M-1}(t)]^\top$   
 b. 对时间  $t$  使用 ODE 离散化, 使用  $U_m^n$  表达, 得到 Linear Equation.
- Example: Crank-Nicolson for Heat Eq.  $u_t = \kappa u_{xx}$ : 对  $x$  用 centred difference; 对  $t$  用 trapezoidal:  
 $\frac{U_{m+1}^{n+1} - U_m^{n+1}}{\Delta t} = \frac{\kappa}{2} \left( \frac{U_{m+1}^{n+1} - 2U_m^{n+1} + U_{m-1}^{n+1}}{(\Delta x)^2} + \frac{U_{m+1}^n - 2U_m^n + U_{m-1}^n}{(\Delta x)^2} \right)$   
 $\Rightarrow -r U_{m-1}^{n+1} + (2 + 2r) U_m^{n+1} - r U_{m+1}^{n+1} = r U_{m-1}^n + (2 - 2r) U_m^n + r U_{m+1}^n$ ,  $r = \frac{\kappa \Delta t}{(\Delta x)^2}$

**Solve it:** By iteration: e.g.  $A \mathbf{U}^{n+1} = B \mathbf{U}^n$ , where  $\mathbf{U}^n = [U_1^n, U_2^n, \dots, U_{M-1}^n]^\top$ .

**Truncation for MoL:** 对于  $\tau_m^n$  离散  $x$  后  $U_m$  替换为  $u_m$ ; 对于  $\tau_m^n$ , 再离散  $t$  后  $U_m^n$  替换为  $u_m^n$ .

e.g. For Crank-Nicolson above:  $\tau_m = \frac{\partial u_m}{\partial t} - \kappa \frac{u_{m+1} - 2u_m + u_{m-1}}{(\Delta x)^2}$

e.g. For Crank-Nicolson above:  $\tau_m^n = \frac{u_{m+1}^n - u_m^n}{\Delta t} - \frac{\kappa}{2}(\dots)$

## 4 Lax-Richtmyer Stability, von Neumann Analysis

**Abs. Stability for ODEs:** For ODEs  $\frac{du}{dt} = A u$ ,  $u$  vector. A 1-step method for  $t$  (i.e.  $U^{n+1} = B U^n$ )

It's absolutely stable if  $\|B \mathbf{w}\|_2 \leq \|\mathbf{w}\|_2$  for all  $\mathbf{w} \in \mathbb{R}^{N-1}$

**Property:** If  $B$  normal, then absolutely stable  $\Leftrightarrow |\lambda_i(B)| \leq 1, \forall i$

**Property:** Absolute stability  $\Rightarrow$  Lax-Richtmyer stability in 2-norm and weighted 2-norm

**Lax-Richtmyer Stability:** For PDE which discrete form  $\frac{1}{\Delta t} U^{n+1} = \frac{1}{\Delta t} B U^n + F^n$ ,  $B$  fixed,  $F$  can change.

It's Lax-Richtmyer stable if  $\|B^n \mathbf{w}\| \leq C \|\mathbf{w}\|$  for all  $\mathbf{w} \in \mathbb{R}^{N-1}$ ,  $n \leq \frac{N}{\Delta t} = N$

**Property:** If consistent with  $\max_n \|\mathbf{x}^n\|_{2, \Delta x} = \mathcal{O}((\Delta x)^\beta)$  and L-R stable  $\|B^n \mathbf{w}\|_{2, \Delta x} \leq C \|\mathbf{w}\|_{2, \Delta x}$ , then, it converges with  $\max_n \|\mathbf{e}^n\|_{2, \Delta x} = \mathcal{O}((\Delta x)^\beta)$  Remark: Always Assume  $\Delta t = K(\Delta x)^\alpha$

**von Neumann Analysis:** PDE: 1. periodic BCs in spatial  $x$ , length  $D$ . 2. 满足 L-R 中 PDE 框架, 3.  $F^n = 0$

$\Rightarrow$  Let  $v_{k,m} = \exp(2\pi i k \frac{m \Delta x}{D})$  for  $m = 0, 1, \dots, M-1$ . Then vector  $\mathbf{v}_k$  为  $B$  的正交特征向量.

\* By  $U_m^n = \exp(2\pi i k \frac{m \Delta x}{D})$ ,  $U_m^{n+1} = \alpha_k U_m^n$ , 可得特征值(放大因子)  $\alpha_k$  表达式 (i.e.  $B \mathbf{v}_k = \alpha_k \mathbf{v}_k$ ).

$\Rightarrow$  If all  $\alpha_k$  for  $k = 0, 1, \dots, M-1$  can be found  $\Rightarrow B$  is normal and then:

If  $|\alpha_k| \leq 1, \forall k \Rightarrow B$  is absolutely stable and L-R stable in weighted 2-norm

**Remark:** 假设  $\Delta t = K(\Delta x)^\alpha$  会得  $K, \alpha$  的限制条件,  $\|$  For  $F^n \neq 0$ , 讨论  $F^n = 0$  足够.

## 5 Boundary Condition

**2nd ODE:** For  $u_x x = \beta(x)$  with  $x \in (a, b)$ , there are three types of BCs:

- Two Dirichlet:**  $u(x=a) = c$   $u(x=b) = d$
- One Dirichlet, One Neumann:**  $u(x=a) = c$   $u_x(x=b) = d$  or  $u_x(x=a) = c$   $u(x=b) = d$
- One Neumann, One Constraint:**  $u_x(x=a) = c$  or  $\int_a^b u(x) dx = c$

**Heat PDE:** For  $u_t = \kappa u_{xx}$  with  $x \in (a, b)$ , there are three types of BCs: \* Basic IC:  $u(x, t_0) = P(x)$

- Two Dirichlet:**  $u(x=a, t) = c$   $u(x=b, t) = d$
- One Dirichlet, One Neumann:**  $u(x=a, t) = c$   $u_x(x=b, t) = d$  or  $u_x(x=a, t) = c$   $u(x=b, t) = d$
- Two Neumann:**  $u_x(x=a, t) = c$   $u_x(x=b, t) = d$

**Remark:** For  $u_x(x, a, t) = c \Rightarrow$  describe  $\frac{u_m^n - u_0^n}{\Delta x} = c$  Boundary condition itself can effect the accuracy

For  $u_t(x, t_0) = P(x) \Rightarrow$  describe  $\frac{u_m^n - u_0^n}{\Delta t} = P(a + m\Delta x)$

## 6 One Time Solve Two Variables PDEs

**Elliptic Equation:**  $L u = F(x_1, \dots, x_d)$  is an elliptic PDE if  $L$  is an elliptic operator.

**Theorem:**  $L = \sum_{r=1}^d \sum s = 1^d A_{r,s} \frac{\partial^2}{\partial x_r \partial x_s} + \sum_{n=1}^d B_n \frac{\partial}{\partial x_n} + C$  is elliptic operator if:

A is symmetric and A is positive/negative definite ( $\lambda_i > 0 | \lambda_i < 0$ ).

**Poisson Equation:**  $\Delta u = F(x_1, \dots, x_d)$  is a Poisson equation and it's elliptic. with BCs:

Dirichlet:  $u = u_D$  on  $\partial \Omega$ ; Neumann:  $\Delta u \cdot \hat{\mathbf{n}} = g$  on  $\partial \Omega$  where  $\int_{\Omega} F = \int_{\partial \Omega} g$  and  $\int_{\Omega} u = \int_{\partial \Omega} u_D$ .

e.g. For 2D:  $u_{xx} + u_{yy} = F(x, y)$  with BCs:  $u(x, y) = u_D$  on  $\partial \Omega$

$\rightarrow$  5-point stencil:  $\frac{u_{m-1,p} + u_{m+1,p} + u_{m,p-1} + u_{m,p+1} - 4u_{m,p}}{(\Delta x)^2} = F(a + m\Delta x, a + p\Delta x)$

$\rightarrow$  Using  $\mathbf{u} = [U_{1,1}, U_{2,1}, \dots, U_{M-1,1}, U_{1,2}, \dots, U_{M-1,M-1}]^\top$  and

$\mathbf{b}_{m+(p-1)}(M-1) = F(a + m\Delta x, a + p\Delta x)$  then  $A \mathbf{u} = \mathbf{b}$  where  $A$  is a symmetric matrix.

$\rightarrow$  Eigenvalue/vectors of  $A$ :  $\mathbf{v}_{k,l} = [v_{k,l,1,1}, v_{k,l,1,2}, \dots, v_{k,l,1,2}, \dots, v_{k,l,M-1,M-1}]^\top$   
 $v_{k,l,m,p} = \sin(\pi k m \Delta x) \sin(\pi l p \Delta x)$ ,  $\lambda_{k,l} = \frac{2}{(\Delta x)^2} [\cos(\pi k \Delta x) + \cos(\pi l \Delta x) - 2] < 0$ ,  $\lambda_{\min} = \lambda_{1,1}$

**Remark:** 这一节都是用同一个  $\Delta x$  离散  $x$  和  $y$  的, 和前面  $t, x$  分别不同.

**Consistency:** If  $\lim_{\Delta x \rightarrow 0} \frac{\|\tau(\Delta x)\|}{\Delta x} = 0$ , the method is consistent.

**Convergence:** If  $\|\mathbf{e}(\Delta x)\| = \mathcal{O}((\Delta x)^p)$ , the method is  $p$ -th order accurate.

**Stability:** A numerical method is stable if for the linear system  $A \mathbf{u} = \mathbf{b}$ :

- $A(\Delta x)$  is invertible  $\forall \Delta x < \Delta x_0$
- $\|A(\Delta x)^{-1} \mathbf{w}\| \leq C \|\mathbf{w}\|$  for all  $\mathbf{w} \in \mathbb{R}^{N-1}$

**Property:** If  $A$  normal, then  $\|A^{-1} \mathbf{w}\|_2 \leq \frac{1}{|\lambda_1|} \|\mathbf{w}\|_2$  where  $0 < |\lambda_1| \leq \dots \leq |\lambda_{N-1}|$  are eigenvalues of  $A$ .

## 7 Spectral Methods

**Inner Product Space for Function:**  $\langle \phi_i, \phi_j \rangle = \int_a^b \phi_i \phi_j w dx$  ps:  $w(x)$  is weight function.

For Fourier Series:  $w(x) = 1, a = -\pi, b = \pi$ :  $\sin(n x), \cos(m x)$  are orthogonal basis.

**Fourier Series:**  $f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos(n x) + b_n \sin(n x) = \sum_{n=-\infty}^{\infty} A_k e^{i k x}$  where

$\rightarrow a_0 = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) dx$   $a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(n x) dx$   $b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(n x) dx$ .

$\rightarrow k > 0: A_k = \frac{1}{2}(a_k - i b_k)$ ;  $k < 0: A_k = \frac{1}{2}(a_{-k} + i b_{-k})$ ;  $k = 0: A_0 = \frac{a_0}{2}$ . For real func:  $A_{-k} = \overline{A_k}$

**Property:**  $A_k = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-i k x} dx$  and  $f'(x) = \sum_{k=-\infty}^{\infty} (i k) A_k e^{i k x}$

**More on Fourier Series:** Assume  $f$  is periodic with period  $2\pi$ , then:

- The Fourier Transform:  $\mathcal{F}\{f(x)\} = \hat{f}(k) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) e^{-i k x} dx$
- The inverse Fourier Transform:  $\mathcal{F}^{-1}\{\hat{f}(k)\} = f(x) = \sum_{k=-\infty}^{\infty} \hat{f}(k) e^{i k x}$

**Change  $L \rightarrow 2\pi$  period:** If  $f(x)$  has period  $L \Rightarrow g(x) = f(\frac{L}{2\pi} x)$  has  $2\pi$  period.

**Discrete Fourier Transform (DFT):** If  $x_n = n\Delta x$  equally-spaced with  $n = 0, 1, \dots, N-1$ ;  $\Delta x = \frac{2\pi}{N}$ :

1. **Fourier Transform:**  $\mathcal{F}\{f\} = \hat{f}_k = \frac{2\pi}{N} \sum_{n=0}^{N-1} f(x_n) e^{-i k x_n}$   $k = -\frac{N}{2} + 1, \dots, \frac{N}{2}$

2. **Inverse FT:**  $\mathcal{F}^{-1}\{\hat{f}\} = f(x) = \frac{1}{2\pi} \sum_{k=-N/2+1}^{N/2} \hat{f}_k e^{i k x_n} = \frac{1}{2\pi} \sum_{k=-N/2}^{N/2} \hat{f}_k e^{i k x_n}$

where  $\hat{f}_k = \hat{f}_k$  for  $k = -\frac{N}{2} + 1, \dots, \frac{N}{2} - 1$ ;  $\hat{f}_{-N/2} = \hat{f}_{N/2} = \frac{1}{2} \hat{f}_{N/2}$ .

**Property:**  $\hat{f}'_k = (i k) \hat{f}_k \Leftrightarrow \mathcal{F}\{f'\} = (i k) \mathcal{F}\{f\}$

Remark: Quickly compute DFT: Fast Fourier Transform (FFT)

**Using Fourier to solve PDE:** For PDE with periodic BCs

- Take Fourier Transform in PDE (in space) e.g.  $u_t = u_{xx}$   $\rightarrow \frac{d}{dt} \hat{u}_k = -k^2 \hat{u}_k$
- Take Fourier Transform in IC (in space) e.g.  $u(x, 0) = f(x) \rightarrow \hat{u}_k(t=0) = \hat{f}_k$
- Using Transformed PDE and IC  $\rightarrow$  a. Directly Solve ODE. b. numerical method for ODE.  $\rightarrow \hat{u}_k(t)$
- Inverse Fourier Transform to get  $u(x_n, t)$  对  $x_n$  点  $t$  的估计或精确解.

**Ideal of Interpolants:** 给定  $(x_n, f(x_n))$ , 构造插值函数  $p(x)$  s.t.  $p(x_n) = f(x_n) \Rightarrow$  用  $p'(x_n)$  估计  $f'(x_n)$ .

**Spectral Collocation:** Given  $(x_n, f(x_n))$ ,  $f$  has period  $2\pi$ , construct Fourier Interpolant  $p(x)$  s.t.  $p(x_n) = f(x_n)$

- DFT:  $\hat{f}_k = \frac{2\pi}{N} \sum_{n=0}^{N-1} f(x_n) e^{-i k x_n} \Rightarrow p(x) = \frac{1}{2\pi} \sum_{k=-N/2}^{N/2} \hat{f}_k e^{i k x}$
- (Method I) (即 FFT, 无法得到微分算子的矩阵) Direct differentiate  $p(x)$ , and  $f'(x_n) \approx p'(x_n)$ .
- (Method II) Assume  $p(x) = \sum_{m=0}^{N-1} f(x_m) P_\delta(x - x_m)$  where  $P_\delta(x_j - x_m) = \delta_{j-m}$   
 $\Rightarrow P_\delta(x - x_m) = \frac{1}{2\pi} \sum_{k=-N/2}^{N/2} \hat{P}_{\delta k-m} e^{i k x} = \frac{1}{2\pi} \sum_{k=-N/2}^{N/2} e^{i k(x-x_m)} \text{ and } P_\delta(x) = \frac{\sin(\pi x / \Delta x)}{(2\pi / \Delta x) \tan(x/2)}$   
 $\Rightarrow f'(x_n) \approx p'(x_n) = \sum_{m=0}^{N-1} f(x_m) P'_\delta(x_n - x_m)$  and  $P'_\delta(x_n - x_m) = \begin{cases} 0 & j \equiv 0 \pmod{N} \\ \frac{1}{2}(-1)^j \cot(\frac{i h}{2}) & j \not\equiv 0 \pmod{N} \end{cases}$

$\Rightarrow$  Can be written in matrix form:  $\mathbf{w} = D \mathbf{v}$  其中  $\mathbf{w}$  是  $f'(x_n)$  的估计向量,  $\mathbf{v}$  是  $f(x_n)$  的向量 and  $f^{(n)} \Leftrightarrow D^n$

**Chebyshev Spectral Method:** For non-periodic BCs. Assume points are  $x_n = \cos(\frac{n\pi}{N})$  for  $n = 0, 1, \dots, N$ .

Then  $\mathbf{w} = D \mathbf{v}$  其中  $\mathbf{w}$  是  $f'(x_n)$  的估计向量,  $\mathbf{v}$  是  $f(x_n)$  的向量 and  $f^{(n)} \Leftrightarrow D^n$  where

$D_N \in \mathbb{R}^{(N+1) \times (N+1)}$  is:

10 The Steepest Descent Method

**Theorem:** Let  $\phi(\mathbf{x}) = \frac{1}{2} \mathbf{x}^T A \mathbf{x} - \mathbf{b}^T \mathbf{x}$ , then it has unique minimum if  $A$  is *positive definite*.  
If  $A$  is symmetric positive definite  $\Rightarrow \mathbf{x}: A\mathbf{x} = \mathbf{b} \Leftrightarrow \mathbf{x}: \phi(\mathbf{x}) = \min_{\tilde{\mathbf{x}}} \phi(\tilde{\mathbf{x}}) = \min_{\tilde{\mathbf{x}}} [\frac{1}{2} \tilde{\mathbf{x}}^T A \tilde{\mathbf{x}} - \mathbf{b}^T \tilde{\mathbf{x}}]$   
**Following need  $A$  to be symmetric positive definite.**

**Steepest Descent Method:** To solve  $\min \phi(x)$  or  $A\mathbf{x} = \mathbf{b}$ , we can use the following iteration:  
 $\mathbf{x}^{(k)} = \mathbf{x}^{(k-1)} + \alpha_{k-1} \nabla \phi(\mathbf{x}^{(k-1)})$  and  $\alpha_{k-1}$  s.t.  $\min_{\alpha} \phi(\mathbf{x}^{(k-1)} - \alpha \nabla \phi(\mathbf{x}^{(k-1)}))$ .  
1. **Property 1:**  $\nabla \phi(\mathbf{x})$  is the direction of steepest ascent (most rapid increase) of  $\phi$  at  $\mathbf{x}$ .  
2. **Property 2:**  $\alpha_{k-1} = \frac{(\mathbf{r}^{(k-1)})^T \mathbf{r}^{(k-1)}}{(\mathbf{r}^{(k-1)})^T A \mathbf{r}^{(k-1)}}$  and  $\nabla \phi(\mathbf{x}^{(k)}) = A\mathbf{x}^{(k)} - \mathbf{b} =: -\mathbf{r}^{(k)}$  is  $k$ -th residual.  
3. **Property 3:**  $\mathbf{r}^{(k)} = \mathbf{r}^{(k-1)} - \alpha_{k-1} A \mathbf{r}^{(k-1)}$

Remark: Convergence behaviour is *highly problem-dependent*. (如果  $\phi$  的等高线是圆形则快, 椭圆形则慢)  
**The Conjugate Gradient Method:** To solve  $\min \phi(x)$  or  $A\mathbf{x} = \mathbf{b}$ , we can use the following iteration:  
 $\mathbf{x}^{(k)} = \mathbf{x}^{(k-1)} + \alpha_{k-1} \mathbf{p}^{(k-1)}$  and  $\alpha_{k-1}$  s.t.  $\min_{\alpha} \phi(\mathbf{x}^{(k-1)} - \alpha \mathbf{p}^{(k-1)})$   
and  $\mathbf{p}^{(k)}$  is  $A$ -conjugate to previous search directions:  $\mathbf{p}^{(k-1)}: (\mathbf{p}^{(k)})^T A \mathbf{p}^{(k-1)} = 0$   
1. **Property 1:**  $\alpha_{k-1} = \frac{(\mathbf{r}^{(k-1)})^T \mathbf{r}^{(k-1)}}{(\mathbf{p}^{(k-1)})^T A \mathbf{p}^{(k-1)}}$  and  $-\mathbf{r}^{(k)} = A\mathbf{x}^{(k)} - \mathbf{b}$  is  $k$ -th residual.  
2. **Property 2:**  $\mathbf{p}^{(k)} = \mathbf{r}^{(k)} + \beta_{k-1} \mathbf{p}^{(k-1)}$  where  $\beta_{k-1} = \frac{(\mathbf{r}^{(k)})^T \mathbf{r}^{(k)}}{(\mathbf{r}^{(k-1)})^T \mathbf{r}^{(k-1)}}$   
3. **Property 3:**  $(\mathbf{p}^{(k)})^T A \mathbf{p}^{(j)} = 0$  and  $(\mathbf{r}^{(k)})^T \mathbf{r}^{(j)} = 0$  for  $j = 0, 1, \dots, k-1$   
4. **Property 4:**  $\text{span}\{\mathbf{p}^{(0)}, \dots, \mathbf{p}^{(k-1)}\} = \text{span}\{A^0 \mathbf{r}^{(0)}, \dots, A^{k-1} \mathbf{r}^{(0)}\} = \text{span}\{A \mathbf{e}^{(0)}, \dots, A^k \mathbf{e}^{(0)}\}$   
where  $\mathbf{e}^{(k)} = \mathbf{x}^{(k)} - \mathbf{x}$  is the error at  $k$ -th iteration. (Quicker than JAC/GAU)  
5. **Exact Property:** For  $A \in \mathbb{R}$  has  $n$  distinct eigenvalues  $\Rightarrow$  solve in at most  $n$  iterations. (By exact arithmetic)

**A-Norm:** The  $A$ -norm of  $\mathbf{x}$  is defined as  $\|\mathbf{x}\|_A = \sqrt{\mathbf{x}^T A \mathbf{x}}$ .  
**Condition Number:** The condition number of  $A$  is  $\kappa(A) = \|A\| \|A^{-1}\| \geq 1$  and  $\kappa_2(A) = \frac{\lambda_{\max}(A)}{\lambda_{\min}(A)}$

**Error of Conjugate Gradient:** To solve  $A\mathbf{x} = \mathbf{b}$ , error is:  $\frac{\|\mathbf{e}^{(k)}\|_A}{\|\mathbf{e}^{(0)}\|_A} \leq 2 \left( \frac{\sqrt{\kappa(A)}-1}{\sqrt{\kappa(A)}+1} \right)^k$

11 Appendix

**11.1 Other**  
**Divergence Theorem:**  $\int_{\Omega} \nabla \cdot \mathbf{F} dV = \int_{\partial \Omega} \mathbf{F} \cdot \hat{\mathbf{n}} dS$   $\hat{\mathbf{n}}$  outward normal.  $\nabla^2 = \nabla \cdot \nabla = \Delta$   
**How to improve Approximation Accuracy?:** a. Using more neighboring points. (more terms from Taylor expansion) b. Make  $\Delta t, \Delta x$  smaller. c. For Periodic BCs, use Fourier Transform.