

1 Notation & Perturbation Method

Big $\mathcal{O}$: $f(z) = \mathcal{O}(g(z))$ as $z \rightarrow z_0$ if $\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)}$ bdd $\Leftrightarrow |f(z)| \leq A|g(z)|$ as $z \rightarrow z_0$
Small o : $f(z) = o(g(z))$ as $z \rightarrow z_0$ if $\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = 0 \Leftrightarrow f(z) \ll g(z)$ as $z \rightarrow z_0$
Asymptotic $\sim$: $f(z) \sim g(z)$ as $z \rightarrow z_0$ if $\lim_{z \rightarrow z_0} \frac{f(z)}{g(z)} = 1$ **ps:** $e^{-x} = o(x^{-n}), \forall n > 0$ as $x \rightarrow \infty$
Asymptotic Sequence: Seq $\{\varphi_n(z)\}$ is asy. if $\forall n \in \mathbb{N} \varphi_{n+1}(z) = o(\varphi_n(z))$ as $z \rightarrow z_0$
Asymptotic Expansion: $f(z) \sim \sum_{n=0}^M a_n \varphi_n(z)$ as $z \rightarrow z_0$ to M if $\phi_n(z)$ is asy. seq. and
 $f(z) - \sum_{n=0}^N a_n \varphi_n(z) = o(\varphi_N(z)) = \mathcal{O}(\varphi_{N+1}(z)), \forall N \leq M$ as $z \rightarrow z_0$
 $f(x, \varepsilon) \sim \sum_{n=0}^{\infty} a_n(x) \varepsilon^n$ uniformly in $x \in I$ as $\varepsilon \rightarrow 0$ if ε 与 x 无关 (i.e. $\sup_{x \in I} |f(x, \varepsilon) - \sum_{n=0}^N a_n(x) \varepsilon^n| = o(\varepsilon^N)$)
Singular|Regular - Perturbation Method: a. Let $\varepsilon = 0$ solve $f(x, 0) = 0$.
b.i. 如果 f 结构变了 (阶数降低), Let $x = \varepsilon^{-\alpha} X$, PDB 找 α c. 假设 $X = a_0 + a_1 \varepsilon^\beta + a_2 \varepsilon^{2\beta} + \dots$ 代入求解
b.ii. 如果 f 结构不变/变, 但有重根 x_0 , Let $x = x_0 + \varepsilon^\alpha X$, PDB 找 α . $\uparrow \beta$ 看化简后的 X 方程得出
Principle of dominant balance (PDB): For Singular Perturbation $p(x, \varepsilon) = 0$, Let $x = \varepsilon^\alpha X$, 代入选择 α :
对于 ε 来说 a. 至少有两项阶数相同且最低; b. 其他项阶数均更高. (即可忽略)

2 Integral By Parts, Laplace's Method & Method of Steepest Descents

Taylor: $f(x) = \sum_{n=0}^{N-1} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n + \frac{f^{(N)}(\xi)}{N!} (x - x_0)^N \Rightarrow f(x) \sim \sum_{n=0}^{\infty} \frac{f^{(n)}(x_0)}{n!} (x - x_0)^n$ as $x \rightarrow x_0$
Exp. Integral: $E_1(x) = \int_x^\infty \frac{e^{-t}}{t} dt, (x \rightarrow 0 \text{ 发散}) \quad E_1(x) \sim e^{-x} \sum_{n=0}^{\infty} \frac{(-1)^n}{x^{n+1}}, x \rightarrow \infty$
Gamma Func: $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt \quad \Gamma(x+1) = x\Gamma(x)$ and $\Gamma(n) = (n-1)!$ $\Gamma(\frac{1}{2}) = \sqrt{\pi}$
Incomplete Gamma Func: **Lower:** $\gamma(a, x) = \int_0^x t^{a-1} e^{-t} dt$ **Upper:** $\Gamma(a, x) = \int_x^\infty t^{a-1} e^{-t} dt$
- $\gamma(a, x) = \Gamma(a) - \Gamma(a, x) \quad \Gamma(a, x) \sim e^{-x} x^{a-1} \sum_{n=0}^{\infty} \frac{\Gamma(a)}{\Gamma(a-n)} x^{-n}, x \rightarrow \infty$
Laplace's Method: For $I(x) = \int_a^b e^{x f(t)} g(t) dt$ as $x \rightarrow \infty$
1. If $f(t)$ has unique max at $t = t_0 \in (a, b) \Rightarrow I \sim \int_{-\infty}^\infty e^{x[f(t_0) + \frac{f''(t_0)}{2}(t-t_0)^2 + \dots]} g(t_0) dt$
(留 n 项非常数展开为 n term approximation) (ps: $\max \leftrightarrow f'(t_0) = 0, f''(t_0) < 0$)
2. 对于留 n 项展开: e.g. $n = 2$. 由 1 基础 $\rightarrow x f(t_0) \sim 1 \rightarrow t = \mathcal{O}(x^{-\frac{1}{2}}) \rightarrow u = t_0 + x^{\frac{1}{2}} t$ 换元求解
3. **Exact Substitution:** 由 1 的基础, 得到 $f(t)$ 在 t_0 处的展开式, 换元令 $f(t)$ 就等于展开式前几项求解.
4. If $g = g(t, x): e^{x f(t)} g(t, x) = e^{x f(t) + \ln(g(t, x))}$, 看 $x f(t) + \ln(g(t, x))$ 的最大值, 类似 1 处理.
Method of Steepest Descents: For $I(x) = \int_C e^{x f(z)} g(z) dz$ as $x \rightarrow \infty$. f 容易避开 singular point
If z_0 is saddle point: $f'(z_0) = 0, f''(z_0) \neq 0 \quad f(z)$ Taylor at $z_0: f(z) = f(z_0) + \frac{f''(z_0)}{2}(z - z_0)^2 + \dots$
1. Let $z = z_0 + e^{i\theta} y$ (θ 选让 $f''(z_0) e^{2i\theta}$ 为负实数) $\rightarrow I(x) \sim \int_{-\infty}^\infty e^{x[f(z_0) - \frac{|f''(z_0)|}{2} y^2 + \dots]} g(z_0) e^{i\theta} dy$
2. If f 不容易避开 singular point: a. $\text{Im}(f(z)) = \text{Im}(f(z_0))$ and $z_0 \in C_0$ to find steepest descent path C_0
3. b. 画图看哪些 singular points 在 $C \rightarrow C_0$ 变形路径上, 计算 residues. c. $I = 1$ 部分 $+ 2\pi i \sum$ residues.
Residue: If z_0 pole of order m of $f, \Rightarrow \text{Res}(f, z_0) = \frac{1}{(m-1)!} \lim_{z \rightarrow z_0} \frac{d^{m-1}}{dz^{m-1}} [(z - z_0)^m f(z)]$

3 Appendix

x, x^2, x^3 as $x \rightarrow 0, \quad x^{-1}, x^{-2}, x^{-3}$ as $x \rightarrow \inf$
 $y' + Py = Q: \mu y = \int Q(x) e^{\mu x} dx, \quad \mu = e^{\int P(x) dx} \quad \downarrow x \geq 0$
Gaussian Integral: $\int_{-\infty}^\infty e^{-ax^2} dx = \frac{\sqrt{\pi}}{\sqrt{a}}$ **Inequalities:** $\ln(x+1) < x, \quad e^x > x+1, \quad (1+x)^a < e^{ax}$
Binomial Expansion: $(1+x)^\alpha = \sum_{n=0}^\infty \binom{\alpha}{n} x^n = 1 + \alpha x + \frac{\alpha(\alpha-1)}{2!} x^2 + \dots, |x| < 1$
Taylor: $f(x+h) = \sum_{n=0}^\infty \frac{f^{(n)}(x)}{n!} h^n = f(x) + f'(x)h + \frac{f''(x)}{2!} h^2 + \dots$
 $\sin(x) = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots \quad \cos(x) = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots \quad e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$
 $\ln(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \dots \quad \sinh(x) = x + \frac{x^3}{3!} + \frac{x^5}{5!} + \dots \quad \cosh(x) = 1 + \frac{x^2}{2!} + \frac{x^4}{4!} + \dots$
 $\sum_{n=0}^\infty x^n = \frac{1}{1-x} \quad \tan(x) = x + \frac{x^3}{3} + \frac{2x^5}{15} + \dots \quad \cot(x) = \frac{1}{x} - \frac{x}{3} - \frac{x^3}{45} - \dots$
Integral: $\int \tan x = -\ln |\cos x| = \ln |\sec x| \quad \int \cot x = \ln |\sin x| = -\ln |\csc x| \quad \int_0^\infty e^{-x} t^n dx = \frac{1}{n} \Gamma(\frac{m+1}{n}) x^{-\frac{m+1}{n}}$