

Soft-constrained Schrödinger Bridge for Diffusion Model Sampling

Presenter: Quan Zhou

Department of Statistics, Texas A&M University

Joint work with Jhanvi Garg and Xianyang Zhang

The research presented in this talk is supported by NSF
DMS-2245591, DMS-2311307.



Introduction: Langevin diffusion

Given a distribution π on \mathbb{R}^d , we can generate samples from it by simulating the Langevin diffusion (LD):

$$dX_t = \sigma^2 \nabla \log \pi(X_t) dt + \sqrt{2\sigma^2} dW_t$$

where W_t denotes the d -dimensional standard Brownian motion.

Comments on Langevin diffusion

- Under some regularity conditions, X_t converges in distribution to π as $t \rightarrow \infty$ regardless of the initial distribution.
 - If π satisfies a Poincaré inequality, then the convergence is exponentially fast.
 - Convergence can be quite slow if the target is multimodal.
 - To simulate the Langevin diffusion, we need access to $\nabla \log \pi(x)$.
 - When simulating the unadjusted Langevin diffusion, the step size needs to be sufficiently small.


A Gaussian Mixture Example

Consider the following Gaussian mixture on \mathbb{R}^2 :

$$\begin{aligned} & 0.1 \mathcal{N}((1, 1), 0.05^2 I) \\ & + 0.2 \mathcal{N}((-1, 1), 0.05^2 I) \\ & + 0.3 \mathcal{N}((1, -1), 0.05^2 I) \\ & + 0.4 \mathcal{N}((-1, -1), 0.05^2 I). \end{aligned}$$

Simulate the (unadjusted) LD with $\sigma = 0.02$ and time increment size $1/200 = 0.005$.

A Gaussian Mixture Example



1000 simulated trajectories with $X_0 = 0$.

Introduction: Schrödinger Bridge

Consider the SDE

$$\begin{aligned} dX_t &= u^*(X_t, t)dt + \sigma dW_t, \\ X_0 &= 0, \end{aligned}$$


with

$$u^*(x, t) = \sigma^2 \nabla_x \log \int \frac{\pi(y)}{\phi_\sigma(y)} \phi_{\sigma\sqrt{1-t}}(y - x) dy.$$

where ϕ_σ denotes the density of $\mathcal{N}(0, \sigma^2 I)$.

We have $X_1 \sim \pi$ (this is a special case of the Schrödinger bridge).


A Gaussian Mixture Example




1000 simulated trajectories with $\sigma = 1$ and time increment size
 $1/200 = 0.005$.

Comments on the Schrödinger Bridge

- The SB process behaves like a Brownian motion for a long time, and the “denoising” effect becomes increasingly significant as the simulation gets close to completion.
- Under some regularity conditions, it is guaranteed that $X_1 \sim \pi$, but the drift term depends on the initial distribution.
- To simulate the SB process, we need access to the score of some “perturbed” version of π instead of π itself.



$t = 0$



$t = 0.31$



$t = 0.60$



$t = 0.63$


From De Bortoli et al. [6]

Schrödinger Bridge Problem

Let $X = (X_t)_{0 \leq t \leq 1}$ be a diffusion process with $X_0 \sim f_0$, where f_0 denotes the density w.r.t. a dominating measure ν . Denote the density of X_1 by p_1 .

How to “optimally” modify the dynamics of X so that its distribution at time 1 has density $f_1 \neq p_1$ (w.r.t. the Lebesgue measure)?

Hot Gas Experiment



Schrödinger Bridge Problem

Let $X = (X_t)_{0 \leq t \leq 1}$ denote a weak solution to the SDE

$$X_0 = \xi,$$

$$\mathrm{d}X_t = b(X_t, t)dt + \sigma dW_t,$$

where $b: \mathbb{R}^d \times [0, 1] \rightarrow \mathbb{R}^d$, $\sigma \in (0, \infty)$, and $\xi \sim f_0$ is independent of W .

Given a control $u = (u_t)_{0 \leq t \leq 1}$, define the *controlled process* X^u by

$$X_0^u = \xi,$$

$$\mathrm{d}X_t^u = [b(X_t^u, t) + u_t] dt + \sigma dW_t.$$

Schrödinger Bridge Problem

Let \mathcal{U} denote the set of *admissible* controls, and

$$\mathcal{U}_0 = \{u \in \mathcal{U} : \text{Law}(X_1^u) \text{ has density } f_1\}.$$

Schrödinger Bridge (SB) Problem

Find $V = \inf_{u \in \mathcal{U}_0} J(u)$, where

$$J(u) = \mathbb{E} \int_0^1 \frac{\|u_t\|^2}{2\sigma^2} dt,$$

and find the optimal control u^* such that $J(u^*) = V$.

Schrödinger Bridge Problem

Let \mathcal{U} denote the set of admissible controls, and

$$\mathcal{U}_0 = \{u \in \mathcal{U} : \text{Law}(X_1^u) \text{ has density } f_1\}.$$

Schrödinger Bridge (SB) Problem

Find $V = \inf_{u \in \mathcal{U}_0} J(u)$, where

$$J(u) = \mathbb{E} \int_0^1 \frac{\|u_t\|^2}{2\sigma^2} dt = \mathcal{D}_{\text{KL}}(\mathbb{P}_X^u, \mathbb{P}_X),$$

and find the optimal control u^* such that $J(u^*) = V$.

$\mathcal{D}_{\text{KL}}(\nu, \mu) = \int \log\left(\frac{d\nu}{d\mu}\right) d\nu$ denotes the Kullback-Leibler divergence.

Application to Denoising Diffusion Probabilistic Models

Many existing score-based generative modeling methods are essentially numerical approximations to the solution of SB problem.

- Denoising diffusion probabilistic models of [18, 21]
- Two-stage Schrödinger bridge algorithm of [22]
- Diffusion Schrödinger bridge algorithm of [6]
- Time-series Schrödinger bridge algorithm of [16]

In these problems, f_0 is some reference distribution (e.g. normal), and f_1 is the target distribution (e.g. distribution of the images in CelebA). Often f_1 is unknown but we have *samples from f_1* .

Solution to the SB Problem

Let $p_{1|0}(x | y)$ denote the conditional density of $X_1 = x$ given $X_0 = y$.

Theorem 3.2 of Dai Pra [5]

Suppose there exist integrable functions $\rho_0, \rho_1 \geq 0$ such that

$$\begin{aligned}f_0(y) &= \rho_0(y) \int p_{1|0}(x | y) \rho_1(x) dx, \\f_1(x) &= \rho_1(x) \int p_{1|0}(x | y) \rho_0(y) \nu(dy).\end{aligned}$$

Under some regularity conditions, the optimal control for the SB problem is $u_t^* = u^*(X_t^{u^*}, t)$, where

$$u^*(x, t) = \sigma^2 \nabla_x \log E[\rho_1(X_1) | X_t = x].$$

Solution to the SB Problem

That is, under the optimal control u^* , the joint distribution of $(X_0^{u^*}, X_1^{u^*})$ has density

$$\rho(y, x) = \rho_0(y)p_{1|0}(x | y)\rho_1(x).$$

By matching the marginal distributions, one gets the Schrödinger system in the previous slide.

This is also a well studied problem in the statistical literature [8, 20].

Two Examples for SB

Assume $b = 0$ for both examples, and let ϕ_σ be the density of $N(0, \sigma^2 I)$.
The optimally controlled process satisfies

$$dX_t^{u^*} = u^*(X_t^{u^*}, t)dt + \sigma dW_t.$$

We now give two examples where $u(x, t)$ can be explicitly characterized.

Example 1 for SB

Example 1

If the initial distribution is a Dirac measure at x_0 , then


$$\rho_1(x) = \frac{f_1(x)}{\phi_\sigma(x - x_0)}.$$

If f_1 is known up to a normalizing constant (e.g. a posterior distribution in Bayesian statistics), one can use Monte Carlo sampling to approximate

$$u^*(x, t) = \sigma^2 \nabla_x \log \int \rho_1(y) \phi_{\sigma\sqrt{1-t}}(y - x) dy.$$

See [17] for more sophisticated schemes.

Example 1 for SB



The Gaussian mixture example.

Example 2 for SB

Example 2

If $f_0(x) = \int f_1(y)\phi_\sigma(x - y)dy$, then $\rho_1(x) = f_1(x)$,


$$u^*(x, t) = \sigma^2 \nabla_x \log \int f_1(y)\phi_{\sigma\sqrt{1-t}}(x - y)dy.$$

This integral is the density of $Y + \sigma\sqrt{1-t}Z$, where $Y \sim f_1$ and $Z \sim N(0, I)$. The function

$$s(x, \sigma) = \nabla_x \log \int f_1(y)\phi_{\sigma\sqrt{1-t}}(x - y)dy$$

is called the *score*. How to estimate $s(x, \sigma)$ if one has samples from f_1 .

Score Matching



By adding Gaussian noise to these samples, one can train a neural network for approximating the score function $s(x, \sigma)$ [19].
What happens when $\sigma = 0$?


Example 2 for SB

Example 2 (continued)

To numerically simulate the solution to the SB problem, one still needs samples from $f_0(x) = \int f_1(y)\phi_\sigma(x - y)dy$. Some possible solutions:

- If σ is sufficiently large, one can assume f_0 is approximately gaussian. This yields the denoising diffusion model sampling algorithm of [21].
- One can train another SB process such that the terminal distribution coincides with f_0 . This is the approach taken in Wang et al. [22].
- Assuming the score $s(x, \sigma)$ is available, one can run a Langevin diffusion targeting f_0 for a sufficiently long time.

Schrödinger's Cat Denoising Herself



Soft-constrained Schrödinger Bridge

Recall \mathcal{U} denotes the set of admissible controls. Let μ_1 be the prob measure with density f_1 .

Soft-constrained Schrödinger Bridge (SSB) Problem

For $\beta > 0$, find $V = \inf_{u \in \mathcal{U}} J_\beta(u)$, where

$$J_\beta(u) = \beta \mathcal{D}_{\text{KL}}(\mathcal{L}aw(X_1^u), \mu_1) + \mathbb{E} \int_0^1 \frac{\|u_t\|^2}{2\sigma^2} dt,$$

and find the optimal control u^* such that $J_\beta(u^*) = V$.

Thanks to Tiziano!

Solution to the SSB Problem

Theorem 4 of Garg et al. [14]

Suppose there exist integrable functions $\rho_0, \rho_1 \geq 0$ such that

$$f_0(y) = \rho_0(y) \int p_{1|0}(x | y) \rho_1(x) dx,$$

$$f_1(x) = \rho_1(x)^{(1+\beta)/\beta} \int p_{1|0}(x | y) \rho_0(y) dy.$$

Under some regularity conditions, the optimal control for the SSB problem is $u_t^* = u^*(X_t^{u^*}, t)$, where

$$u^*(x, t) = \sigma^2 \nabla_x \log E[\rho_1(X_1) | X_t = x].$$

Comparison between SB and SSB

- For SB, $\mathcal{L}aw(X_1^{u^*})$ has density f_1 . For SSB, the density of $X_1^{u^*}$ is proportional to

$$f_1(x)^{\beta/(1+\beta)} \left(\int p_{1|0}(x | y) \rho_0(y) dy \right)^{1/(1+\beta)}.$$

So its law is a geometric mixture of f_1 and another distribution.

- For SB, the solution does not exist if $\mathcal{D}_{KL}(\mu_1, \mathcal{L}aw(X_1)) = \infty$ (e.g. when f_1 is the Cauchy distribution and X is a Wiener process). For SSB, the solution always exists.
- As $\beta \rightarrow \infty$, the solution of SSB converges to that of SB. (See Garg et al. [14] for precise statements.)

Example 1 for SSB

Example 1

If the initial distribution is a Dirac measure at x_0 ,

$$\rho_1(x) = \left(\frac{f_1(x)}{p_{1|0}(x | x_0)} \right)^{\beta/(1+\beta)}.$$

If f_1 is known up to a normalizing constant, Monte Carlo sampling can be used to simulate the resulting solution to the SSB problem.

$\mathcal{L}aw(X_1^{u^*})$ has density proportional to

$$f_1(x)^{\beta/(1+\beta)} p_{1|0}(x | x_0)^{1/(1+\beta)}.$$

Example 2 for SSB

Example 2

Assume $b = 0$. If

$$f_0(y) = c^{-1} \int \phi_\sigma(x - y) f_1(x)^{\frac{\beta}{1+\beta}} dx,$$

where $c = \int f_1(x)^{\beta/(1+\beta)} dx$ is the normalizing constant assumed to be finite. Then,

$$\rho_0(y) = c^{-(1+\beta)}, \quad \rho_1(x) = c^\beta f_1(x)^{\beta/(1+\beta)}.$$

Hence,

$$u^*(x, t) = \sigma^2 \nabla_x \log \int f_1(y)^{\beta/(1+\beta)} \phi_{\sigma\sqrt{1-t}}(x - y) dy.$$

Numerical Example for Normal Mixtures

Let the uncontrolled process X be such that $\mathcal{L}aw(X_1)$ is

$$0.1 N((1, 1), 0.05^2 I) + 0.2 N((-1, 1), 0.05^2 I) + \\ 0.3 N((1, -1), 0.05^2 I) + 0.4 N((-1, -1), 0.05^2 I).$$


Let our target terminal distribution μ_1 be

$$0.5 N((1.2, 0.8), 0.5^2 I) + 0.5 N((-1.5, -0.5), 0.5^2 I).$$


We solve the resulting SSB problem; that is, minimize

$$J_\beta(u) = \beta \mathcal{D}_{\text{KL}}(\mathcal{L}aw(X_1^u), \mu_1) + \mathsf{E} \int_0^1 \frac{\|u_t\|^2}{2\sigma^2} dt.$$

Numerical Example for Normal Mixtures



SSB trajectories for normal mixture targets.



Application to Generative Modeling

Suppose we have access to two data sets.

- \mathcal{D}_{ref} : a large set of high-quality samples with distribution μ_{ref}
- \mathcal{D}_{obj} : a small set of noisy samples with distribution μ_{obj}

Our objective is to generate realistic samples resembling those in \mathcal{D}_{obj} . We can use SSB as a regularization method to mitigate overfitting to \mathcal{D}_{obj} .

For simplicity, we set $X_0 = 0$, and we know that $\mathcal{L}aw(X_1^{u^*})$ has density proportional to

$$f_{\text{ref}}(x)^{1/(1+\beta)} f_{\text{obj}}(x)^{\beta/(1+\beta)}.$$

Score Matching


To simulate a diffusion process with terminal distribution being the geometric mixture, we need to learn the score

$$s(x, \tilde{\sigma}) = \nabla_x \log \int f_{\text{ref}}(x)^{1/(1+\beta)} f_{\text{obj}}(x)^{\beta/(1+\beta)} \phi_{\tilde{\sigma}}(x - y) dy.$$

Given only samples from μ_{ref} and μ_{obj} , we can combine the existing score matching algorithm with *importance sampling* to train a neural network for approximating $s(x, \tilde{\sigma})$; see Garg et al. [14] for details.



MNIST Example

- \mathcal{D}_{obj} : 50 noisy images labeled as “8”
- \mathcal{D}_{ref} : all 54,149 clean images not labeled as “8”



(added entrywise noise $\sim N(0, 0.4^2)$)

MNIST Example

 $\beta = 0$  $\beta = 0.25$  $\beta = 0.7$  $\beta = 1.5$  $\beta = 4$  $\beta = 100$ 

MNIST Example

Generated 40K samples and compared them to clean digit “8” images (sample size \approx 6K) using the Fréchet Inception Distance (FID) metric.

β	0	0.25	0.7	1.5	4	100
FID	67.4	66.4	61.0	56.3	110.9	182.4

On the Existence of Solution to SSB

How to find the pair (ρ_0, ρ_1) that satisfies the following system?

$$(1) \quad f_0(y) = \rho_0(y) \int p_{1|0}(x | y) \rho_1(x) dx, \quad (1)$$

$$(2) \quad f_1(x) = \rho_1(x)^{(1+\beta)/\beta} \int p_{1|0}(x | y) \rho_0(y) dy. \quad (2)$$

Initial guess $\hat{\rho}_0 \Rightarrow$ calculate $\hat{\rho}_1$ by (2) \Rightarrow update $\hat{\rho}_0$ by (1) $\Rightarrow \dots$

- If this iteration has a fixed point, then SSB has a solution.
- When $\beta = \infty$, this algorithm is known as iterative proportional fitting procedure (IPFP) or Sinkhorn algorithm [8, 20].

On the Existence of Solution to SSB



David Hilbert (from Wiki)

Under a compact support assumption, we show that this iteration is a strict contraction mapping with respect to the Hilbert metric [1].

The proof is similar to existing results for the SB problem [13, 15, 2, 9, 7]. However, the exponent $(1 + \beta)/\beta$ simplifies the argument significantly.

Time Series Extension

Time series SSB

Consider N fixed time points $0 < t_1 < \dots < t_N = 1$. Let μ_N be a probability distribution on $\mathbb{R}^{d \times N}$ such that $\mu_N \ll \lambda$. For $\beta > 0$, find $V = \inf_{u \in \mathcal{U}} J_\beta^N(u)$, where


$$J_\beta^N(u) = \beta \mathcal{D}_{\text{KL}}(\mathcal{L}\text{aw}((X_{t_i})_{1 \leq i \leq N}), \mu_N) + \mathsf{E} \int_0^1 \frac{\|u_t\|^2}{2\sigma^2} dt,$$

and find the optimal control u^* such that $J_\beta^N(u^*) = V$.

See our paper [14] for the solution.

Concluding Remarks

- Major contribution of our paper is theoretical: a rigorous solution to the SSB problem using the log transformation technique [10, 11, 12].
- Future direction: more general generative modeling algorithms based on SSB.
- Future direction: comparison between the convergence rate of IPFP for SB and that for SSB.
- There are interesting connections between SSB and the optimal transport [4]. In particular, Chen et al. [3] studied a matrix OT problem which is a discrete-time analogue to SSB on finite spaces.



Slides available at <https://zhouquan34.github.io>

Comics created with the help of ChatGPT & DALL·E 3.

Jhanvi Garg, Xianyang Zhang and Quan Zhou. "Soft-constrained Schrödinger bridge: a stochastic control approach." International Conference on Artificial Intelligence and Statistics (AISTATS 2024).

References I

- [1] Peter J Bushell. Hilbert's metric and positive contraction mappings in a Banach space. *Archive for Rational Mechanics and Analysis*, 52:330–338, 1973.
- [2] Yongxin Chen, Tryphon Georgiou, and Michele Pavon. Entropic and displacement interpolation: a computational approach using the Hilbert metric. *SIAM Journal on Applied Mathematics*, 76(6):2375–2396, 2016.
- [3] Yongxin Chen, Tryphon T Georgiou, Michele Pavon, and Allen Tannenbaum. Relaxed Schrödinger bridges and robust network routing. *IEEE Transactions on Control of Network Systems*, 7(2):923–931, 2019.
- [4] Marco Cuturi. Sinkhorn distances: Lightspeed computation of optimal transport. *Advances in neural information processing systems*, 26, 2013.
- [5] Paolo Dai Pra. A stochastic control approach to reciprocal diffusion processes. *Applied Mathematics and Optimization*, 23(1):313–329, 1991.
- [6] Valentin De Bortoli, James Thornton, Jeremy Heng, and Arnaud Doucet. Diffusion Schrödinger bridge with applications to score-based generative modeling. *Advances in Neural Information Processing Systems*, 34:17695–17709, 2021.
- [7] George Deligiannidis, Valentin De Bortoli, and Arnaud Doucet. Quantitative uniform stability of the iterative proportional fitting procedure. *The Annals of Applied Probability*, 34(1A):501–516, 2024.

References II

- [8] W Edwards Deming and Frederick F Stephan. On a least squares adjustment of a sampled frequency table when the expected marginal totals are known. *The Annals of Mathematical Statistics*, 11(4):427–444, 1940.
- [9] Montacer Essid and Michele Pavon. Traversing the Schrödinger bridge strait: Robert Fortet's marvelous proof redux. *Journal of Optimization Theory and Applications*, 181(1):23–60, 2019.
- [10] Wendell H Fleming. Exit probabilities and optimal stochastic control. *Applied Mathematics and Optimization*, 4:329–346, 1977.
- [11] Wendell H Fleming. Logarithmic transformations and stochastic control. In *Advances in Filtering and Optimal Stochastic Control: Proceedings of the IFIP-WG 7/1 Working Conference Cocoyoc, Mexico, February 1–6, 1982*, pages 131–141. Springer, 2005.
- [12] Wendell H Fleming and Raymond W Rishel. *Deterministic and stochastic optimal control*, volume 1. Springer Science & Business Media, 2012.
- [13] Robert Fortet. Résolution d'un système d'équations de m. Schrödinger. *Journal de Mathématiques Pures et Appliquées*, 19(1-4):83–105, 1940.
- [14] Jhanvi Garg, Xianyang Zhang, and Quan Zhou. Soft-constrained schrödinger bridge: a stochastic control approach. In *International Conference on Artificial Intelligence and Statistics*, pages 4429–4437. PMLR, 2024.

References III

- [15] Tryphon T Georgiou and Michele Pavon. Positive contraction mappings for classical and quantum Schrödinger systems. *Journal of Mathematical Physics*, 56(3), 2015.
- [16] Mohamed Hamdouche, Pierre Henry-Labordere, and Huyêñ Pham. Generative modeling for time series via Schrödinger bridge. *arXiv preprint arXiv:2304.05093*, 04 2023. doi: 10.13140/RG.2.2.25758.00324.
- [17] Jeremy Heng, Valentin De Bortoli, and Arnaud Doucet. Diffusion Schrödinger bridges for Bayesian computation. *Statistical Science*, 39(1):90–99, 2024.
- [18] Jonathan Ho, Ajay Jain, and Pieter Abbeel. Denoising diffusion probabilistic models. *Advances in Neural Information Processing Systems*, 33:6840–6851, 2020.
- [19] Aapo Hyvärinen. Estimation of non-normalized statistical models by score matching. *Journal of Machine Learning Research*, 6(4):695–709, 2005.
- [20] Ludger Rueschendorf. Convergence of the iterative proportional fitting procedure. *Annals of Statistics*, 23, 08 1995. doi: 10.1214/aos/1176324703.
- [21] Yang Song, Jascha Sohl-Dickstein, Diederik P. Kingma, Abhishek Kumar, Stefano Ermon, and Ben Poole. Score-based generative modeling through stochastic differential equations, 2021.
- [22] Gefei Wang, Yuling Jiao, Qian Xu, Yang Wang, and Can Yang. Deep generative learning via Schrödinger bridge. In Marina Meila and Tong Zhang, editors, *Proceedings of the 38th International Conference on Machine Learning*, volume 139 of *Proceedings of Machine Learning Research*, pages 10794–10804. PMLR, 18–24 Jul 2021.