

# ICFP M2 – SOFT MATTER PHYSICS

## Tutorial 7. Manning condensation on DNA

*Jean-Marc Di Meglio and Thomas Salez*

DNA is a polymer containing charges : a so-called polyelectrolyte. In addition to potentially altering the monomer-monomer interactions and thus the chain conformation, these charges interact with counterions from the solvent. This interaction can even lead to a massive condensation of the counterions around the chain – the latter becoming “dressed” by counterions and thus bearing a reduced effective charge. Here, we would like to understand this phenomenon using Manning’s seminal argument, and estimate further the free-counterion fraction related to the osmotic pressure. For the latter purpose, we recall the general Debye-Hückel framework for electrolyte solutions.

### I Condensation criterion

We describe the chain as an infinite and straight line, along the axis of which identical charges  $e$  are equally spaced with spatial period  $b$ . The chain is in a solvent of permittivity  $\epsilon$ .

- 1 Comment on the validity of the straight-cylinder assumption.
- 2 Give the electric field  $E$  and the potential  $\Phi$  at distance  $d$  from the axis.
- 3 We consider a monovalent counterion of charge  $-e$  placed at distance  $d$  from the axis. Calculate the interaction energy and show that the average counterionic concentration scales as  $c \sim d^{-2\xi}$ , where  $\ell_B = \xi b$  is a characteristic length – to be described.
- 4 Evaluate the number of counterions, per charge on the chain, between the axis and an arbitrary distance  $d_{\max}$ .
- 5 Show that condensation of counterions along the chain is expected for  $\xi > 1$ .
- 6 Assuming that, when  $\xi > 1$ , the condensation brings back  $\xi$  to 1 by reducing the effective chain charges, estimate the fraction of neutralized chain charges as a function of the initial  $\xi$ . How is this result modified for a counterion of charge  $ze$ ?
- 7 Estimate the condensation-induced neutralization of charges on DNA ( $b \approx 0.17$  nm) at  $37^\circ\text{C}$ .

### II Debye-Hückel theory

We consider an assembly of different ions of types  $i$ , with charges  $z_i e$ , and global average concentrations  $c_i^0$ , thermalized in a solvent at temperature  $T$ .

- 1 Express the electroneutrality constraint.
- 2 In the following, we consider a given – fixed – test ion of type  $j$  creating an electrostatic potential  $\phi(r)$  at a distance  $r$ . What is the energy of another ion, of type  $i$ , placed in this potential?
- 3 How is the local average concentration  $c_i(r)$ , of ionic species  $i$  at distance  $r$  from the test ion, modified with respect to  $c_i^0$  due to the interaction with the test ion?
- 4 What is the local charge density  $\rho(r)$  at distance  $r$  from the test ion?
- 5 In a static description, how are  $\phi$  and  $\rho$  related in a solvent of permittivity  $\epsilon$ ? Obtain the so-called Poisson-Boltzmann equation satisfied by the electric potential  $\phi$ .
- 6 Assuming a dilute solution, linearize the previous equation. What length scale  $\lambda_D$  does naturally appear?
- 7 Solve for  $\phi(r)$  and interpret the meaning of  $\lambda_D$ . One can invoke a vanishingly small ionic size  $a$  in order to discuss prefactors.

### III Free counterions and osmotic pressure

We come back to the polyelectrolyte modelled in the first section, and we consider a situation where  $\xi < 1$ . We call  $n$  the net number of charges on one chain,  $N$  the number of chains,  $V$  the volume, and we further assume  $\lambda_D$  to be much larger than the monomeric size  $a$ . Our aim is now to study the free counterions in solution in presence of the chains, by invoking the Debye-Hückel framework.

- 1 Express the total potential  $\psi(d)$  created by one chain at a given point located at distance  $d$  from the axis. Show that it depends only on the variable  $s = d/\lambda_D$ .
- 2 Provide a low- $s$  asymptotic expression for  $\psi$ , and separate it into two contributions – a chain one  $\psi_1$  and a counterionic one  $\psi_2$ .
- 3 Calculate the work  $W = n \int_0^e \psi_2 \, de'$ , needed to establish the counterionic atmosphere of the chain, by effectively loading the chain charge (integration variable :  $ne'$ ) from 0 to  $ne$ .
- 4 What is the associated energy per unit volume? How does it depend on the total polyelectrolytic charge concentration  $c_{\text{tot}}$ ?
- 5 Recall the bare osmotic pressure  $\Pi_0$  corresponding to a dilute homogeneous solution of monovalent counterions of concentration  $c_{\text{tot}}$ . Explain the latter choice of concentration.
- 6 Express the real osmotic pressure  $\Pi = \Pi_0 + \Delta\Pi$ , in presence of the chain.
- 7 Determine the osmotic coefficient  $\Pi/\Pi_0$  and interpret it as the fraction of free counterions. Discuss the case  $\xi > 1$ .