F2 Internal-bond response to complex formation

Provisional data audit | distances retain the workbook's unspecified source unit

H-end Cl-end strong-interaction candidate

(a) H-end: Δ*r*(O–H), all 28

(b) H-end: 25 other systems (zoom)

(c) Cl-end: Δ*r*(O–Cl), all 28

L1 CAACme
L2 CAACmePPh3
L3 CAACmeSPh2
L4 CO
L5 CS
L6 N2
L7 N2CAACme
L8 N2CO
L9 N2CS
L10 N2NHCme
L11 N2PPh3
L12 N2SPh2
L13 NHCme
L14 NHCmeCAACme
L15 NHCmePPh3
L16 NHCmeSPh2
L17 OCCAACme
L18 OCCS
L19 OCNHCme
L20 OCPPh3
L21 OCSPh2
L22 PPh3
L23 PPh3SPh2
L24 SCCAACme
L25 SCNHCme
L26 SCPPh3
L27 SCSPh2
L28 SPh2

0.752
0.661
0.779

in panel (a)
in panel (a)
in panel (a)

0.333

0.0 0.2 0.4 0.6 0.8
Distance change
(source unit; not specified)

0.000 0.025 0.050 0.075 0.100 0.125 0.150 0.175
Distance change
(source unit; not specified)

0.00 0.05 0.10 0.15 0.20 0.25 0.30 0.35 0.40
Distance change
(source unit; not specified)

Changes are recalculated relative to the isolated HOCl entries in Structure data!E59 (O–H) and F59 (O–Cl).
Panel (b) magnifies the 25 H-end systems outside the three highlighted H-end candidates; no points are silently discarded.
Geometry alone does not demonstrate proton transfer, chlorine transfer, or a specific bond category.