

Throughout this section, κ denotes the membrane permeability, T the target-hit time, τ_{out} the first exit time from the corridor band, and τ_{mem} the first genuine membrane crossing. The three windows **peak1**, **valley**, and **peak2** are compared using two complementary descriptions: *normalized* observables, reported as ratios or percentages of T , and *absolute* budgets, reported in expected step counts. More specifically, for an event time τ , the relative timing ratio is

$$\frac{E[\tau \mid \tau < T, T \in W]}{E[T \mid T \in W]},$$

i.e., the mean event time expressed as a fraction of the mean hit time in window W ; the outside-time share is the fraction of T spent outside the corridor band among trajectories whose hit times fall in the corresponding window; the outside budget is the expected number of pre-hit steps spent outside the corridor band; and the post-crossing budget is the expected number of steps remaining after the first membrane crossing and before target hit. In Figure 1, “outside” is the union of the left-side pocket and the merged outer/right-side bulk.

Figure 1: Top: spatial partition for the symmetric one-target baseline. Bottom: hit-time curves $f(t)$ for $\kappa = 0$ and $\kappa = 0.0040$, with embedded stacked bars for the **peak1**, **valley**, and **peak2** windows marked on the curves. The three segments give the normalized time shares in the left side, corridor, and merged outer/right side. The t_1 , t_v , and t_2 markers indicate the true window centres; the bars are shifted horizontally only for legibility.




Figure 1 provides the geometric baseline for the comparison between the valley and the second-peak window. For both $\kappa = 0$ and $\kappa = 0.0040$, the hit-time window centred on **peak2** allocates a larger fraction of its pre-hit budget outside the corridor band than the valley window does: the outside share increases from about 48% to 56% in the control and from about 47% to 56% at $\kappa = 0.0040$. The stacked bars show that this increase is carried mainly by the left-side pocket, whose share rises to about 30% in the second-peak window. This decomposition indicates that the delayed arrivals contributing to the second peak are associated primarily with increased pre-hit residence outside the corridor, especially in the left-side pocket, rather than with a uniform increase of residence in the outer/right-side bulk.

Figure 2: Relative timing ratio for τ_{out} , defined as $E[\tau_{\text{out}} \mid \tau_{\text{out}} < T, T \in W]/E[T \mid T \in W]$, that is, the mean exit time expressed as a fraction of the mean hit time in window W . Black labels give the relative timing ratios, colour-matched percentages give the outside-time shares, and the top-right annotation gives the corresponding outside budgets, namely the expected numbers of pre-hit steps spent outside the corridor band, for the valley and **peak2** windows. The three series correspond to $\kappa = 0$, $\kappa = 0.0040$, and the boundary value $\kappa = 0.0152$.




Figure 2 shows that **peak1** remains the direct-corridor benchmark across all three permeabilities: its ratio stays near 0.76 to 0.79, while the outside-time share stays near 9% to 10%. By contrast, the valley and **peak2** windows both display small exit ratios, so the normalized exit time alone does not separate them well. Their difference is seen more clearly in the outside budget. At $\kappa = 0$, the outside budget increases from about 718 steps in the valley to about 1311 steps in **peak2**; at $\kappa = 0.0040$, it increases from about 705 to about 1288 steps. Under the present parameters, these comparisons indicate that the second-peak window is not only a geometric timing effect, but is associated with longer outside excursions accumulated before absorption. In this sense, the late-time probability mass is generated by trajectories that remain outside the corridor band for substantially longer than the valley-window trajectories; Figure 1 indicates that, in this case, much of this contrast is carried by the left-side pocket.

Figure 3: Relative timing ratio for τ_{mem} , i.e., the mean first-crossing time expressed as a fraction of the mean hit time in window W . Black labels give the relative timing ratios, colour-matched percentages give the membrane-crossing probabilities $P(\tau_{\text{mem}} < T \mid T \in W)$, and the top-right annotation gives the post-crossing budgets, namely the expected numbers of steps remaining after the first crossing and before target hit. The label ‘n/a’ denotes that the ratio is undefined when no crossing occurs before target hit. The three series correspond to $\kappa = 0$, $\kappa = 0.0040$, and the boundary value $\kappa = 0.0152$.




Figure 3 isolates the membrane contribution to this picture. At $\kappa = 0$, all crossing probabilities and post-crossing budgets vanish, as expected. For $\kappa > 0$, the valley and peak2 windows remain relatively close in crossing probability, about 18% versus 20% at $\kappa = 0.0040$, so crossing probability by itself again provides only limited separation. The difference is more evident in the time that remains after the first crossing, about 213 versus 336 steps. At the boundary value $\kappa = 0.0152$, these post-crossing budgets increase further to about 604 and 785 steps. This comparison indicates that permeability acts primarily by enlarging the post-crossing budget, rather than merely by increasing the probability of crossing. The membrane therefore modifies the delayed residence mechanism mainly after crossing: it lengthens the remaining path to absorption and amplifies the late-time shoulder/second-peak contribution without making crossing probability alone a sufficient explanation.